

APPLICATION FOR UNITED STATES LETTERS PATENT
IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

(MBHB Case no. 01-019-A)

Title: BENZIMIDAZOLE AND PYRIDYLIMIDAZOLE DERIVATIVES

Inventors:

Guiying Li, #8 Valley Court, Branford, CT 06405

John M. Peterson, 28 Highland Terrace, Madison, CT 06443

Pamela Albaugh, 10475 Trebah Circle, Carmel, IN 46032

Kevin S. Currie, 33 Clear Lake Manor Road, North Branford,
CT 06471

Guolin Cai, 2382 Whitechapel Place, Thousand Oaks, CA 91362

Linda M. Gustavson, 3 Chestnut Court, Guilford, CT 06437

Kyungae Lee, 6 Finch Lane, Guilford, CT 06437

Alan Hutchison, 29 Kimberly Lane, Madison, CT 06443

Vinod Singh, 4056, IIT Campus, Kanpur - 208016, India

George D. Maynard, 27 Glenwood Road, Clinton, CT 06413

Jun Yuan, 40 Spruce Hill Drive, Guilford, CT 06437

Ling Hong Xie, 136 Renee's Way, Guilford, CT 06437

Manuka Ghosh, 211 E. Main Street, #83, Branford, CT 06405

Nian Liu, 20 Country Road, North Branford, CT 06471

George P. Luke, 17 Vienna Lane, Clinton, CT 06413

Scott Mitchell, 13 Morris Road, East Haven, CT 06513

Martin Patrick Allen, 63 Mystic Road, North Stonington CT
06359

Spiros Liras, 12 Kidds Way, Stonington CT 06378

Background of the Invention

Field of the Invention

This invention relates to benzimidazole and pyridylimidazole derivatives, and, more specifically, to such derivatives that bind with high selectively and/or high affinity to the benzodiazepine site of GABA_A receptors. This invention also relates to pharmaceutical compositions comprising such compounds and to the use of such compounds in the treatment of central nervous system (CNS) diseases.

Description of the Related Art

The GABA_A receptor superfamily represents one of the classes of receptors through which the major inhibitory neurotransmitter, γ -aminobutyric acid, or GABA, acts. Widely, although unequally, distributed throughout the mammalian brain, GABA mediates many of its actions through a complex of proteins called the GABA_A receptor, which causes alteration in chloride conductance and membrane polarization. In addition to being the site of neurotransmitter action, a number of drugs including the anxiolytic and sedating benzodiazepines bind to this receptor. The GABA_A receptor comprises a chloride channel that generally, but not invariably, opens in response to GABA, allowing chloride to enter the cell. This, in turn, effects a slowing of neuronal activity through hyperpolarization of the cell membrane potential.

GABA_A receptors are composed of five protein subunits. A number of cDNAs for these GABA_A receptor subunits have been cloned and their primary structures determined. While these subunits share a basic motif of 4 membrane-spanning helices, there is sufficient sequence diversity to classify them into several groups. To date at least 6 α , 3 β , 3 γ , 1 ϵ , 1 δ and 2 ρ subunits have been identified. Native GABA_A receptors are

typically composed of 2α , 2β , and 1γ . Various lines of evidence (such as message distribution, genome localization and biochemical study results) suggest that the major naturally occurring receptor combinations are $\alpha_1\beta_2\gamma_2$, $\alpha_2\beta_3\gamma_2$, $\alpha_3\beta_3\gamma_2$, and $\alpha_5\beta_3\gamma_2$ (Mohler et al. *Neuroch. Res.* 1995; 20(5):631-36).

The GABA_A receptor binding sites for GABA (2 per receptor complex) are formed by amino acids from the α and β subunits. Amino acids from the α and γ subunits together form one benzodiazepine site per receptor. Benzodiazepines exert their pharmacological actions by interacting with the benzodiazepine binding sites associated with the GABA_A receptor. In addition to the benzodiazepine site (sometimes referred to as the benzodiazepine or BDZ receptor), the GABA_A receptor contains sites of interaction for several other classes of drugs. These include a steroid binding site, a picrotoxin site, and a barbiturate site. The benzodiazepine site of the GABA_A receptor is a distinct site on the receptor complex that does not overlap with the site of interaction for other classes of drugs that bind to the receptor or for GABA (see, e.g., Cooper, et al., *The Biochemical Basis of Neuropharmacology*, 6th ed., 1991, pp. 145-148, Oxford University Press, New York).

In a classic allosteric mechanism, the binding of a drug to the benzodiazepine site increases the affinity of the GABA receptor for GABA. Benzodiazepines and related drugs that enhance the ability of GABA to open GABA_A receptor channels are known as agonists or partial agonists depending on the level of GABA enhancement. Other classes of drugs, such as β -carboline derivatives, that occupy the same site and negatively modulate the action of GABA are called inverse agonists. A third class of compounds exists which occupy the same site as both the agonists and inverse agonists and yet have little or no effect on GABA activity. These compounds will, however, block the action of

agonists or inverse agonists and are thus referred to as GABA_A receptor antagonists.

5 The important allosteric modulatory effects of drugs acting at the benzodiazepine site were recognized early, and the distribution of activities at different subtype receptors has been an area of intense pharmacological discovery. Agonists that act at the benzodiazepine site are known to exhibit anxiolytic, sedative, and hypnotic effects, while compounds that act as inverse agonists at this site elicit anxiogenic, cognition enhancing, and proconvulsant effects. While benzodiazepines have enjoyed long pharmaceutical use as anxiolytics, these compounds are known to exhibit a number of unwanted side effects. These may include cognitive impairment, sedation, ataxia, potentiation of ethanol effects, and a tendency for tolerance and drug dependence.

15 GABA_A selective ligands may also act to potentiate the effects of certain other CNS active compounds. For example, there is evidence that selective serotonin reuptake inhibitors (SSRIs) may show greater antidepressant activity when used in combination with GABA_A selective ligands than when used alone.

SUMMARY OF THE INVENTION

This invention provides benzimidazole and pyridylimidazole derivatives that bind to the benzodiazepine site of GABA_A receptors, including human GABA_A receptors. Preferred compounds of the invention bind with high selectivity and/or high affinity to GABA_A receptors. Preferred compounds act as agonists, antagonists or inverse agonists of such receptors. As such, they are useful in the treatment of various CNS disorders.

The invention provides compounds of Formula I (shown below), and pharmaceutical compositions comprising compounds of Formula I.


The invention provides methods for synthesizing compounds of Formula I.

The invention further provides methods of treating patients suffering from certain CNS disorders with an effective amount of a compound of the invention. The patient may be a human or other mammal. Treatment of humans, domesticated companion animals (pets) or livestock animals suffering from certain CNS disorders with an effective amount of a compound of the invention is encompassed by the invention.

In a separate aspect, the invention provides methods of potentiating the actions of other CNS active compounds. These methods comprise administering an effective amount of a compound of the invention in conjunction with the administration of another CNS active compound.

Additionally this invention relates to the use of compounds of Formula I as probes for the localization of GABA_A receptors in tissue sections.

In a first aspect, the invention provides compounds of Formula I



and the pharmaceutically acceptable salts thereof.

In this aspect, Z_1 is nitrogen or CR_1 ; Z_2 is nitrogen or CR_2 ; Z_3 is nitrogen or CR_3 ; and Z_4 is nitrogen or CR_4 ; provided that no more than two of Z_1 , Z_2 , Z_3 , and Z_4 are nitrogen.


Further in Formula I,

R_1 , R_2 , R_3 , and R_4 are independently selected from

i) hydrogen, halogen, hydroxy, nitro, cyano, amino, haloalkyl, and haloalkoxy,


ii) alkyl, alkoxy, cycloalkyl, alkenyl, alkynyl, (cycloalkyl)alkyl, $-NH(R_{10})$, $-N(R_{10})(R_{11})$, hydroxyalkyl, aminoalkyl, $(R_{10})NHalkyl-$, $(R_{10})(R_{11})Nalkyl-$, alkanoyl, alkoxycarbonyl, (heterocycloalkyl)alkyl, alkylsulfonyl, alkylthio, mono- or dialkylaminocarbonyl, heterocycloalkyl, aryl, and heteroaryl, each of which is optionally substituted with 1, 2, 3, or 4 of R_{20} , wherein R_{10} and R_{11} are independently selected from the group consisting of alkyl, alkenyl, alkynyl, alkoxy, cycloalkyl, (cycloalkyl)alkyl, aryl, arylalkyl, alkanoyl, and mono and dialkylaminoalkyl; and

iii) a group of the formula:



where G is a bond, alkyl, $-O-$, $-C(=O)-$, or $-CH_2C(=O)-$, and R_A is a saturated, partially unsaturated, or aromatic carbocycle, consisting of 1 ring or 2 fused, pendant, or spiro rings, each ring containing 0, 1, or 2 heteroatoms independently chosen from N, S, and O, said saturated, partially unsaturated, or aromatic carbocycle is optionally substituted with 1, 2, 3, or 4 of R_{20} ,

iv) a group of the formula



where J is N, CH, or C-alkyl, and R_B and R_C are independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, alkoxy, cycloalkyl, (cycloalkyl)alkyl, heterocycloalkyl, aryl, arylalkyl, alkanoyl, heteroaryl, and mono and dialkylaminoalkyl, each of which is optionally substituted by 1 or 2 substituents independently chosen from halogen, hydroxy, cyano, amino, nitro, alkoxy, and alkyl; R_B and R_C and the atom to which they are attached form a 4- to 10-membered monocyclic or bicyclic ring, which may contain:

a) one or more double bonds,

b) one or more of oxo, O, S, SO, SO₂, or N-R_D wherein R_D is hydrogen, Ar₁, alkyl, cycloalkyl, heterocycloalkyl, or Ar₁alkyl; wherein Ar₁ is aryl or heteroaryl, each of which is optionally substituted by 1 or 2 substituents independently chosen from halogen, hydroxy, cyano, amino, nitro, alkoxy, and alkyl,

c) one or more substituents R₂₀;

v) -OC(=O)R_E, -C(=O)OR_E, -C(=O)NH₂, -C(=O)NHR_E, -C(=O)NR_ER_F, -S(O)_nR_E, -S(O)_nNH₂, -S(O)_nNHR_E, -S(O)_nNR_ER_F, -NHC(=O)R_E, -C(=NR_E)R_F, -HC=N-OH, -HC=N(alkoxy), -HC=N(alkyl), -NR_EC(=O)R_F, -NHS(O)_mR_E, and -NR_ES(O)_mR_F, where m is 0, 1 or 2, and

R_E and R_F are independently selected at each occurrence from alkyl, cycloalkyl, heterocycloalkyl, alkoxy, mono- or

dialkylamino, aryl, or heteroaryl each of which is optionally substituted by 1, 2, or 3 of R₃₀.

R₂₀, in this aspect of the invention, is independently selected at each occurrence from the group consisting of: halogen; hydroxy; nitro; cyano; amino; alkyl; alkoxy optionally substituted with amino or mono- or dialkylamino; cycloalkyl; cycloalkylalkyl; cycloalkylalkoxy; alkenyl; alkynyl; haloalkyl;

oxo; haloalkoxy; mono- and dialkylamino; aminoalkyl; and mono- and dialkylaminoalkyl.

R_{30} is independently selected at each occurrence from halogen, hydroxy, nitro, cyano, amino, alkyl, alkoxy optionally substituted with amino or mono- or dialkylamino, cycloalkyl, cycloalkylalkyl, cycloalkylalkoxy, heterocycloalkyl, alkenyl, alkynyl, haloalkyl, haloalkoxy, oxo, mono- and dialkylamino, aminoalkyl, and mono- and dialkylaminoalkyl.

R_5 represents hydrogen or haloalkyl; or


R_5 represents alkyl, cycloalkyl, or (cycloalkyl)alkyl, each of which may contain one or more double or triple bonds, and each of which is optionally substituted with 1, 2, or 3 of R_{30} , or

R_5 represents aryl, arylalkyl, heteroaryl, or heteroarylalkyl each of which is optionally substituted with 1, 2, or 3 substituents selected from the group consisting of haloalkyl, amino, $-NH(R_{10})$, $-N(R_{10})(R_{11})$, carboxamido, $(R_{10})NHcarbonyl$, $(R_{10})(R_{11})Ncarbonyl$, halogen, hydroxy, nitro, cyano, amino, alkyl, alkoxy optionally substituted with amino or mono- or dialkylamino, cycloalkyl, cycloalkylalkyl, cycloalkylalkoxy, heterocycloalkyl, alkenyl, alkynyl, haloalkyl, haloalkoxy, aminoalkyl, and mono- and dialkylaminoalkyl.

Q represents $-C(R_6)(R_7)$ or oxygen, with the proviso that Q is not oxygen when X_2 is nitrogen.

R_6 and R_7 independently represent hydrogen, fluorine, or alkyl.

The group:



represents a 5 to 7 membered heteroaryl or heterocycloalkyl ring containing up to 4 heteroatoms selected from nitrogen, sulfur, and oxygen, said 5 to 7 membered heteroaryl or heterocycloalkyl

ring is substituted at each carbon atom by R, and substituted at each nitrogen atom available for substitution by R'.

R is independently chosen at each occurrence from hydrogen, halogen, amino, alkyl, alkenyl, alkynyl, alkoxy, cycloalkyl, (cycloalkyl)alkyl, haloalkyl, haloalkoxy, carboxamido, and 3- to 7-membered carbocyclic or heterocyclic groups which are saturated, unsaturated, or aromatic, which may be further substituted with one or more substituents independently selected from halogen, oxo, hydroxy, alkyl, and alkoxy.

R' is independently chosen at each occurrence from alkyl, hydrogen, cycloalkyl, cycloalkyl(alkyl), and 3- to 7-membered carbocyclic or heterocyclic groups which are saturated, unsaturated, or aromatic, which 3- to 7-membered carbocyclic or heterocyclic groups are optionally substituted with one or more substituents independently selected from halogen, oxo, hydroxy, alkyl, and alkoxy.

X₁ and X₂ independently represent nitrogen, carbon or CH.

Y is nitrogen, oxygen, carbon, -CH-, -CH₂-, or absent.

W represents aryl or heteroaryl, wherein the aryl or heteroaryl group is optionally substituted with up to 4 groups independently selected from R₃₀, -CO₂H, -C(=O)OR_E, -C(=O)NHR_E, -C(=O)NR_ER_F, -C(O)R_E, and -S(O)_mR_E, -OR_E, where R₃₀ and R_E are as defined above and m is 0, 1, or 2.

DETAILED DESCRIPTION


In addition to the compounds and salts of Formula I, described above, the invention further provides compounds of Formula I wherein

5 R_1 , R_2 , R_3 , and R_4 are independently selected from

i) hydrogen, halogen, hydroxy, nitro, cyano, amino, halo(C_1 - C_6) alkyl, and halo(C_1 - C_6)alkoxy,

10 ii) (C_1 - C_6)alkyl, (C_1 - C_6)alkoxy, (C_3 - C_8)cycloalkyl, (C_2 - C_6)alkenyl, alkynyl, ((C_3 - C_8)cycloalkyl) (C_1 - C_4)alkyl, -NH(R_{10}), -N(R_{10}) (R_{11}), hydroxy(C_1 - C_6)alkyl, amino(C_1 - C_6)alkyl, (R_{10})NH (C_1 - C_6)alkyl, (R_{10}) (R_{11})N(C_1 - C_6)alkyl, (C_1 - C_6)alkanoyl, (C_1 - C_6)alkoxycarbonyl, (C_1 - C_6)alkylsulfonyl, (C_1 - C_6)alkylthio, mono- or di(C_1 - C_6)alkylaminocarbonyl, heterocycloalkyl, (heterocycloalkyl) C_1 - C_4 alkyl, aryl, and heteroaryl, each of which
15 is optionally substituted with 1, 2, 3, or 4 of R_{20} , wherein R_{10} and R_{11} are independently selected from the group consisting of (C_1 - C_6)alkyl, (C_2 - C_6)alkenyl, (C_1 - C_6)alkoxy, (C_3 - C_8)cycloalkyl, (C_3 - C_8)cycloalkylalkyl, aryl, aryl(C_1 - C_6)alkyl, (C_1 - C_6)alkanoyl, and mono and di(C_1 - C_6)alkylaminoalkyl;


20 iii) a group of the formula:



where G is (C_1 - C_6)alkyl, -O-, -C(=O)-, or -CH₂C(=O)-, and

25 R_A is a saturated, partially unsaturated, or aromatic carbocycle, consisting of 1 ring or 2 fused, pendant, or spiro rings, each ring consisting of from 3 to 8 ring atoms, and each ring containing 0, 1, or 2 heteroatoms independently chosen from N, S, and O; said saturated, partially unsaturated, or aromatic carbocycle is optionally substituted with 1, 2, 3, or 4 of R_{20} ,

iv) a group of the formula



where J is N, CH, or C-(C₁-C₆)alkyl and R_B and R_C are independently selected from the group consisting of hydrogen, (C₁-C₆)alkyl, (C₂-C₆)alkenyl, (C₂-C₆)alkynyl, (C₁-C₆)alkoxy, (C₃-C₈)cycloalkyl, (C₃-C₈cycloalkyl) (C₁-C₄)alkyl, heterocycloalkyl, aryl, aryl(C₁-C₄)alkyl, (C₁-C₆)alkanoyl, heteroaryl, and mono and di(C₁-C₆)alkylamino(C₁-C₆)alkyl, each of which is optionally substituted by 1 or 2 substituents independently chosen from halogen, hydroxy, cyano, amino, nitro, C₁-C₆alkoxy, and C₁-C₆alkyl; or R_B and R_C and the atom to which they are attached form a 4- to 10-membered monocyclic or bicyclic ring, which may contain

a) one or more double bonds

b) one or more of oxo, O, S, SO, SO₂, and N-R_D wherein R_D is hydrogen, Ar₁, (C₁-C₆)alkyl, (C₃-C₈)cycloalkyl, heterocycloalkyl, or Ar₁(C₁-C₆)alkyl; wherein Ar₁ is aryl or heteroaryl, each of which is optionally substituted by 1 or 2 substituents independently chosen from halogen, hydroxy, cyano, amino, nitro, C₁-C₆alkoxy, and C₁-C₆alkyl;

c) one or more substituents R₂₀;

v) -OC(=O)R_E, -C(=O)OR_E, -C(=O)NH₂, -C(=O)NHR_E, -C(=O)NR_ER_F, -S(O)_nR_E, -S(O)_nNH₂, -S(O)_nNHR_E, -S(O)_nNR_ER_F, -NHC(=O)R_E, -C(=NR_E)R_F, -HC=N-OH, -HC=N(C₁-C₆alkoxy), -HC=N(C₁-C₆alkyl), -NR_EC(=O)R_F, -NHS(O)_mR_E, and -NR_ES(O)_mR_F, where m is 0, 1 or 2, and R_E and R_F are independently selected at each occurrence from (C₁-C₆)alkyl, (C₃-C₈)cycloalkyl, heterocycloalkyl, (C₁-C₆)alkoxy, mono- and di(C₁-C₆)alkylamino, aryl, and heteroaryl each of which is optionally substituted by 1, 2, or 3 of R₃₀.

R₂₀ is independently selected at each occurrence from the group consisting of halogen; hydroxy; nitro; cyano; amino; (C₁-

C₆)alkyl; (C₁-C₆)alkoxy optionally substituted with amino or mono- or di(C₁-C₆)alkylamino; (C₃-C₈)cycloalkyl; (C₃-C₈)cycloalkyl(C₁-C₄)alkyl; (C₃-C₈)cycloalkyl(C₁-C₄)alkoxy; (C₂-C₆)alkenyl; (C₂-C₆)alkynyl; halo(C₁-C₆)alkyl; halo(C₁-C₆)alkoxy; oxo; mono- and
5 di(C₁-C₆)alkylamino; amino(C₁-C₆)alkyl; and mono- and di(C₁-C₆)alkylamino(C₁-C₆)alkyl.

R₃₀ is independently selected at each occurrence from halogen, hydroxy, nitro, cyano, amino, (C₁-C₆)alkyl, (C₁-C₆)alkoxy optionally substituted with amino or mono- or di(C₁-
10 C₆)alkylamino, (C₃-C₈)cycloalkyl, (C₃-C₈)cycloalkyl(C₁-C₄)alkyl, (C₃-C₈)cycloalkyl(C₁-C₄)alkoxy, heterocycloalkyl, (C₂-C₆)alkenyl, (C₂-C₆)alkynyl, halo(C₁-C₆)alkyl, halo(C₁-C₆)alkoxy, oxo, mono- and di(C₁-C₆)alkylamino, amino(C₁-C₆)alkyl, and mono- and di(C₁-C₆)alkylamino(C₁-C₆)alkyl.


15 R₅ represents hydrogen or halo(C₁-C₆)alkyl; or

R₅ represents (C₁-C₆)alkyl, (C₃-C₈)cycloalkyl, or (C₃-C₈)cycloalkyl(C₁-C₄)alkyl, each of which may contain one or more double or triple bonds, and each of which is optionally substituted with 1, 2, or 3 of R₃₀, or

20 R₅ represents aryl, aryl(C₁-C₄)alkyl, heteroaryl, or heteroaryl(C₁-C₄)alkyl each of which is optionally substituted with 1, 2, or 3 substituents selected from the group consisting of: halo(C₁-C₆)alkyl, amino, NH(R₁₀), N(R₁₀)(R₁₁), carboxamido, NH(R₁₀)carbonyl, N(R₁₀)(R₁₁)carbonyl, halogen, hydroxy, nitro, cyano, amino, (C₁-C₆)alkyl, (C₁-C₆)alkoxy optionally substituted
25 with amino or mono- or di(C₁-C₆)alkylamino, (C₃-C₈)cycloalkyl, (C₃-C₈)cycloalkyl(C₁-C₄)alkyl, (C₃-C₈)cycloalkyl(C₁-C₄)alkoxy, heterocyclo(C₁-C₄)alkyl, (C₂-C₆)alkenyl, (C₂-C₆)alkynyl, halo(C₁-C₆)alkyl, halo(C₁-C₆)alkoxy, amino(C₁-C₆)alkyl, and mono- and
30 di(C₁-C₆)alkylamino(C₁-C₆)alkyl;

Q represents -C(R₆)(R₇) or oxygen, with the proviso that Q is not oxygen when X₂ is nitrogen;

R₆ and R₇ independently represent hydrogen, fluorine, or C₁-C₆alkyl; and
the group:



5 represents a 5 to 7 membered heteroaryl or heterocycloalkyl ring containing up to 4 heteroatoms selected from nitrogen, sulfur, and oxygen, said 5 to 7 membered heteroaryl or heterocycloalkyl ring is substituted at each carbon atom by R, and is substituted at each nitrogen atom available for substitution by R';


10 R is independently chosen at each occurrence from hydrogen, halogen, amino, C₁-C₆alkyl, (C₂-C₆)alkenyl, (C₂-C₆)alkynyl, C₁-C₆alkoxy, (C₃-C₈)cycloalkyl, (C₃-C₈cycloalkyl)(C₁-C₄)alkyl, halo(C₁-C₆)alkyl, haloalkoxy, carboxamido, and 3- to 7-membered carbocyclic or heterocyclic groups which are saturated,
15 unsaturated, or aromatic, which may be further substituted with one or more substituents independently selected from halogen, oxo, hydroxy, C₁₋₄alkyl, and -O(C₁₋₄alkyl);

R' is independently chosen at each occurrence from hydrogen, C₁-C₆alkyl, C₃-C₈cycloalkyl, C₃-C₈cycloalkyl(C₁-C₄alkyl), and 3- to
20 7-membered carbocyclic or heterocyclic groups which are saturated, unsaturated, or aromatic, which 3- to 7-membered carbocyclic or heterocyclic groups are optionally substituted with one or more substituents independently selected from halogen, oxo, hydroxy, C₁₋₄alkyl, and -O(C₁₋₄alkyl); and

25 X₁, X₂, W, and Y are as defined for Formula I, above.

Such compounds will be referred to as compounds of Formula IA.

A particular aspect of the invention is directed to
30 compounds and pharmaceutically acceptable salts of Formula II



Formula II

In Formula II, the variables Z_1 , Z_2 , Z_3 , Z_4 , R_5 , Q , X_1 , X_2 , and W carry the definition set forth for Formula I, or more preferably, for Formula IA;

X_3 and X_4 are independently selected from the group consisting of carbon, CR, N, O, S, NH, and $N(C_1-C_6)alkyl$; provided that at least one of X_1 , X_2 , X_3 , and X_4 is carbon or CR; and

R is independently chosen at each occurrence from hydrogen, halogen, amino, $(C_1-C_6)alkyl$, $(C_1-C_6)alkoxy$, $(C_3-C_8)cycloalkyl$, $(C_3-C_8)cycloalkyl(C_1-C_6)alkyl$, $(C_2-C_6)alkenyl$, $(C_2-C_6)alkynyl$, halo $(C_1-C_6)alkyl$, halo $(C_1-C_6)alkoxy$, carboxamido, and 3- to 7-membered carbocyclic or heterocyclic groups which are saturated, unsaturated, or aromatic, which may be further substituted with one or more substituents independently selected from halogen, oxo, hydroxy, $C_{1-4}alkyl$, and $-O(C_{1-4}alkyl)$.

The invention is particularly directed to compounds of Formula I, Formula IA, and Formula II, in which Z_1 is CR_1 , Z_2 is CR_2 , Z_3 is CR_3 , and Z_4 is CR_4 .

The invention is also directed to compounds of Formula I, Formula IA, and Formula II, in which one, and only one, of Z_1 , Z_2 , Z_3 , and Z_4 is nitrogen.


Another particular aspect of the invention provides compounds of Formula I, Formula IA, and Formula II, in which Z_1 is CR_1 , Z_4 is CR_4 , and only one, of Z_2 and Z_3 is nitrogen.

The invention is further directed to compounds of Formula I, Formula IA, and Formula II wherein:


- i) X_2 is carbon; and Q is oxygen;
- ii) X_2 is N; and Q is $C(R_6)(R_7)$;
- iii) X_2 is carbon; and Q is $C(R_6)(R_7)$;
- iv) X_1 is carbon; X_2 is N; and Q is $C(R_6)(R_7)$;
- v) X_1 is nitrogen; X_2 is carbon; and Q is $C(R_6)(R_7)$; or wherein
- vi) Q is $C(R_6)(R_7)$.

For each of i) through vi) preferred compounds are those where Z_1 is CR_1 , Z_2 is CR_2 , Z_3 is CR_3 , and Z_4 is CR_4 . For each of i) through vi) preferred compounds are those in which one, and only one, of Z_1 , Z_2 , Z_3 , and Z_4 is nitrogen. For each of i) through vi) compounds in which Z_1 is CR_1 , and only one, of Z_2 and Z_3 is nitrogen are particularly preferred.

In another aspect, the invention provides compounds of Formula III and Formula IV:



Formula III



Formula IV

wherein R, R_1 , R_2 , R_3 , R_4 , R_5 , Q, and W carry the definitions set forth for Formula I, or more preferably for Formula IA.

Particular compounds of Formula III included in the invention are those wherein Q is C(R₆)(R₇).

Preferred compounds of Formula IV include those where R₆ and R₇ are hydrogen, methyl or fluoro and the other is ethyl, or where one of R₆ and R₇ is hydrogen, methyl or fluoro and the other is ethyl.

Other compounds of the invention include compounds of Formula III or Formula IV, wherein R is independently selected at each occurrence from the group consisting of:

- i) hydrogen, halogen, (C₁-C₆)alkyl, (C₃-C₈)cycloalkyl, (C₃-C₈)cycloalkyl(C₁-C₆)alkyl, (C₁-C₆)alkoxy, halo(C₁-C₆)alkyl, halo(C₁-C₆)alkoxy, and
- ii) phenyl and pyridyl each of which is optionally substituted with up to 3 substituents independently chosen from halogen, hydroxy, C₁₋₄alkyl, and -O(C₁₋₄alkyl).

Q (in Formula III) is C(R₆)(R₇).

In Formula III and IV, R₁, R₂, R₃, and R₄ are independently selected from hydrogen, halogen, hydroxy, nitro, cyano, amino, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, (C₃-C₈)cycloalkyl, (C₃-C₈)cycloalkyl(C₁-C₆)alkyl, (C₂-C₆)alkenyl, (C₂-C₆)alkynyl, heterocycloalkyl, halo(C₁-C₆)alkyl, halo(C₁-C₆)alkoxy, mono or di(C₁-C₆)alkylamino, amino(C₁-C₆)alkyl, and mono- and di(C₁-C₆)alkylamino(C₁-C₆)alkyl;

R₅ represents hydrogen, (C₁-C₆)alkyl, (C₃-C₈)cycloalkyl, (C₃-C₈)cycloalkyl(C₁-C₆)alkyl, phenyl, benzyl, thiophenyl, thiazoyl, pyridyl, imidazolyl, pyrazolyl, or pyrimidinyl;

R₆ and R₇ independently represent hydrogen, fluorine, or C₁-C₆ alkyl; and

W represents phenyl, thienyl, thiazoyl, pyridyl, imidazolyl, oxazolyl, triazolyl, tetrazolyl, pyrazolyl, isoxazolyl, or pyrimidinyl, each of which is optionally substituted with up to 4 R₃₀ groups, where R₃₀ carries the definition set forth for Formula

I, or more preferably R_{30} carries the definition set forth for Formula IA.

Particularly included in the invention are compounds of
5 Formula III and Formula IV in which wherein R , R_1 , R_2 , R_3 , R_4 , R_5 , Q , and W carry the definition set forth for Formula I, or more preferably for Formula IA, and

W represents a 6-membered aryl or heteroaryl groups, wherein the 6-membered aryl or heteroaryl group is optionally substituted
10 with up to 4 groups independently selected from R_{30} , $-\text{CO}_2\text{H}$, $-\text{C}(=\text{O})\text{OR}_E$, $-\text{C}(=\text{O})\text{NHR}_E$, $-\text{C}(=\text{O})\text{NR}_E\text{R}_F$, $-\text{C}(\text{O})\text{R}_E$, $-\text{S}(\text{O})_n\text{R}_E$, and $-\text{OR}_E$ or wherein

W represents a 5-membered heteroaryl group, wherein the 5-membered heteroaryl group is optionally substituted with up to 4
15 groups independently selected from R_{30} , $-\text{CO}_2\text{H}$, $-\text{C}(=\text{O})\text{OR}_E$, $-\text{C}(=\text{O})\text{NHR}_E$, $-\text{C}(=\text{O})\text{NR}_E\text{R}_F$, $-\text{C}(\text{O})\text{R}_E$, $-\text{S}(\text{O})_m\text{R}_E$, and $-\text{OR}_E$.

In these embodiments of the invention m is 0, 1, or 2, and R_E carries the definition set forth for Formula I, or more preferably R_E carries the definition set forth for Formula IA and
20 R_{30} carries the definition set forth above with respect to Formula IA.

In another aspect, the compounds of Formula III or Formula IV are those in which one of R_2 or R_3 carries the definition set
25 forth for Formula I, or more preferably for Formula IA.

In this aspect of the invention,

R is independently selected at each occurrence from the group consisting of hydrogen, halogen, and $(\text{C}_1\text{-C}_2)\text{alkyl}$;

R_1 , R_4 , and the other of R_2 and R_3 are independently selected
30 from hydrogen, halogen, hydroxy, nitro, cyano, amino, $(\text{C}_1\text{-C}_6)\text{alkyl}$, $(\text{C}_1\text{-C}_6)\text{alkoxy}$, $(\text{C}_3\text{-C}_8)\text{cycloalkyl}$, $(\text{C}_3\text{-C}_8)\text{cycloalkyl}(\text{C}_1\text{-C}_6)\text{alkyl}$, halo $(\text{C}_1\text{-C}_6)\text{alkyl}$, halo $(\text{C}_1\text{-C}_6)\text{alkoxy}$,

mono or di(C₁-C₆)alkylamino, amino(C₁-C₆)alkyl, and mono- and di(C₁-C₆)alkylamino(C₁-C₆)alkyl;

R₅ represents (C₁-C₆)alkyl; and

Q (in Formula III) is CH₂, and R₆ and R₇ in Formula IV are hydrogen; and

W represents phenyl, furanyl, thienyl, thiazoyl, pyridyl, imidazolyl, oxazolyl, triazolyl, tetrazolyl, pyrazolyl, isoxazolyl, pyrimidinyl, benzimidazolyl, quinolinyl, isoquinolinyl each of which is optionally substituted with up to 4 R₃₀ groups, where R₃₀ carries the definition set forth for Formula I, or more preferably R₃₀ carries the definition set forth for Formula IA.

Still other preferred W groups are 4-pyrimidinyl, 5-halo-2-pyrimidinyl, 3,6-dihalopyrimidin-2-yl, and 2,6-, 4,6-, and 5,6-dihalopyridin-2-yl. Other preferred W groups are phenyl substituted with one or two independently selected C₁-C₂ alkyl, C₁-C₂ alkoxy, amino, halogen, trifluoromethyl, or cyano groups. Still other preferred W groups are 2-thiazolyl groups carrying one or two independently selected C₁-C₂ alkyl, amino, (C₁-C₃) alkyl, hydroxy, (C₁-C₃)alkyl, or trifluoromethyl groups.

Another aspect of the invention includes compounds of Formula III or Formula IV wherein

R is independently selected at each occurrence from the group consisting of hydrogen, halogen, and (C₁-C₂)alkyl;

R₁, R₄, and one of R₂ and R₃ are independently selected from hydrogen, halogen, trifluoromethyl, C₁-C₂ alkyl, and cyano;

the other of R₂ and R₃ carries the definition set forth for Formula I, or more preferably for Formula IA; and

R₅ represents (C₁-C₆)alkyl, and preferably C₂-C₄ alkyl.

Preferably R₁ and R₄ are hydrogen;

Preferred R groups are independently selected from hydrogen and C₁-C₃ alkyl, more preferably hydrogen and methyl, and most preferably are hydrogen.

More preferred R₅ groups are ethyl and n-propyl.

In this aspect,

Q (in Formula III) is CH₂, and R₆ and R₇ in Formula IV are hydrogen; and

5 W, is phenyl, pyridyl, or thiazolyl, each which is optionally substituted by one or more substituents independently chosen from halogen, cyano, hydroxy, oxo, C₁-C₂haloalkyl, C₁-C₂alkyl, and C₁-C₂ alkoxy, or more preferably

10 W is 2-thiazolyl, 2-pyrimidinyl, 3-fluorophenyl, or 6-fluoro-2-pyridinyl. Such compounds will be referred to as compounds of Formula III-A and Formula IV-A.

A particular aspect of the invention provides compounds of Formula III-A and Formula IV-A, wherein:

15 one of R₂ and R₃ is chosen from hydrogen, halogen, hydroxy, nitro, cyano, amino, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, (C₃-C₈)cycloalkyl, (C₃-C₈)cycloalkyl(C₁-C₆)alkyl, halo(C₁-C₆)alkyl, halo(C₁-C₆)alkoxy, mono or di(C₁-C₆)alkylamino, amino(C₁-C₆)alkyl, and mono- and di(C₁-C₆)alkylamino(C₁-C₆)alkyl;

20 the other of R₂ and R₃ is chosen from

i) hydrogen, halogen, hydroxy, nitro, cyano, amino, halo(C₁-C₆)alkyl, and halo(C₁-C₆)alkoxy, and

25 ii) C₁-C₆alkyl, C₁-C₆alkoxy, C₃-C₈cycloalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, (C₃-C₈cycloalkyl)C₁-C₄alkyl, -NH(R₁₀), -N(R₁₀)(R₁₁), (R₁₀)NH(C₁-C₆)alkyl, (R₁₀)(R₁₁)N(C₁-C₆)alkyl, (heterocycloalkyl)alkyl, and heterocycloalkyl, each of which is optionally substituted with 1, 2, 3, or 4 of R₂₀; and

R₂₀ carries the definition set forth for Formula I, or more

30 preferably R₂₀ carries the definition set forth with respect to Formula IA.


In this aspect, preferred compounds of Formula III-A and IV-A include those where one of R₂ and R₃ is hydrogen, halogen, hydroxy, nitro, cyano, amino, C₁-C₃ alkyl, C₁-C₂ alkoxy,

cyclopropyl, cyclopropylmethyl, trifluoromethyl, or mono- or di(C₁-C₂)alkylamino, and the other is hydrogen, halogen, or C₁-C₃ alkyl, preferably hydrogen or methyl. More preferred compounds of Formula IV-A include those where R₂ is hydrogen, halogen, more preferably fluoro or chloro, cyano, amino, C₁-C₂ alkyl or C₁-C₂ alkoxy and R₃ is hydrogen or methyl. Other more preferred compounds of Formula IV-A include those where R₂ is hydrogen, methyl, or ethyl, and R₃ is hydrogen, halogen, preferably fluoro or chloro, cyano, amino, or C₁-C₃ alkoxy.

Another aspect of the invention provides compounds of Formula III-A and Formula IV-A, wherein:

one of R₂ and R₃ is chosen from hydrogen, halogen, hydroxy, nitro, cyano, amino, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, (C₃-C₈)cycloalkyl, (C₃-C₈)cycloalkyl(C₁-C₆)alkyl, halo(C₁-C₆)alkyl, halo(C₁-C₆)alkoxy, mono or di(C₁-C₆)alkylamino, amino(C₁-C₆)alkyl, and mono- and di(C₁-C₆)alkylamino(C₁-C₆)alkyl;

the other of R₂ and R₃ is chosen from



where J is N, CH, or C-(C₁-C₆)alkyl and

R_B and R_C are independently selected from the group consisting of hydrogen, (C₁-C₆)alkyl, (C₂-C₆)alkenyl, (C₂-C₆)alkynyl, C₃-C₈)cycloalkyl, and (C₃-C₈cycloalkyl) (C₁-C₄)alkyl; or

R_B and R_C and the atom to which they are attached form a 4- to 10-membered monocyclic or bicyclic ring, which may contain


- one or more double bonds,
- one or more of oxo, O, S, SO, SO₂, and N-R_D wherein R_D is hydrogen or (C₁-C₆)alkyl;
- one or more of R₂₀; and

R₂₀ carries the definition set forth with respect to Formula I, or more preferably R₂₀ carries the definition set forth for R₂₀ Formula IA.

5 The invention also provides compounds of Formula III-A and Formula IV-A, wherein:

one of R₂ and R₃ is chosen from hydrogen, halogen, hydroxy, nitro, cyano, amino, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, (C₃-C₈)cycloalkyl, (C₃-C₈)cycloalkyl(C₁-C₆)alkyl, halo(C₁-C₆)alkyl, halo(C₁-C₆)alkoxy, mono or di(C₁-C₆)alkylamino, amino(C₁-C₆)alkyl, and mono- and di(C₁-C₆)alkylamino(C₁-C₆)alkyl;

the other of R₂ and R₃ is a group of the formula:



where G is a bond or C₁-C₂alkyl;

15 R_A is a saturated, partially unsaturated, or aromatic carbocycle, consisting of 1 ring or 2 fused, pendant, or spiro rings, each ring containing 0, 1, or 2 heteroatoms independently chosen from N, S, and O, said saturated, partially unsaturated, or aromatic carbocycle is optionally substituted with 1, 2, 3, or 4 of R₂₀; and

20 R₂₀ carries the definition set forth for R_a in Formula I, or more preferably R₂₀ carries the definition set forth for R₂₀ in Formula IA R₂₀.

25 Preferably R_A is chosen from phenyl, pyrrolyl, pyrazolyl, thiazolyl, isoxazolyl, triazolyl, tetrazolyl, oxadiazolyl, and oxazolyl each of which is optionally substituted with 1, 2, 3, or 4 of R₂₀.

30 Other compounds of the invention are compounds of Formula III-A and Formula IV-A, wherein:

one of R_2 and R_3 is chosen from hydrogen, halogen, hydroxy, nitro, cyano, amino, (C_1-C_6) alkyl, (C_1-C_6) alkoxy, (C_3-C_8) cycloalkyl, (C_3-C_8) cycloalkyl (C_1-C_6) alkyl, halo (C_1-C_6) alkyl, halo (C_1-C_6) alkoxy, mono or di (C_1-C_6) alkylamino, amino (C_1-C_6) alkyl, and mono- and di (C_1-C_6) alkylamino (C_1-C_6) alkyl; and the other of R_2 and R_3 is $-HC=N-OH$ or $-HC=N(C_1-C_6\text{alkoxy})$.

Another aspect of the invention is directed to compounds of Formula I, Formula IA and Formula II wherein


one and only one of Z_1 is CR_1 , Z_4 is CR_4 , either Z_2 or Z_3 is nitrogen; and

i) W represents a 5-membered heteroaryl group, and the 5-membered heteroaryl group is optionally substituted with up to 4 groups independently selected from R_{30} , $-CO_2H$, $-C(=O)OR_E$, $-C(=O)NHR_E$, $-C(=O)NR_ER_F$, $-C(O)R_E$, and $-S(O)_mR_E$, $-OR_E$; or


ii) W represents a 6-membered aryl or heteroaryl group, wherein the 6-membered aryl or heteroaryl group is optionally substituted with up to 4 groups independently selected from R_{30} , $-CO_2H$, $-C(=O)OR_E$, $-C(=O)NHR_E$, $-C(=O)NR_ER_F$, $-C(O)R_E$, and $-S(O)_mR_E$, $-OR_E$; where

R_{30} and R_E carry the definitions set forth for those groups in Formula I, or preferably for Formula IA, and m is 0, 1, or 2.

Thus, the invention includes compounds represented by Formula V and Formula VI



Formula V



Formula VI

wherein R, R₁, R₂, R₃, R₄, R₅, and Q carry the definitions set forth for Formula I, or more preferably for Formula IA, and W is a 5-membered heteroaryl group as described above.

5 The invention further includes compounds of Formula V and Formula VI wherein

R is independently selected at each occurrence from the group consisting of

- 10 i) hydrogen, halogen, (C₁-C₆)alkyl, (C₃-C₈)cycloalkyl, (C₃-C₈)cycloalkyl(C₁-C₆)alkyl, (C₁-C₆)alkoxy, halo(C₁-C₆)alkyl, halo(C₁-C₆)alkoxy, and
- ii) phenyl and pyridyl each of which is optionally substituted with up to 3 substituents independently chosen from halogen, hydroxy, C₁₋₄alkyl, and -O(C₁₋₄alkyl);

15 R₁, R₂, R₃, and R₄, are independently selected from hydrogen, halogen, hydroxy, nitro, cyano, amino, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, (C₃-C₈)cycloalkyl, (C₃-C₈)cycloalkyl(C₁-C₆)alkyl, (C₂-C₆)alkenyl, (C₂-C₆)alkynyl, heterocycloalkyl, halo(C₁-C₆)alkyl, halo(C₁-C₆)alkoxy, mono or di(C₁-C₆)alkylamino, amino(C₁-C₆)alkyl, and mono- and di(C₁-C₆)alkylamino(C₁-C₆)alkyl;

20 R₅ represents hydrogen, (C₁-C₆)alkyl, (C₃-C₈)cycloalkyl, (C₃-C₈)cycloalkyl(C₁-C₆)alkyl, phenyl, benzyl, thiophenyl, thiazoyl, pyridyl, imidazolyl, pyrazolyl, or pyrimidinyl;

25 R₆ and R₇ independently represent hydrogen, fluorine, or C₁-C₆ alkyl;

W represents either a 5-membered heteroaryl group chosen from thienyl, thiazolyl, imidazolyl, oxazolyl, triazolyl, tetrazolyl, pyrazolyl, or isoxazolyl each of which is optionally substituted with up to 4 R₃₀ groups; or

30 W represents a 6-membered aryl or heteroaryl group chosen from phenyl, pyrimidinyl, pyridyl, pyridizinyl, or pyrazinyl, each of which is optionally substituted with up to 4 R₃₀ groups; and

R₃₀ is as defined for Formula I, or preferably as defined for Formula IA.

Preferred compounds of Formula V and VI include those where R₂ and R₃ independently represent hydrogen, halogen, preferably fluoro or chloro, C₁-C₃ alkyl, cyclopropyl, cyclopropylmethyl, C₁-C₃ alkoxy, trifluoromethyl, nitro, cyano, amino, or mono- or di(C₁-C₃) alkylamino. Other preferred R₂ and R₃ groups are mono- or di(C₁-C₃) alkylamino(C₂-C₃) alkoxy, morpholinyl(C₂-C₃) alkoxy, piperidin-1-yl(C₂-C₃) alkoxy, and piperazin-1-yl(C₂-C₃) alkoxy.

In another aspect, the invention is directed to compounds of Formula V and Formula VI wherein:

R is independently selected at each occurrence from the group consisting of hydrogen, halogen, and (C₁-C₂) alkyl; and

R₁ and R₄ are independently selected from hydrogen, halogen, hydroxy, nitro, cyano, amino, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₃-C₈) cycloalkyl, (C₃-C₈) cycloalkyl(C₁-C₆) alkyl, halo(C₁-C₆) alkyl, halo(C₁-C₆) alkoxy, mono or di(C₁-C₆) alkylamino, amino(C₁-C₆) alkyl, and mono- and di(C₁-C₆) alkylamino(C₁-C₆) alkyl.

In this aspect of the invention,

R₂ (Formula V) and R₃ (Formula VI) carry the definitions set forth with respect to Formula I, or more preferably for Formula IA;

R₅ represents (C₁-C₆) alkyl, preferably ethyl or n-propyl;

R₆ and R₇ are hydrogen;

W represents a 5-membered heteroaryl group chosen from furanyl, thienyl, thiazoyl, imidazolyl, oxazolyl, triazolyl, tetrazolyl, pyrazolyl, or isoxazolyl, each of which is optionally substituted with up to 4 R₃₀ groups, or

W represents a 6-membered aryl or heteroaryl group chosen from phenyl, pyrimidinyl, pyridyl, pyridiziny, or pyrazinyl each of which is optionally substituted with up to 4 R₃₀ groups; and

R₃₀ is as defined for Formula I, or more preferably for Formula IA.

Compounds of this aspect of the invention will be referred to as compounds of Formula V-A and Formula VI-A.

5 Preferred R groups are hydrogen and C₁-C₃ alkyl, more preferably hydrogen and methyl, and most preferably hydrogen.

Preferred R₁ and R₄ groups for this aspect of the invention include hydrogen, halogen, trifluoromethyl, C₁-C₂ alkyl, and cyano. Preferably R₁ and R₄ are hydrogen. More preferably R, R₁,
10 and R₄ are all hydrogen.

Preferred compounds of this aspect of the invention in which W is a 5-membered heteroaryl group include those wherein W is thiazolyl which is optionally substituted by one or more
15 substituents independently chosen from halogen, cyano, hydroxy, oxo, C₁-C₂haloalkyl, C₁-C₂alkyl, and C₁-C₂ alkoxy. Other preferred compounds are those wherein W is 2-thiazolyl.

Preferred compounds of this aspect in which W is a 6-membered heteroaryl group include those wherein W is phenyl or pyridyl, each of which is optionally substituted by one or more
20 substituents independently chosen from halogen, cyano, hydroxy, oxo, C₁-C₂haloalkyl, C₁-C₂alkyl, and C₁-C₂ alkoxy. Also preferred are compounds wherein W is 2-pyrimidinyl, 3-fluorophenyl, or 6-fluoro-2-pyridinyl.

Still other preferred W groups are 4-pyrimidinyl, 5-halo-2-pyrimidinyl, 3,6-dihalopyrimidin-2-yl, and 2,6-, 4,6-, and 5,6-dihalopyridin-2-yl. Other preferred W groups are phenyl
25 substituted with one or two independently selected C₁-C₂ alkyl, C₁-C₂ alkoxy, amino, halogen, trifluoromethyl, or cyano groups. Still other preferred W groups are 2-thiazolyl groups carrying
30 one or two independently selected C₁-C₂ alkyl, amino, (C₁-C₃) alkyl, hydroxy, (C₁-C₃)alkyl, or trifluoromethyl groups.

Other aspects of the invention include compounds of Formula V-A and Formula VI-A wherein


R₂ (for Formula V-A) or R₃ (for Formula VI-A) is chosen from

i) hydrogen, halogen, hydroxy, nitro, cyano, amino, halo(C₁-C₆)alkyl, and halo(C₁-C₆)alkoxy,

ii) C₁-C₆alkyl, C₁-C₆alkoxy, C₃-C₈cycloalkyl, C₂-C₆alkenyl, C₁-C₆alkynyl, (C₃-C₈cycloalkyl)C₁-C₄alkyl, -NH(R₁₀), -N(R₁₀)(R₁₁), (R₁₀)NH(C₁-C₆)alkyl, (R₁₀)(R₁₁)N(C₁-C₆)alkyl, (heterocycloalkyl)C₁-C₄alkyl, and heterocycloalkyl, each of which is optionally substituted with 1, 2, 3, or 4 of R₂₀, where R₂₀ carries the definition set forth for compounds of Formula I, or preferably for compounds of Formula IA.

Another aspect of the invention is directed to compounds of Formula V-A and Formula VI-A wherein R₂ (for Formula V-A) or R₃ (for Formula VI-A) is chosen from hydrogen, halogen, hydroxy, nitro, cyano, amino, halo(C₁-C₆)alkyl, and halo(C₁-C₆)alkoxy.

Further aspects of the invention include compounds of Formula V-A and Formula VI-A wherein R₂ (for Formula V-A) or R₃ (for Formula VI-A) is a group of the formula




where J is N, CH, or C-(C₁-C₆)alkyl and

R_B and R_C are independently selected from the group consisting of hydrogen, (C₁-C₆)alkyl, (C₂-C₆)alkenyl, (C₂-C₆)alkynyl, C₃-C₈cycloalkyl, and (C₃-C₈cycloalkyl)(C₁-C₄)alkyl; or

R_B and R_C and the atom to which they are attached form a 4- to 10-membered monocyclic or bicyclic ring, which may contain a) one or more double bonds, b) one or more of oxo, O, S, SO, SO₂, and N-R_D wherein R_D is hydrogen or (C₁-C₆)alkyl; and/or c) one or

more substituents R_{20} , where R_{20} carries the definition set forth for compounds of Formula I, or more preferably that set forth for compounds of Formula IA.


- 5 Also provided by the invention are compounds of Formula V-A and Formula VI-A wherein R_2 (for Formula V-A) or R_3 (for Formula VI-A) is a group of the formula




- 10 where G is a bond or C_1 - C_2 alkyl; and R_A is a saturated, partially unsaturated, or aromatic carbocycle consisting of 1 ring or 2 fused, pendant, or spiro rings, each ring containing 0, 1, or 2 heteroatoms independently selected from N, S, and O, where the saturated, partially unsaturated, or aromatic carbocycle is optionally substituted with 1, 2, 3, or 4 of R_{20} . Preferred
15 compounds of this class are those where R_A is chosen from phenyl, pyrrolyl, pyrazolyl, thiazolyl, isoxazolyl, triazolyl, tetrazolyl, oxadiazolyl, and oxazolyl each of which is optionally substituted with 1, 2, 3, or 4 of R_{20} . R_{20} carries the definition set forth for compounds of Formula I, or more preferably that set
20 forth for compounds of Formula IA.

The invention also includes compounds of Formula V-A and Formula VI-A wherein R_2 (for Formula V-A) or R_3 (for Formula VI-A) is $-HC=N-OH$ or $-HC=N(C_1-C_6\text{alkoxy})$.


25 Other benzimidazole and pyridylimidazole compounds of the invention are represented by Formula X - Formula XVIII, below.




Formula X




Formula XI




Formula XII



Formula XIII




Formula XIV




where R' is H or C₁-C₆alkyl


Formula XV



Formula XVI



Formula XVII



Formula XVIII

The variables Z₁, Z₂, Z₃, Z₄, R, Q, and W, which appear in Formulae X - XVIII carry the definition set forth for Formula I, or more preferably for Formula IA.

- 5 Compounds of Formulae X - XVIII wherein Z₁ is CR₁, Z₂ is CR₂, Z₃ is CR₃, and Z₄ is CR₄ are preferred. Compounds of Formulae X - XVIII wherein one and only one Z₁, Z₂, Z₃, Z₄ is nitrogen are also preferred. Compounds of Formulae X - XVIII wherein one and only one Z₁, Z₂, Z₃, Z₄ is nitrogen, and either Z₂ or Z₃ is nitrogen are
- 10 particularly preferred.

Particularly embodied in the invention are compounds of Formulae X - XVIII wherein Q (when present) is $C(R_6)(R_7)$. Preferably R_6 and R_7 are hydrogen.

5 Other embodiments of the invention are directed to compounds of Formulae X - XVIII wherein W represents a 5-membered heteroaryl group, and the 5-membered heteroaryl group is optionally substituted with up to 4 groups independently selected from R_{30} , $-CO_2H$, $-C(=O)OR_E$, $-C(=O)NHR_E$, $-C(=O)NR_ER_F$, $-C(O)R_E$, and -
10 $S(O)_mR_E$, $-OR_E$, where R_{30} and R_E are as defined above and m is 0, 1, or 2. Preferred compounds of this class are compounds wherein wherein Z_1 is CR_1 , Z_2 is CR_2 , Z_3 is CR_3 , and Z_4 is CR_4 or wherein one and only one Z_1 , Z_2 , Z_3 , Z_4 is nitrogen are also preferred; compounds of this class wherein one and only one Z_1 , Z_2 , Z_3 , Z_4 is
15 nitrogen, and either Z_2 or Z_3 is nitrogen are particularly preferred.

This invention provides benzimidazole and pyridylimidazole derivatives, preferred examples of which bind with high affinity
20 to the benzodiazepine site of $GABA_A$ receptors, including human $GABA_A$ receptors. The affinity of compounds of Formula I for the benzodiazepine site may be determined using a $GABA_A$ receptor binding assay, such as the assay presented in Example 53. Preferred compounds of Formula I that bind with high affinity to
25 the benzodiazepine site of the $GABA_A$ receptor exhibit K_i values of less than $1\mu M$ in that assay. Very high affinity compounds of the invention exhibit K_i values of less than 100 nM or more preferably less than 10 nM in the assay presented in Example 53. Without wishing to be bound to any particular theory, it is
30 believed that the interaction of the compounds of Formula I with the benzodiazepine site results in the pharmaceutical utility of these compounds.

Syndrome, neurodegenerative diseases such as Alzheimer's disease and Parkinson's disease, and stroke related dementia. Compounds of the invention that act as inverse agonists at the $\alpha_5\beta_3\gamma_2$ are particularly useful in treating cognitive disorders through the enhancement of memory, and particularly short-term memory, in memory-impaired patients. Compounds of the invention that act as agonists at the $\alpha_1\beta_2\gamma_2$ receptor subtype are useful in treating convulsive disorders such as epilepsy. Compounds that act as antagonists at the benzodiazepine site are useful in reversing the effect of benzodiazepine overdose and in treating drug and alcohol addiction.

The diseases and/ or disorders that can also be treated using compounds and compositions according to the invention include:

Depression, e.g. depression, atypical depression, bipolar disorder, depressed phase of bipolar disorder.

Anxiety, e.g. general anxiety disorder (GAD), agoraphobia, panic disorder +/- agoraphobia, social phobia, specific phobia, Post traumatic stress disorder, obsessive compulsive disorder (OCD), dysthymia, adjustment disorders with disturbance of mood and anxiety, separation anxiety disorder, anticipatory anxiety acute stress disorder, adjustment disorders, cyclothymia.

Sleep disorders, e.g. sleep disorders including primary insomnia, circadian rhythm sleep disorder, dyssomnia NOS, parasomnias, including nightmare disorder, sleep terror disorder, sleep disorders secondary to depression and/or anxiety or other mental disorders, substance induced sleep disorder.

Cognition Impairment, e.g. cognition impairment, memory impairment, short-term memory impairment, Alzheimer's disease, Parkinson's disease, mild cognitive impairment (MCI), age-related cognitive decline (ARCD), stroke, traumatic brain injury, AIDS associated dementia, and dementia associated with depression, anxiety or psychosis.

Attention Deficit Disorder, e.g. attention deficit disorder (ADD), and attention deficit and hyperactivity disorder (ADHD).

Speech disorders, e.g. stuttering, including motor tic, clonic stuttering, dysfluency, speech blockage, dysarthria, Tourette syndrome or logospasm.

Psychosis e.g. schizophrenia, hallucinatory disorders

The invention also provides pharmaceutical compositions comprising one or more compounds of the invention together with a pharmaceutically acceptable carrier or excipient, for treating disorders responsive to GABA_A receptor modulation, e.g., treatment of anxiety, depression, sleep disorders or cognitive impairment by GABA_A receptor modulation. Pharmaceutical compositions include packaged pharmaceutical compositions comprising a container holding a therapeutically effective amount of at least one GABA_A receptor modulator as described supra and instructions (e.g., labeling) indicating the contained GABA_A receptor ligand is to be used for treating a disorder responsive to GABA_A receptor modulation in the patient.

In a separate aspect, the invention provides a method of potentiating the actions of other CNS active compounds, which comprises administering an effective amount of a compound of the invention in combination with another CNS active compound. Such CNS active compounds include, but are not limited to the following: for anxiety, serotonin receptor (e.g. 5-HT_{1A}) agonists and antagonists; for anxiety and depression, neurokinin receptor antagonists or corticotropin releasing factor receptor (CRF₁) antagonists; for sleep disorders, melatonin receptor agonists; and for neurodegenerative disorders, such as Alzheimer's dementia, nicotinic agonists, muscarinic agents, acetylcholinesterase inhibitors and dopamine receptor agonists. Particularly the invention provides a method of potentiating the antidepressant activity of selective serotonin reuptake

inhibitors (SSRIs) by administering an effective amount of a GABA agonist compound of the invention in combination with an SSRI.

Combination administration can be carried out in a fashion analogous to that disclosed in Da-Rocha, et al., *J. Psychopharmacology* (1997) 11(3) 211-218; Smith, et al., *Am. J. Psychiatry* (1998) 155(10) 1339-45; or Le, et al., *Alcohol and Alcoholism* (1996) 31 Suppl. 127-132. Also see, the discussion of the use of the GABA_A receptor ligand 3-(5-methylisoxazol-3-yl)-6-(1-methyl-1,2,3-triazol-4-yl) methoxy-1,2,4-triazolo [3,4-
10 alphthalzine in combination with nicotinic agonists, muscarinic agonists, and acetylcholinesterase inhibitors, in PCT International publications Nos. WO 99/47142, WO 99/47171, and WO 99/47131, respectively. Also see in this regard PCT International publication No. WO 99/37303 for its discussion of
15 the use of a class of GABA_A receptor ligands, 1,2,4-triazolo[4,3-b]pyridazines, in combination with SSRIs.

The present invention also pertains to methods of inhibiting the binding of benzodiazepine compounds, such as Ro15-1788, or GABA to the GABA_A receptors which methods involve contacting a
20 solution containing compound of the invention with cells expressing GABA_A receptors, wherein the compound is present at a concentration sufficient to inhibit benzodiazepine binding or GABA binding to GABA_A receptors *in vitro*. This method includes inhibiting the binding of benzodiazepine compounds to GABA_A
25 receptors *in vivo*, e.g., in a patient given an amount of a compound of Formula I that would be sufficient to inhibit the binding of benzodiazepine compounds or GABA to GABA_A receptors *in vitro*. In one embodiment, such methods are useful in treating benzodiazepine drug overdose. The amount of a compound that would
30 be sufficient to inhibit the binding of a benzodiazepine compound to the GABA_A receptor may be readily determined via a GABA_A receptor binding assay, such as the assay described in Example 53. The GABA_A receptors used to determine *in vitro* binding may be

obtained from a variety of sources, for example from preparations of rat cortex or from cells expressing cloned human GABA_A receptors.

10033069 "110111
The invention also provides methods for altering the signal-transducing activity, particularly the chloride ion conductance of GABA_A receptors, said method comprising exposing cells expressing such receptors to an effective amount of a compound of the invention. This method includes altering the signal-transducing activity of GABA_A receptors *in vivo*, e.g., in a patient given an amount of a compound of Formula I that would be sufficient to alter the signal-transducing activity of GABA_A receptors *in vitro*. The amount of a compound that would be sufficient to alter the signal-transducing activity of GABA_A receptors may be determined via a GABA_A receptor signal transduction assay, such as the assay described in Example 54. The cells expressing the GABA receptors *in vivo* may be, but are not limited to, neuronal cells or brain cells. Such cells may be contacted with compounds of the invention through contact with a body fluid containing the compound, for example through contact with cerebrospinal fluid. Alteration of the signal-transducing activity of GABA_A receptors *in vitro* may be determined from a detectable change in the electrophysiology of cells expressing GABA_A receptors, when such cells are contacted with a compound of the invention in the presence of GABA. For example, a change in the electrophysiology of cells expressing GABA_A receptors may be detected using a voltage-clamp assay performed on oocytes injected with GABA_A receptor mRNA. Such an assay is shown in Example 54.

Intracellular recording or patch-clamp recording may be used to quantitate changes in electrophysiology of cells. A reproducible change in behavior of an animal given a compound of the invention may also be used to indicate that changes in the

electrophysiology of the animal's cells expressing GABA_A receptors has occurred.

The GABA_A receptor ligands provided by this invention and labeled derivatives thereof are also useful as standards and reagents in determining the ability of a potential pharmaceutical to bind to the GABA_A receptor. Radiolabeled derivatives the GABA_A receptor ligands provided by this invention are also useful as radiotracers for positron emission tomography (PET) imaging or for single photon emission computerized tomography (SPECT).


More particularly compounds of the invention may be used for demonstrating the presence of GABA_A receptors in cell or tissue samples. This may be done by preparing a plurality of matched cell or tissue samples, at least one of which is prepared as an experimental sample and at least one of which is prepared as a control sample. The experimental sample is prepared by contacting (under conditions that permit binding of R015-1788 to GABA_A receptors within cell and tissue samples) at least one of the matched cell or tissue samples that has not previously been contacted with any compound or salt of the invention with an experimental solution comprising the detectably-labeled preparation of the selected compound or salt at the first measured molar concentration. The control sample is prepared in the same manner as the experimental sample and also contains an unlabelled preparation of the same compound or salt of the invention at a greater molar concentration.

The experimental and control samples are then washed to remove unbound detectably-labeled compound. The amount of remaining bound detectably-labeled compound is then measured and the amount of detectably-labeled compound in the experimental and control samples is compared. A comparison that indicates the detection of a greater amount of detectable label in the at least one washed experimental sample than is detected in any of control

samples demonstrates the presence of GABA_A receptors in that experimental sample.


The detectably-labeled compound used in this procedure may be labeled with a radioactive label or a directly or indirectly luminescent label. When tissue sections are used in this procedure and the detectably-labeled compound is radiolabeled, the bound, labeled compound may be detected autoradiographically to generate an autoradiogram. The amount of detectable label in an experimental or control sample may be measured by viewing the autoradiograms and comparing the exposure density of the autoradiograms.

The invention provides a method for preparing a compound of Formula A




Formula A

which comprises reacting a compound of Formula B



Formula B

with a compound of Formula C



Formula C.

In Formula A and C, above Z_1 , Z_2 , Z_3 , Z_4 , and R_5 carry the definitions forth for Formula I, or more preferably Z_1 , Z_2 , Z_3 , Z_4 , and R_5 carry the definitions forth for Formula Ia.

R_6 and R_7 independently represent hydrogen, fluorine, or alkyl.

R in Formula B, above, is independently chosen at each occurrence from hydrogen, halogen, amino, C_1 - C_6 alkyl, (C_2 - C_6)alkenyl, (C_2 - C_6)alkynyl, C_1 - C_6 alkoxy, (C_3 - C_8)cycloalkyl, (C_3 - C_8 cycloalkyl) (C_1 - C_4)alkyl, halo(C_1 - C_6)alkyl, haloalkoxy, carboxamido, and 3- to 7-membered carbocyclic or heterocyclic groups which are saturated, unsaturated, or aromatic, which may be further substituted with one or more substituents independently selected from halogen, oxo, hydroxy, C_{1-4} alkyl, and $-O(C_{1-4}alkyl)$.

W in Formula B represents aryl or heteroaryl, wherein the aryl or heteroaryl group is optionally substituted with up to 4 groups independently selected from R_{30} , $-CO_2H$, $-C(=O)OR_E$, $-C(=O)NHR_E$, $-C(=O)NR_ER_F$, $-C(O)R_E$, and $-S(O)_mR_E$, $-OR_E$, where R_{30} and R_E are as for Formula I or preferably as defined for Formula Ia and m is 0, 1, or 2. This process will be referred to as Process 1.

In particular embodiments the invention includes a process of preparing a compound of Formula A as described above wherein: Z_1 is CR_1 , Z_2 is CR_2 , Z_3 is CR_3 , and Z_4 is CR_4 .

R is independently selected at each occurrence from the group consisting of hydrogen, halogen, and (C_1 - C_2)alkyl;

R_1 , R_4 , and one of R_2 and R_3 are independently selected from hydrogen, halogen, hydroxy, nitro, cyano, amino, (C_1 - C_6)alkyl, (C_1 - C_6)alkoxy, (C_3 - C_8)cycloalkyl, (C_3 - C_8)cycloalkyl(C_1 - C_6)alkyl,

halo(C₁-C₆)alkyl, halo(C₁-C₆)alkoxy, mono or di(C₁-C₆)alkylamino, amino(C₁-C₆)alkyl, and mono- and di(C₁-C₆)alkylamino(C₁-C₆)alkyl.

The other of R₂ and R₃ carries the definition set forth for Formula I, or preferably that set forth for Formula Ia, or in

certain preferred embodiments this group is chosen from

i) hydrogen, halogen, hydroxy, nitro, cyano, amino, halo(C₁-C₆)alkyl, and halo(C₁-C₆)alkoxy,

ii) C₁-C₆alkyl, C₁-C₆alkoxy, C₃-C₈cycloalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, (C₃-C₈cycloalkyl) C₁-C₄alkyl, -NH(R₁₀), -

N(R₁₀)(R₁₁), (R₁₀)NH(C₁-C₆)alkyl, (R₁₀)(R₁₁)N(C₁-C₆)alkyl, (heterocycloalkyl)C₁-C₄alkyl, and heterocycloalkyl, each of which is optionally substituted with 1, 2, 3, or 4 of R₂₀.

In certain preferred embodiments R, R₁, and R₄ are all hydrogen.

R₅ represents (C₁-C₆)alkyl. Preferred definitions of R₅ include ethyl and n-propyl.

R₆ and R₇ are hydrogen.

W represents phenyl, furanyl, thienyl, thiazolyl, imidazolyl, oxazolyl, triazolyl, tetrazolyl, pyrazolyl, isoxazolyl, pyrimidinyl, benzimidazolyl, quinolinyl, isoquinolinyl each of which is optionally substituted with up to 4 R₃₀ groups, where R₃₀ is as defined in the above process. Preferred W groups include, 2-thiazolyl, 2-pyrimidinyl, 3-fluorophenyl, or 6-fluoro-2-pyridinyl.

In other preferred embodiments the invention is directed to a process, as described as for Process 1, wherein Z₁ is CR₁; one and only one of Z₂ or Z₃ is nitrogen; Z₄ is CR₄.

R₁ and R₄ may carry the definition set forth in Process 1.

Preferred definitions of R₁ and R₄ include hydrogen, halogen, trifluoromethyl, C₁-C₂ alkyl, and cyano.

In certain preferred embodiments R, R₁, and R₄ are all hydrogen.

R₂ or R₃ (for whichever one of Z₂ or Z₃ is CR₂ or CR₃) is chosen from

i) hydrogen, halogen, hydroxy, nitro, cyano, amino, halo(C₁-C₆)alkyl, and halo(C₁-C₆)alkoxy,

5 ii) C₁-C₆alkyl, C₁-C₆alkoxy, C₃-C₈cycloalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, (C₃-C₈cycloalkyl) C₁-C₄alkyl, -NH(R₁₀), -N(R₁₀)(R₁₁), (R₁₀)NH(C₁-C₆)alkyl, (R₁₀)(R₁₁)N(C₁-C₆)alkyl, (heterocycloalkyl)C₁-C₄alkyl, and heterocycloalkyl, each of which is optionally substituted with 1, 2, 3, or 4 of R₂₀.

10 R₅ represents (C₁-C₆)alkyl. Preferred definitions of R₅ include ethyl and n-propyl.

R₆ and R₇ are hydrogen.

W represents a 5-membered heteroaryl group, the 5-membered heteroaryl group is optionally substituted with up to 4 groups
15 independently selected from R₃₀, -CO₂H, -C(=O)OR_E, -C(=O)NHR_E, -C(=O)NR_ER_F, -C(O)R_E, and -S(O)_mR_E, -OR_E, where R₃₀ and R_E are as defined above and m is 0, 1, or 2; or

W represents a 6-membered aryl or heteroaryl group, wherein the 6-membered aryl or heteroaryl group is optionally substituted
20 with up to 4 groups independently selected from R₃₀, -CO₂H, -C(=O)OR_E, -C(=O)NHR_E, -C(=O)NR_ER_F, -C(O)R_E, and -S(O)_mR_E, -OR_E, where R₃₀ and R_E are as defined above and m is 0, 1, or 2.

When W represents a 5-membered heteroaryl group W is preferably thiazolyl, thienyl, imidazolyl, oxazolyl, triazolyl,
25 tetrazolyl, pyrazolyl, or isoxazolyl, each of which is optionally substituted by one or more substituents independently chosen from halogen, cyano, hydroxy, oxo, C₁-C₂haloalkyl, C₁-C₂alkyl, and C₁-C₂alkoxy. Unsubstituted 2-thiazolyl is a particularly preferred W group.

30 When W represents a 6-membered aryl or heteroaryl group, W is preferably phenyl, pyrimidinyl, pyridyl, pyrazinyl, or pyridizynyl, each of which is optionally substituted by one or more substituents independently chosen from halogen, cyano,

hydroxy, oxo, C₁-C₂haloalkyl, C₁-C₂alkyl, and C₁-C₂ alkoxy. Particularly preferred W groups include 2-pyrimidinyl, 3-fluorophenyl, or 6-fluoro-2-pyridinyl.

In this process the reactants B and C are generally combined in a polar aprotic solvent, such as THF, DMF, or 1,4-dioxane, at temperatures ranging from 0 - 100 degrees C. A reducing agent such as NaH or other base, for example sodium hydroxide, potassium butoxide, potassium carbonate, or cesium carbonate, is then added, and the reaction is allowed to proceed. Choice of solvent, reaction temperature, and reducing agent will depend on the identity of the reactants B and C, but will be readily determined by a worker of ordinary skill in the art of chemical synthesis. Scheme I, step 4, provides further illustration of this process.

Chemical description and terminology

Formula I includes, but is not limited to the subformulae exemplified as Formula Ia, Formulae II-VI and Formulae X - XVIII and their pharmaceutically acceptable acid and base addition salts.

The phrase "pharmaceutically acceptable" is employed herein to refer to those compounds, materials, compositions, and/or dosage forms which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of human beings or animals without excessive toxicity, irritation, allergic response, or other problem or complication, commensurate with a reasonable benefit/risk ratio. As used herein, "pharmaceutically acceptable salts" refer to derivatives of the disclosed compounds wherein the parent compound is modified by making acid or base salts thereof. Those skilled in the art will recognize a wide variety of non-toxic pharmaceutically acceptable addition salts.

Examples of pharmaceutically acceptable salts include, but are not limited to, mineral or organic acid salts of basic residues such as amines; alkali or organic salts of acidic residues such as carboxylic acids; and the like. The pharmaceutically acceptable salts include the conventional non-toxic salts or the quaternary ammonium salts of the parent compound formed, for example, from non-toxic inorganic or organic acids. For example, such conventional non-toxic salts include those derived from inorganic acids such as hydrochloric, hydrobromic, hydroiodic, sulfuric, sulfamic, sulfinic, phosphoric, nitric and the like; and the salts prepared from organic acids such as alkanolic such as acetic, $\text{HOOC}-(\text{CH}_2)_n\text{-ACOOH}$ where n is 0-4, and the like, tartaric, maleic, propionic, succinic, glycolic, stearic, lactic, malic, tartaric, citric, ascorbic, pantoic, maleic, hydroxymaleic, phenylacetic, glutamic, benzoic, salicylic, sulfanilic, 2-acetoxybenzoic, fumaric, toluenesulfonic, methanesulfonic, ethane disulfonic, oxalic, isethionic, $\text{HOOC}-(\text{CH}_2)_n\text{-COOH}$ where n is 0-4, and the like. The pharmaceutically acceptable salts of the present invention can be synthesized from the parent compound which contains a basic or acidic moiety by conventional chemical methods. Generally, such salts can be prepared by reacting the free acid or base forms of these compounds with a stoichiometric amount of the appropriate base or acid in water or in an organic solvent, or in a mixture of the two; generally, nonaqueous media like ether, ethyl acetate, ethanol, isopropanol, or acetonitrile are preferred. Lists of suitable salts are found in *Remington's Pharmaceutical Sciences*, 17th ed., Mack Publishing Company, Easton, PA, p. 1418 (1985).

The invention includes hydrates of compounds of Formula I.

The invention includes all crystalline forms of the compounds of Formula I. Certain crystalline forms may be preferred.

The present invention also encompasses the acylated prodrugs of the compounds of Formula I. Those skilled in the art will recognize various synthetic methodologies that may be employed to prepare non-toxic pharmaceutically acceptable addition salts and acylated prodrugs of the compounds encompassed by Formula I. The invention further encompasses all enantiomers and diastereomers of the disclosed compounds. Those of ordinary skill in the art will readily recognize methods by which mixtures of enantiomers and diastereomers may be resolved. The definition of Formula I as used in herein include possible isomers, such as tautomers and rotamers.

The compounds herein described may have one or more asymmetric centers or planes. Compounds of the present invention containing an asymmetrically substituted atom may be isolated in optically active or racemic forms. It is well known in the art how to prepare optically active forms, such as by resolution of racemic forms (racemates), by asymmetric synthesis, or by synthesis from optically active starting materials. Resolution of the racemates can be accomplished, for example, by conventional methods such as crystallization in the presence of a resolving agent, or chromatography, using, for example a chiral HPLC column. Many geometric isomers of olefins, C=N double bonds, and the like can also be present in the compounds described herein, and all such stable isomers are contemplated in the present invention. *Cis* and *trans* geometric isomers of the compounds of the present invention are described and may be isolated as a mixture of isomers or as separated isomeric forms. All chiral (enantiomeric and diastereomeric), and racemic forms, as well as all geometric isomeric forms of a structure are intended, unless the specific stereochemistry or isomeric form is specifically indicated.

When any variable occurs more than one time in any constituent or formula for a compound, its definition at each

occurrence is independent of its definition at every other occurrence. Thus, for example, if a group is shown to be substituted with 0 to 3 R*, (where R* indicates any variable group such as R) then said group may optionally be substituted
5 with up to three R* groups and R* at each occurrence is selected independently from the definition of R*. Also, combinations of substituents and/or variables are permissible only if such combinations result in stable compounds.

When any group, such as an aryl group, heteroaryl group,
10 carbocyclic group, heterocyclic group, or monocyclic or bicyclic ring is said to be "optionally substituted by one or more substituents" that group may contain 0 or from 1 to the maximum number of substituents allowable without exceeding the valency of the atoms of the substituted group. Preferably such groups are
15 substituted with 0 or from 1 to 4 substituents, and more preferably such groups are substituted with 0 or from 1 to 3 substituents. Preferably such groups are not substituted with more than one oxo substituent.

A dash "-" that is not between two letters or symbols is
20 used to indicate a point of attachment for a substituent. For example -C(=O)NH₂ is attached through the carbon atom.

As used herein, "alkyl" is intended to include both branched and straight-chain aliphatic hydrocarbon groups, having the specified number of carbon atoms. Alkyl groups of 2 or more
25 carbon atoms may contain double or triple bonds. Examples of alkyl include, but are not limited to, methyl, ethyl, n-propyl, i-propyl, n-butyl, s-butyl, t-butyl, n-pentyl, and s-pentyl. Preferred alkyl groups are C₁-C₆ alkyl groups. "C₁-C₆ alkyl" indicates alkyl groups having from 1 to about 6 carbon atoms.

30 As used herein, "alkoxy" represents an alkyl group as defined above with the indicated number of carbon atoms attached through an oxygen bridge. Examples of alkoxy include, but are not limited to, methoxy, ethoxy, n-propoxy, i-propoxy, n-butoxy, 2-

butoxy, t-butoxy, n-pentoxy, 2-pentoxy, 3-pentoxy, isopentoxy, neopentoxy, n-hexoxy, 2-hexoxy, 3-hexoxy, and 3-methylpentoxy. "C₁-C₆ alkoxy" indicates alkoxy groups having from 1 to about 6 carbon atoms.

5 "Alkenyl" is intended to include hydrocarbon chains of either a straight or branched configuration comprising one or more unsaturated carbon-carbon bonds which may occur in any stable point along the chain, such as ethenyl and propenyl. Alkenyl groups typically will have 2 to about 8 carbon atoms,
10 more typically 2 to about 6 carbon atoms.

"Alkynyl" is intended to include hydrocarbon chains of either a straight or branched configuration comprising one or more triple carbon-carbon bonds which may occur in any stable point along the chain, such as ethynyl and propynyl. Alkynyl
15 groups typically will have 2 to about 8 carbon atoms, more typically 2 to about 6 carbon atoms.

"Aryl" refers to aromatic groups having 1 or more rings, wherein the members of the aromatic ring or rings are carbon. When indicated such groups may be substituted. Preferred aryl
20 groups include optionally substituted phenyl and optionally substituted naphthyl.

The term "Cycloalkyl" is intended to include saturated ring groups, having the specified number of carbon atoms, such as cyclopropyl, cyclobutyl, cyclopentyl, or cyclohexyl. Cycloalkyl
25 groups typically will have 3 to about 8 ring members.

In the term "(cycloalkyl)alkyl", Cycloalkyl and alkyl are as defined above and the point of attachment is on the alkyl group. This term encompasses, but is not limited to, cyclopropylmethyl, cyclohexylmethyl, cyclohexylmethyl.

30 As used herein, "haloalkyl" is intended to include both branched and straight-chain saturated aliphatic hydrocarbon groups having the specified number of carbon atoms, substituted with 1 or more halogen atoms (for example -C_vF_w where v = 1 to 3

and $w = 1$ to $(2v+1)$. Examples of haloalkyl include, but are not limited to, trifluoromethyl, difluoromethyl, trichloromethyl, pentafluoroethyl, and pentachloroethyl.

As used herein, "haloalkoxy" indicates a haloalkyl group as defined above with the indicated number of carbon atoms attached through an oxygen bridge. Examples of haloalkoxy groups include, but are not limited to, trifluoromethoxy and trichloromethoxy.

As used herein the term "heteroaryl" is intended to mean a stable 5-to 7-membered monocyclic or 7-to 10-membered bicyclic heterocyclic aromatic ring which consists of carbon atoms and from 1 to 4 heteroatoms independently selected from the group consisting of N, O and S. It is preferred that the total number of S and O atoms in the heteroaryl group is not more than 1.

Examples of heteroaryl groups include, but are not limited to, pyrimidinyl, pyridyl, quinolinyl, benzothienyl, indolyl, pyridazinyl, pyrazinyl, isoindolyl, isoquinolyl, quinazolinyl, quinoxalinyl, phthalazinyl, imidazolyl, isoxazolyl, pyrazolyl, oxazolyl, thienyl, thiazolyl, indolizynyl, indazolyl, benzothiazolyl, benzimidazolyl, benzofuranyl, benzoisoxolyl, dihydro-benzodioxinyl, furanyl, pyrrolyl, oxadiazolyl, thiadiazolyl, triazolyl, tetrazolyl, oxazolopyridinyl, imidazopyridinyl, isothiazolyl, naphthyridinyl, cinnolinyl, carbazolyl, beta-carbolinyl, isochromanlyl, chromanonyl, chromanyl, tetrahydroisoquinolinyl, isoindolinyl, isobenzotetrahydrofuranlyl, isobenzotetrahydrothienyl, isobenzothienyl, benzoxazolyl, pyridopyridinyl, benzotetrahydrofuranlyl, benzotetrahydrothienyl, purinyl, benzodioxolyl, triazinyl, phenoxazinyl, phenothiazinyl, pteridinyl, benzothiazolyl, imidazopyridinyl, imidazothiazolyl, dihydrobenzisoxazinyl, benzisoxazinyl, benzoxazinyl, dihydrobenzisothiazinyl, benzopyranyl, benzothiopyranyl, coumarinyl, isocoumarinyl, chromanyl, tetrahydroquinolinyl, dihydroquinolinyl, dihydroquinolinonyl, dihydroisoquinolinonyl,

1
2
3
4 dihydrocoumarinyl, dihydroisocoumarinyl, isoindolinonyl,
5 benzodioxanyl, benzoxazolinonyl, pyrrolyl N-oxide, pyrimidinyl N-
oxide, pyridazinyl N-oxide, pyrazinyl N-oxide, quinolinyl N-
oxide, indolyl N-oxide, indolinyl N-oxide, isoquinolyl N-oxide,
6 quinazolinyl N-oxide, quinoxalinyl N-oxide, phthalazinyl N-oxide,
7 imidazolyl N-oxide, isoxazolyl N-oxide, oxazolyl N-oxide,
8 thiazolyl N-oxide, indoliziny N-oxide, indazolyl N-oxide,
9 benzothiazolyl N-oxide, benzimidazolyl N-oxide, pyrrolyl N-oxide,
10 oxadiazolyl N-oxide, thiadiazolyl N-oxide, triazolyl N-oxide,
11 tetrazolyl N-oxide, benzothiopyranyl S-oxide, and
12 benzothiopyranyl S,S-dioxide.

13 Preferred heteroaryl groups include imidazolyl, pyrrolyl,
14 pyridyl, thiazolyl, pyrazolyl, thiazolyl, isoxazolyl, triazolyl,
15 tetrazolyl, oxadiazolyl, pyrimidinyl, and oxazolyl.

16 The term "heterocycloalkyl" is intended to include saturated
17 ring groups having at least 1 heteroatom. Heterocycloalkyl
18 groups typically include 3 to 8 ring atoms, preferably 5 to 7
19 ring atoms. Heterocycloalkyl groups typically have from 1 to 3
20 heteroatoms selected from N, S, and O with remaining ring atoms
21 being carbon. Preferably not more than one S atom and one O atom
22 is present in a heterocycloalkyl group. Preferred
23 heterocycloalkyl groups include morpholinyl, piperidinyl,
24 piperazinyl, thiomorpholinyl, and pyrrolidinyl.

25 The phrase "monocyclic or bicyclic ring" refers to
26 saturated, partially unsaturated, or aromatic rings or ring
27 systems, which optionally contain from 1 to 4 heteroatoms
28 independently chosen from N, S, and O with remaining ring members
29 being carbon. Preferred monocyclic and bicyclic rings are
30 saturated and partially unsaturated rings or ring systems.

31 The term "oxo" indicates a carbonyl group. When an oxo
32 group appears as a substituent the allowed valence of the
33 substituted position is not exceeded.

Pharmaceutical Compositions

5 The compounds of general Formulas I may be administered orally, topically, parenterally, by inhalation or spray or rectally in dosage unit formulations containing conventional non-toxic pharmaceutically acceptable carriers, adjuvants and vehicles. Oral administration in the form of a pill, capsule, elixir, syrup, lozenge, troche, or the like is particularly preferred. The term parenteral as used herein includes subcutaneous injections, intradermal, intravascular (e.g., 10 intravenous), intramuscular, spinal, intrathecal injection or like injection or infusion techniques. In addition, there is provided a pharmaceutical formulation comprising a compound of general Formula I and a pharmaceutically acceptable carrier. One or more compounds of general Formula I may be present in association with one or more non-toxic pharmaceutically acceptable carriers and/or diluents and/or adjuvants and if desired other active ingredients. The pharmaceutical compositions containing compounds of general Formula I may be in a form suitable for oral use, for example, as tablets, troches, 20 lozenges, aqueous or oily suspensions, dispersible powders or granules, emulsion, hard or soft capsules, or syrups or elixirs.

25 Compositions intended for oral use may be prepared according to any method known to the art for the manufacture of pharmaceutical compositions and such compositions may contain one or more agents selected from the group consisting of sweetening agents, flavoring agents, coloring agents and preserving agents in order to provide pharmaceutically elegant and palatable preparations. Tablets contain the active ingredient in admixture with non-toxic pharmaceutically acceptable excipients that are 30 suitable for the manufacture of tablets. These excipients may be for example, inert diluents, such as calcium carbonate, sodium carbonate, lactose, calcium phosphate or sodium phosphate; granulating and disintegrating agents, for example, corn starch,

or alginic acid; binding agents, for example starch, gelatin or acacia, and lubricating agents, for example magnesium stearate, stearic acid or talc. The tablets may be uncoated or they may be coated by known techniques to delay disintegration and absorption in the gastrointestinal tract and thereby provide a sustained action over a longer period. For example, a time delay material such as glyceryl monostearate or glyceryl distearate may be employed.

Formulations for oral use may also be presented as hard gelatin capsules wherein the active ingredient is mixed with an inert solid diluent, for example, calcium carbonate, calcium phosphate or kaolin, or as soft gelatin capsules wherein the active ingredient is mixed with water or an oil medium, for example peanut oil, liquid paraffin or olive oil.

Aqueous suspensions contain the active materials in admixture with excipients suitable for the manufacture of aqueous suspensions. Such excipients are suspending agents, for example sodium carboxymethylcellulose, methylcellulose, hydropropylmethylcellulose, sodium alginate, polyvinylpyrrolidone, gum tragacanth and gum acacia; dispersing or wetting agents may be a naturally-occurring phosphatide, for example, lecithin, or condensation products of an alkylene oxide with fatty acids, for example polyoxyethylene stearate, or condensation products of ethylene oxide with long chain aliphatic alcohols, for example heptadecaethyleneoxycetanol, or condensation products of ethylene oxide with partial esters derived from fatty acids and a hexitol such as polyoxyethylene sorbitol monooleate, or condensation products of ethylene oxide with partial esters derived from fatty acids and hexitol anhydrides, for example polyethylene sorbitan monooleate. The aqueous suspensions may also contain one or more preservatives, for example ethyl, or n-propyl p-hydroxybenzoate, one or more

coloring agents, one or more flavoring agents, and one or more sweetening agents, such as sucrose or saccharin.

Oily suspensions may be formulated by suspending the active ingredients in a vegetable oil, for example arachis oil, olive oil, sesame oil or coconut oil, or in a mineral oil such as liquid paraffin. The oily suspensions may contain a thickening agent, for example beeswax, hard paraffin or cetyl alcohol. Sweetening agents such as those set forth above, and flavoring agents may be added to provide palatable oral preparations. These compositions may be preserved by the addition of an anti-oxidant such as ascorbic acid.

Dispersible powders and granules suitable for preparation of an aqueous suspension by the addition of water provide the active ingredient in admixture with a dispersing or wetting agent, suspending agent and one or more preservatives. Suitable dispersing or wetting agents and suspending agents are exemplified by those already mentioned above. Additional excipients, for example sweetening, flavoring and coloring agents, may also be present.

Pharmaceutical compositions of the invention may also be in the form of oil-in-water emulsions. The oily phase may be a vegetable oil, for example olive oil or arachis oil, or a mineral oil, for example liquid paraffin or mixtures of these. Suitable emulsifying agents may be naturally-occurring gums, for example gum acacia or gum tragacanth, naturally-occurring phosphatides, for example soy bean, lecithin, and esters or partial esters derived from fatty acids and hexitol, anhydrides, for example sorbitan monoleate, and condensation products of the said partial esters with ethylene oxide, for example polyoxyethylene sorbitan monoleate. The emulsions may also contain sweetening and flavoring agents.

Syrups and elixirs may be formulated with sweetening agents, for example glycerol, propylene glycol, sorbitol or sucrose.

Such formulations may also contain a demulcent, a preservative and flavoring and coloring agents. The pharmaceutical compositions may be in the form of a sterile injectable aqueous or oleaginous suspension. This suspension may be formulated according to the known art using those suitable dispersing or wetting agents and suspending agents which have been mentioned above. The sterile injectable preparation may also be sterile injectable solution or suspension in a non-toxic parentally acceptable diluent or solvent, for example as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil may be employed including synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid find use in the preparation of injectables.

The compounds of general Formulas I may also be administered in the form of suppositories, e.g., for rectal administration of the drug. These compositions can be prepared by mixing the drug with a suitable non-irritating excipient that is solid at ordinary temperatures but liquid at the rectal temperature and will therefore melt in the rectum to release the drug. Such materials are cocoa butter and polyethylene glycols.

Compounds of general Formulas I may be administered parenterally in a sterile medium. The drug, depending on the vehicle and concentration used, can either be suspended or dissolved in the vehicle. Advantageously, adjuvants such as local anesthetics, preservatives and buffering agents can be dissolved in the vehicle.

For administration to non-human animals, the composition may also be added to the animal feed or drinking water. It will be convenient to formulate these animal feed and drinking water compositions so that the animal takes in an appropriate quantity

of the composition along with its diet. It will also be convenient to present the composition as a premix for addition to the feed or drinking water.

Dosage levels of the order of from about 0.1 mg to about 140 mg per kilogram of body weight per day are useful in the treatment of the above-indicated conditions (about 0.5 mg to about 7 g per patient per day). The amount of active ingredient that may be combined with the carrier materials to produce a single dosage form will vary depending upon the host treated and the particular mode of administration. Dosage unit forms will generally contain between from about 1 mg to about 500 mg of an active ingredient.

Frequency of dosage may also vary depending on the compound used and the particular disease treated. However, for treatment of most disorders, a dosage regimen of 4 times daily or less is preferred. For the treatment of anxiety, depression, or cognitive impairment a dosage regimen of 1 or 2 times daily is particularly preferred. For the treatment of sleep disorders a single dose that rapidly reaches effective concentrations is desirable.

It will be understood, however, that the specific dose level for any particular patient will depend upon a variety of factors including the activity of the specific compound employed, the age, body weight, general health, sex, diet, time of administration, route of administration, and rate of excretion, drug combination and the severity of the particular disease undergoing therapy.

Preferred compounds of the invention will have certain pharmacological properties. Such properties include, but are not limited to high solubility (preferably 500 ng/ ml or more) in aqueous solutions, oral bioavailability, low toxicity, low serum protein binding, lack of clinically relevant EKG effects, and desirable *in vitro* and *in vivo* half-lives. Penetration of the

blood brain barrier for compounds used to treat CNS disorders is necessary, while low brain levels of compounds used to treat peripherreal disorders are often preferred.

Assays may be used to predict these desirable pharmacological properties. Assays used to predict bioavailability include transport across human intestinal cell monolayers, including Caco-2 cell monolayers. Toxicity to cultured hepatocytes may be used to predict compound toxicity. Penetration of the blood brain barrier of a compound in humans may be predicted from the brain levels of the compound in laboratory animals given the compound intravenously.

Serum protein binding may be predicted from albumin binding assays. Such assays are described in a review by Oravcová, et al. (Journal of Chromatography B (1996) volume 677, pages 1-27).

Compound half-life is inversely proportional to the frequency of dosage of a compound. *In vitro* half-lives of compounds may be predicted from assays of microsomal half-life as described by Kuhnz and Gieschen (Drug Metabolism and Disposition, (1998) volume 26, pages 1120-1127).


EXAMPLES

Preparation of Compounds

Representative procedures suitable for the preparation of compounds of Formula I are outlined in Schemes I-X, which are not to be construed as limiting the invention in scope or spirit to the specific reagents and conditions shown in them. Those having skill in the art will recognize that the reagents and conditions may be varied and additional steps employed to produce compounds encompassed by the present invention. In some cases, protection of reactive functionalities may be necessary to achieve the desired transformations. In general, such need for protecting


groups, as well as the conditions necessary to attach and remove such groups, will be apparent to those skilled in the art of organic synthesis. Unless otherwise stated in the schemes below, the variables, e.g., Z₁, Z₂, Z₃, Z₄, R₅, R₂, R₃ and W, are as defined in Formula I.

Scheme I




Scheme I illustrates a route to selected compounds of Formula 6 via coupling of chloromethyl compounds 4 and aryl imidazoles 5. In Step 1, aryl and heteroaryl halides of formula 1 are reacted with appropriate amines in the presence of base to obtain amino adducts of formula 2. In Step 2, reduction of the nitro group in compounds of formula 2 yields diamines 3. In Step 3, diamines of formula 3 are reacted with 2-chloroacetimidic acid methyl ester hydrochloride or a similar electrophile such as 2-chloro-1,1,1-trimethoxy-ethane or chloroacetic acid anhydride. In Step 4, chloromethyl compounds of formula 4 are reacted with aryl and heteroaryl imidazoles of formula 5 in the presence of base and solvent to obtain compounds of formula 6. Depending on the particular nature of 5, a stronger or weaker base may be selected to facilitate the reaction in Step 4.

Scheme II



Scheme II illustrates the synthesis of compounds of formula 10 from diamines 3. In Step 1, reaction of malonyl dichloride with ethyl vinyl ether provides 7. In Step 2, treatment of 7 with triethyl orthoformate in the presence of acid yields 8. Compound 8 is reacted in Step 3 with a variety of aryl and heteroaryl hydrazines to obtain compounds of formula 9 as a mixture with the undesired regioisomer. As illustrated in Steps 4 and 5, compounds of formula 9 can be hydrolyzed to the corresponding acids and coupled with compounds of formula 3 to obtain, following cyclization in refluxing acetic acid, compounds of formula 10. As described in subsequent schemes and examples, compounds of formula 9 may also be directly coupled to compounds of formula 3 in the presence of trimethylaluminum. Depending on the particular example and reaction conditions selected, cyclization may occur without need for heating in acetic acid as described in Step 5.


Scheme III



Scheme III illustrates a method for preparing compounds of formula 21 and 22. Step 1 encompasses hydrolysis of compounds of formula 11 to the corresponding acids followed by dimerization in the presence of a suitable coupling reagent such as 1,1'-carbonyldiimidazole to form compounds of formula 12. In Step 2, deacylation of compounds of formula 12 is accomplished by heating with concentrated sulfuric acid to obtain compounds of formula 13. Heating of compounds of formula 13 with ammonium hydroxide in Step 3 results in formation of compounds of formula 14. Nitration of compounds of formula 14 in Step 4 is accomplished using nitric acid to obtain compounds of formula 15. Compounds of formula 15 are converted to the corresponding chlorides 16 in Step 5 by heating with phosphorous oxychloride. In Step 6, chlorides 16 are reacted with ammonia followed by heating with phosphorous oxychloride to obtain 2-chloropyridines 17, which are

subsequently reduced to diamines **18** in Step 7. In Step 8, diamines **18** are reacted with esters of formula **19** in the presence of trimethylaluminum followed by heating in acetic acid to obtain compounds of formula **20**. Depending on the particular example and reaction conditions selected, cyclization may occur without need for heating in acetic acid. Step 9 illustrates alkylation of compounds of formula **20** with ethyl iodide in the presence of base to obtain a mixture of compounds of formula **21** and **22**. Those skilled in the art will realize that alternate alkylating agents may be employed to obtain similar compounds bearing different R_5 groups.


Scheme IV



Scheme IV illustrates a variation of Scheme III for specifically preparing compounds of formula **22**. In Step 1, compounds of formula **16** are reacted with ethyl amine to form the amino adducts which are subsequently converted to 2-chloropyridines **23** by reaction with phosphorous oxychloride. Those skilled in the art will realize that numerous other suitable amines of formula $R_5\text{NH}_2$ may be employed in Step 1 to yield other variants of Formula I. In Steps 2 and 3, hydrogenation of compounds of formula **23** to diamines of formula **24** followed by trimethylaluminum-facilitated coupling and


cyclization in acetic acid provides compounds of formula 22. Depending on the particular example and reaction conditions selected, cyclization may occur without need for heating in acetic acid

Scheme V




Scheme V illustrates a route employing a protecting group strategy for preparing pyrazole compounds of formula 29. In Step 1, pyrazole 25 is reacted with di-tert-butyl dicarbonate in the presence of 4-dimethylaminopyridine to obtain 26. Reaction with glyoxal and ammonium hydroxide provides 27. Reaction of 27 with chloromethyl compounds of formula 4 in the presence of base provides compounds of formula 28. Deprotection of compounds of formula 28 with acid in Step 4 provides pyrazoles of formula 29.

Scheme VI



Scheme VI provides a route for preparing thiazole compounds of formula 33. Step 1 involves bromination of ketoesters of formula 30 to form α -bromoketones of formula 31. In Step 2, compounds of formula 31 are reacted with thioformamide to obtain thiazoles of formula 32. Condensation of 32 with 3 in Step 3 in the presence of trimethylaluminum provides compounds of formula 33. Depending on the particular reactants and conditions employed in Step 3, the product mixture may require heating in a suitable solvent such as acetic acid to enhance formation of 33.


Scheme VII



Scheme VII provides routes to several heterocyclic systems via common intermediate 35. In Step 1, compounds of formula 34 are reacted with compounds of formula 4 at low temperature in the presence of a suitable base such as lithium diisopropylamide to form compounds of formula 35. Rxn 1 illustrates conversion of ketones of formula 35 to isoxazole derivatives of formula 36 by reaction of compounds of formula 35 with tris(dimethylamino)methane followed by treatment with hydroxylamine. In Rxn 2, compounds of formula 35 are reacted with tris(dimethylamino)methane followed by treatment with hydrazine acetate to obtain pyrazoles 37. In Rxn 3 and 4, compounds of formula 35 are brominated to form α -bromoketones 38 that are subsequently reacted with thioformamide to obtain thiazoles of


formula 39. Rxn 5 illustrates the synthesis of pyrazoles of formulas 40 and 41 by reaction of compounds of formula 35 with tris(dimethylamino)methane followed by treatment with methylhydrazine.

Scheme VIII



Scheme VIII provides a route for preparation of tetrazoles of formula 44. In Step 1, aryl and heteroaryl tetrazoles of formula 42 are heated with bis(tributyltin) oxide to form stannanes of formula 43. In Step 2, heating compounds of formula 43 with compounds of formula 4 in a suitable solvent such as toluene gives compounds of formula 44.


Scheme IX



Scheme IX illustrates the synthesis of triazoles of formula 50. In Step 1, aryl and heteroaryl hydrazines are reacted with

1,3,5-triazine **45** to obtain triazoles of formula **46**. In Step 2, heating compounds of formula **46** with formaldehyde provides alcohols of formula **47**. In Step 3, alcohols of formula **47** are converted to the corresponding chlorides by treatment with thionyl chloride. The chlorides are subsequently converted to nitriles **48** by the action of tetraethyl ammonium cyanide. Cyanides **48** are hydrolyzed in Step 4 to carboxylic acids **49**. In Step 5, carboxylic acids of formula **49** are coupled with diamines **3** in the presence of EDCI [1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride] or other suitable coupling reagents followed by heating in acetic acid to complete cyclization of the intermediate amino amides to compounds of formula **50**.

Scheme X



Scheme X illustrates two routes for the synthesis of imidazoles of formula **52**, which are intermediates in the synthesis of selected compounds of Formula I. In Step 1, aryl and heteroaryl aldehydes are treated with glyoxal and ammonium hydroxide to form imidazoles of formula **52**. In Step 1', imidazole **53** is treated with butyl lithium followed by tri-*n*-butyltin chloride to obtain compounds of formula **54**, which must be handled with care to avoid decomposition. In Step 2',

compounds of formula 54 are utilized in palladium cross-coupling reactions with aryl and heteroaryl halides to obtain compounds of formula 55. Subsequent treatment of compounds of formula 55 with acid in Step 3' provides compounds of formula 52.

5


The invention is illustrated further by the following examples, which are not to be construed as limiting the invention in scope or spirit to the specific procedures described in them. Additional compounds encompassed by this invention beyond the accompanying Examples may be prepared using methods known to those skilled in the art of chemical synthesis. Those having skill in the art will recognize that the starting materials may be varied and additional steps employed to produce compounds encompassed by the present inventions, as demonstrated by the following examples. In some cases, protection of reactive functionalities may be necessary to achieve some the desired transformations. In general, such need for protecting groups, as well as the conditions necessary to attach and remove such groups, will be apparent to those skilled in the art of organic synthesis.

Preparation of starting materials and intermediates

The starting materials and various intermediates may be obtained from commercial sources, prepared from commercially available organic compounds, or prepared using known synthetic methods. Representative examples of methods suitable for preparing intermediates of the invention are set forth below.

Example 1

Synthesis of 1-Propyl-2-([2-(2-fluoropyrid-6-yl)-1H-imidazol-1-yl]methyl)-5-cyano-1H-benzimidazole



1. Preparation of 4-n-Propylamino-3-nitrobenzonitrile

To a stirring suspension of 4-chloro-3-nitrobenzonitrile (7.30g, 40 mmol) in isopropanol (30 mL) is added n-propylamine (9.87 mL, 120 mmol). The mixture is stirred at room temperature for 5 h, and the solid is then collected by filtration to give 4-n-propylamino-3-nitrobenzonitrile as a yellow solid. ¹H NMR (CDCl₃) δ 8.51 (1H, q), 8.42(1H, br s), 7.60(1H, m), 6.91(1H, d), 3.33(2H, q), 1.84-1.74(2H, m), 1.07(3H, t). LRMS 206.3 (MH⁺).

2. Preparation of 3-Amino-4-n-propylaminobenzonitrile

To a Parr bottle containing 4-n-propylamino-3-nitrobenzonitrile (7.63 g, 37.2 mmol) in ethyl acetate (38 mL) is added 5% Pd/C (50% wet, 633 mg). The Parr bottle is sealed in a mechanical shaker, evacuated, and then purged with nitrogen followed by hydrogen. The system is pressurized to 50 PSI of hydrogen at room temperature and mechanical shaking engaged. After 2 hours, shaking is stopped, and the system purged with nitrogen prior to opening the vessel. The reaction mixture is filtered through celite, concentrated in vacuo, and the obtained solid recrystallized by dissolving in ethyl acetate (15 mL), heating, and adding hexanes (15 mL), to give 3-Amino-4-n-propylaminobenzonitrile as gray crystals. ¹H NMR (CDCl₃) δ 7.14(1H, dd), 6.92(1H, d), 6.56(1H, d), 3.98(1H, br s), 3.30(2H, br s), 3.12(2H, t), 1.75-1.65(2H, m), 1.03(3H, t). LRMS calcd 175.23, found 176.2 (MH⁺).

3. Preparation of 1-Propyl-2-{{2-(2-fluoropyrid-6-yl)-1H-imidazol-1-yl}methyl}-5-cyano-1H-benzimidazole

Method A

(A) Preparation of 1-n-Propyl-2-chloromethyl-5-cyanobenzimidazole hydrochloride

i) A solution of 3-amino-4-n-propylaminobenzonitrile (7.38g, 42.1 mmol) and ethyl chloroacetimidate hydrochloride (9.92 g, 63.2 mmol) in ethanol (100 mL) is heated at reflux for 17 h, then cooled and concentrated to give 1-n-Propyl-2-chloromethyl-5-cyanobenzimidazole hydrochloride. Prior to use in the next step, this material is converted to free base by adding aqueous bicarbonate and extracting with dichloromethane, drying (Na₂SO₄), and concentrating.

ii) Alternatively, chloroacetylchloride rather than ethyl chloroacetimidate can be used: To a solution of 3-amino-4-n-propylaminobenzonitrile (5.15 g, 29.4 mmol) and triethylamine (4.51 mL) in ethyl acetate (52 mL) at room temperature, chloroacetyl chloride (2.57 mL) is added slowly. After stirring the reaction mixture for 30 minutes at room temperature, acetic acid (5 mL) is added and the reaction mixture heated to reflux. After heating for 20 h, the reaction mixture is cooled to room temperature and diluted with water (50 mL). The organic solution is washed twice with 1.0 M sodium hydroxide (2 x 50 mL), then washed with an aqueous solution of 0.25 M KH₂PO₄ (50 mL), followed by brine (50 mL). The organic layer is dried (sodium sulfate), concentrated in vacuo, and the solid recrystallized by heating in ethyl acetate (20 mL), adding hexane (40 mL) and cooling to room temperature with stirring to afford 1-n-Propyl-2-chloromethyl-5-cyanobenzimidazole as brown crystals. ¹H NMR (CDCl₃): δ 8.08 (d, J = 0.8 Hz, 1H), 7.57 (dd, J = 1.65, 8.52 Hz, 1H), 7.45 (d, J = 8.24 Hz, 1H), 7.26 (s, 1H), 4.84 (s, 2H), 4.24 (t, J = 7.6 Hz, 2H), 1.94 (pentet, J = 7.4. 7.6 Hz, 2H), 1.03 (t, J = 7.4 Hz, 3H).

(B) Preparation of 1-Propyl-2-{[2-(2-fluoropyrid-6-yl)-1H-imidazol-1-yl]methyl}-5-cyano-1H-benzimidazole

To a stirring suspension of sodium hydride (2.25g of 60% in oil) in DMF (10 mL) at 0° a solution of 2-Fluoro-6-(1H-imidazol-2-yl)-pyridine (7.7 g, 47.2 mmol) in DMF (20 mL) is added. After stirring for 5 minutes, a solution of 1-n-Propyl-2-chloromethyl-5-cyanobenzimidazole (11 g, 47.2 mmol) and sodium iodide (20 mg) in DMF (80 mL) is added. The reaction mixture is stirred for 6 h, gradually warmed to room temperature. The reaction mixture is cooled, water added, the solid collected, rinsed with water and dried to afford the title compound, 1-Propyl-2-{[2-(2-fluoropyrid-6-yl)-1H-imidazol-1-yl]methyl}-5-cyano-1H-benzimidazole.

Method B

Trimethylaluminum (0.54mL of 2.0M in toluene, 1.09mmol) is added dropwise to a solution of 3-Amino-4-propylaminobenzonitrile (152mg; 0.87mmol) in dichloromethane (10mL), and the mixture was stirred at room temperature for 1hr. A solution of [2-(2-Fluoropyridin-6-yl)-imidazol-1-yl] acetic acid methyl ester (102mg; 0.43mmol) in dichloromethane (5mL) is added all at once, and the mixture heated at reflux for 16 hr. The brown solution is cooled to room temperature and treated dropwise with methanol (1mL) then water (2mL) and stirred at room temperature for 15 min. Anhydrous sodium sulfate is added until the gel becomes solid, the mixture is diluted with dichloromethane (100mL) and filtered through celite. The filtrate is concentrated to give a brown oil which is dissolved in acetic acid (7 mL) and heated at 100° for 72 hr. The mixture is cooled to room temperature and concentrated. The residue is dissolved in ethyl acetate (15mL), washed with saturated aqueous NaHCO₃ (1 x 50mL), then brine (1 x 50mL), dried (MgSO₄), and concentrated, and the residue purified by preparative thin layer chromatography to give 1-n-Propyl-2-

{[2-(2-fluoropyrid-6-yl)-1H-imidazol-1-yl]methyl}-5-cyano-1H-benzimidazole as a light brown semi-solid (57mg).

Method C

A solution of [2-(6-Fluoropyridin-2-yl)-imidazol-1-yl]acetic acid hydrochloride (274mg; 1.06mmol) and triethylamine (0.15mL; 1.06mmol) in dichloromethane (10mL) is treated dropwise with oxalyl chloride (0.64mL of a 2M solution in dichloromethane; 1.27mmol), and the resulting suspension stirred at room temperature for 2 hr.

The mixture is concentrated and the residue suspended in dichloromethane (10mL). A solution of 3-Amino-4-propylaminobenzonitrile (185mg; 1.06mmol) in dichloromethane (5mL) is added and the mixture stirred at room temperature for 16hr, then concentrated. The residue is dissolved in acetic acid (10mL) and heated at 100° for 1hr. The mixture is cooled to room temperature, concentrated, taken up in ethyl acetate (150mL), washed with saturated aqueous NaHCO₃ (50mL) and then washed with brine (50mL), dried (MgSO₄), and concentrated to give the crude product as a waxy brown solid (338mg). This crude product is slurried with diethyl ether (ca. 4mL) and a few drops of methanol, then filtered to give 1-n-Propyl-2-{[2-(2-fluoropyrid-6-yl)-1H-imidazol-1-yl]methyl}-5-cyano-1H-benzimidazole as a pale brown solid (230mg).


¹H NMR (399.96MHz, CDCl₃): δ 8.17(dd, J = 2.0, 7.6Hz, 1H, H-18), 8.05(s, 1H, H-4), 7.88(q, J = 8.0Hz, 1H, H-19), 7.52(d, J = 8.4Hz, 1H, H-6), 7.41(d, J = 8.4Hz, 1H, H-7), 7.21(s, 1H, H-14), 7.18(s, 1H, H-15), 6.28(s, 2H, H-13), 4.28(t, J = 7.6Hz, 2H, H-10), 1.68(dt, J = 7.6Hz, 2H, H-11), 0.84(t, J = 7.6Hz, 3H, H-12).

¹³C NMR (100.57MHz, CDCl₃, ¹H decoupled at 399.957MHz): δ, 162.36(d, J_{C-F} = 239.6Hz), 152.53, 148.69(d, J_{C-F} = 13.0Hz), 142.36(d, J_{C-F} = 7.6Hz), 142.31, 142.28, 138.42, 129.96, 126.55, 125.31, 124.23, 120.27(d, J_{C-F} = 3.8Hz), 119.96, 111.23, 108.52(d, J_{C-F} = 35.9Hz), 105.70, 46.10, 44.57, 23.43, 11.27.

Example 2

Synthesis of 1-Ethyl-2-{[2-(3-fluorophenyl)-pyrazol-3-yl]methyl}-5-cyano-1H-benzimidazole

5



1. Preparation of 1-(3-Fluorophenyl)-5-carboxymethylpyrazole

To a stirring solution of ethyl vinyl ether (340 mL, 3.55 mol) in diethyl ether (200 mL), cooled in an ice bath, is added dropwise a solution of malonyl dichloride (69 mL, 0.71 mol) in ether (20 mL). Stirring is continued at 0°C for 2h, then a solution of triethylamine (196 mL) and ethanol (350 mL) in ether (210 mL) is added with cooling. More ether is added to further precipitate triethylamine hydrochloride, the mixture is then filtered, and the filtrate concentrated. The residue is taken up in ethanol (710 mL), then triethyl orthoformate (177 mL, 1.07 mol) and concentrated HCl (5 mL) is added. The mixture is stirred overnight then concentrated. To the residue is added ethanol (500 mL) and 3-Fluorophenylhydrazine hydrochloride (27.6 g, 0.17 mol). The mixture is heated at reflux for 2 h, then cooled and concentrated. Ethyl acetate is added, the mixture is washed with aqueous bicarbonate, then with water, dried (Na₂SO₄), concentrated, and the residue purified by silica gel chromatography. Ethanol (20mL) and 1 N NaOH (100 mL) are added, and heated at reflux for 1 h. The mixture is cooled and washed with ethyl acetate. The aqueous layer is cooled and acidified, the solid collected by filtration, rinsed well with water and dried to give 1-(3-Fluorophenyl)-5-carboxymethylpyrazole as a gold solid. ¹H NMR (CDCl₃): δ 7.68 (d, J = 1.8 Hz, 1H), 7.41-


7.48 (m, 1H), 7.10-7.26 (m, 3H), 6.44 (d, J = 2.1 Hz, 1H), 3.77 (s, 2H). LCMS: 221.2 (MH⁺), 219.2 (MH⁻).

2. Preparation of 1-Ethyl-2-{[2-(3-fluorophenyl)-pyrazol-3-yl]methyl}-5-cyano-1H-benzimidazole

To a stirring solution of 3-amino-4-ethylaminobenzonitrile (887 mg, 5.5 mmol) and 1-(3-Fluorophenyl)-5-carboxymethylpyrazole (1.10 g, 5 mmol) in pyridine (5 mL) is added 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (1.15 g, 6 mmol). The mixture is stirred at room temperature for 17.5 h, then concentrated. The residue is cooled in an ice bath, aqueous HCl is added with stirring, the precipitate is collected by filtration, rinsed well with water and dried. The solid is added to acetic acid (75 mL) and heated at reflux for 5.5 h. The mixture is cooled, concentrated, and purified by silica gel chromatography to give 1-Ethyl-2-{[2-(3-fluorophenyl)-pyrazol-3-yl]methyl}-5-cyano-1H-benzimidazole. The product is converted to the mesylate salt in acetone. ¹H NMR (CDCl₃): δ 8.17 (s, 1H), 7.89 (d, J = 8.52 Hz, 1H), 7.68-7.75 (m 1H), 7.44-7.54 (m, 2H), 7.37-7.40 (m, 1H), 7.21-7.28 (m, 1H), 6.35 (d, J = 1.65 Hz, 1H), 4.65 (s, 2H), 4.29 (q, J = 7.14 Hz, 2H), 2.34 (s, 3H), 1.19 (t, J = 7.14 Hz, 3H). LCMS 346.0 (MH⁺), 344.4 (MH⁻).

Example 3

Synthesis of 1-Ethyl-2-{[1-(3-fluorophenyl)-1,2,4-triazol-5-yl]methyl}-5-cyano-1H-benzimidazole



1. Preparation of 1-(3-Fluorophenyl)-1,2,4-triazole

A mixture of 1,3,5-triazine (1 g, 12.3 mmol) and 3-Fluorophenylhydrazine hydrochloride (2 g, 12.3 mmol) in ethanol (20 mL) is heated at reflux overnight. After concentrating, ethyl acetate is added, the mixture is washed with aqueous bicarbonate followed by saturated aqueous sodium chloride, dried (MgSO₄), and concentrated to give 1.8 g of crude 1-(3-Fluorophenyl)-1,2,4-triazole. ¹H NMR (CDCl₃): δ 8.57 (s, 1H), 8.12 (s, 1H), 7.46-7.51 (m, 3H), 7.08-7.15 (m, 1H). LCMS 164.1 (MH⁺).

2. Preparation of 1-(3-Fluorophenyl)-5-hydroxymethyl-1,2,4-triazole

A mixture of the crude 1-(3-Fluorophenyl)-1,2,4-triazole and formaldehyde (10 mL of 37 wt % in water) is heated at 150°C in a sealed tube for 48 h. After cooling the reaction vessel, the reaction mixture is extracted with dichloromethane, dried (MgSO₄), concentrated, and the residue purified using silica gel chromatography to afford 1-(3-Fluorophenyl)-5-hydroxymethyl-1,2,4-triazole. ¹H NMR (CDCl₃): δ 8.02 (s, 1H), 7.42-7.53 (m, 3H), 7.22-7.29 (m, 1H), 4.81 (s, 2H). LCMS 194.2 (MH⁻).

3. Preparation of 1-(3-Fluorophenyl)-5-cyanomethyl-1,2,4-triazole

1-(3-Fluorophenyl)-5-hydroxymethyl-1,2,4-triazole (1.7 g) is treated with thionyl chloride (10 mL) in dichloromethane (20 mL) at room temperature overnight. The solvent is then removed. To the residue is added acetonitrile (20 mL), tetraethylammonium cyanide (2.75 g, 17.6 mmol), and triethylamine (2.5 mL, 17.6 mmol). The mixture is stirred at room temperature for 2 h. The mixture is diluted with ethyl acetate, washed with aqueous bicarbonate, and then with saturated aqueous sodium chloride, dried (Na₂SO₄), concentrated and the residue purified by silica gel chromatography to give crude 1-(3-Fluorophenyl)-5-cyanomethyl-1,2,4-triazole. ¹H NMR (CDCl₃): δ 8.06 (s, 1H), 7.52-7.62 (m, 1H), 7.20-7.30 (m, 3H), 3.98 (s, 2H). LCMS 203.0 (MH⁺), 201.2 (MH⁻).

4. Preparation of 1-(3-Fluorophenyl)-5-carboxymethyl-1,2,4-triazole

Crude 1-(3-Fluorophenyl)-5-cyanomethyl-1,2,4-triazole (0.9 g) is taken up in ethanol (50 mL) and cooled in an ice bath. HCl gas is bubbled through for 0.5 h. Water (10 mL) is added and the mixture heated at 65°C for 2 h. After cooling, most of the ethanol is removed on a roto-evaporator, 3 N NaOH (25 mL) and ethanol (25 mL) are then added. The solution is heated at reflux for 2 h, cooled, and then extracted with diethyl ether (3X).


The aqueous layer is acidified to pH 2, extracted with ethyl acetate. The organic layer is washed with saturated aqueous sodium chloride, dried (Na₂SO₄), and concentrated to give 1-(3-Fluorophenyl)-5-carboxymethyl-1,2,4-triazole. ¹H NMR (CDCl₃): δ 8.08 (s, 1H), 7.50-7.57 (m, 1H), 7.22-7.32 (m, 3H), 4.01 (s, 2H).

5. Preparation of 1-Ethyl-2-([1(3-fluorophenyl)-1,2,4-triazol-5-yl]methyl)-5-cyano-1H-benzimidazole

Using the procedure described in Example 2, Step 2, 1-(3-fluorophenyl)-5-carboxymethyl-1,2,4-triazole is converted to 1-Ethyl-2-([1-(3-fluorophenyl)-1,2,4-triazol-5-yl]methyl)-5-cyano-1H-benzimidazole. ¹H NMR (CDCl₃): δ 8.03 (s, 1H); 8.01 (s, 1H); 7.42-7.56 (m, 5H); 7.22 (m, 1H); 4.51 (s, 2H); 4.43 (q, 2H); 1.45 (t, 3H). LCMS 347.3 (MH⁺).

Example 4

Synthesis of 1-Propyl-2-[(5-phenyl-1H-tetrazol-1-yl)methyl]-1H-benzimidazole



1. Preparation of 2-(Tri-n-butyltin)-5-phenyl -1H-tetrazole


A mixture of 5-phenyl-1H-tetrazole (200 mg, 1.22 mmol) and bis(tri-n-butyltin)oxide (0.31 mL, 0.61 mmol) in ethanol (2 mL) is heated at reflux for 10 minutes. The mixture is cooled, concentrated, and the crude 2-(Tri-n-butyltin)-5-phenyl -1H-tetrazole used directly.

2. Preparation of 1-Propyl-2-[5-phenyl-1H-tetrazol-1-yl)methyl]-1H-benzimidazole

Crude 2-(tri-*n*-butyltin)-5-phenyl-1H-tetrazole (2.09 g, 4.8 mmol) and 1-propyl-2-chloromethylbenzimidazole (1.0 g, 4.8 mmol) are heated in toluene (5 mL) at reflux overnight. The solvent is removed in vacuo, the residue washed with hexane, and then purified by preparative chromatography to give 1-Propyl-2-[(5-phenyl-1H-tetrazol-1-yl)methyl]-1H-benzimidazole as the minor isomer. ¹H NMR (CDCl₃): δ 8.05-8.12 (m, 2H); 7.77 (m, 1H); 7.58-7.62 (m, 2H); 7.29-7.43 (m, 3H); 5.86 (s, 2H); 4.40 (t, 2H); 1.85 (m, 2H); 1.01 (t, 3H).

Example 5

Synthesis of 1-Ethyl-2-{[2-(3-fluorophenyl)-pyrazol-3-yl]methyl}-5-acetyl-1H-benzimidazole




A mixture of 1-Ethyl-2-{[2-(3-fluorophenyl)-pyrazol-3-yl]methyl}-5-bromo-1H-benzimidazole (200mg, 0.5 mmol), tributyl(1-ethoxyvinyl)tin (0.34 ml, 1.0 mmol), and tetrakis(triphenylphosphine)palladium(0) (29 mg) in toluene (10 mL) is heated at reflux for 1 h under argon. The solvent is removed in vacuo, the residue is then dissolved in 10% HCl (5 mL) and THF (5 mL). The mixture is stirred at room temperature for

0.5 h, then extracted with ethyl acetate. The aqueous layer is adjusted to pH 9, extracted with dichloromethane, and the organic layer dried (Na₂SO₄), concentrated, and the residue purified by preparative silica gel chromatography to give 1-Ethyl-2-{[2-(3-fluorophenyl)-pyrazol-3-yl]methyl}-5-acetyl-1H-benzimidazole. ¹H NMR (CDCl₃): δ 8.36 (d, 1H); 7.99 (dd, 1H); 7.645 (d, 1H); 7.25-7.49 (m, 4H); 7.14 (m, 1H); 6.21 (d, 1H); 4.36 (s, 2H); 4.02 (g, 2H); 2.68 (s, 3H); 1.24 (t, 3H).

10 Example 6


Synthesis of 1-Ethyl-2-{[2-(3-fluorophenyl)-pyrazol-3-yl]methyl}-5-[1-(methoxyimino)ethyl]-1H-benzimidazole



A mixture of 1-Ethyl-2-{[2-(3-fluorophenyl)-pyrazol-3-yl]methyl}-5-acetyl-1H-benzimidazole (23 mg), methoxylamine hydrochloride (15 mg, 3 eq.), and sodium acetate (15 mg, 3 eq.) in methanol (1 mL) is stirred at room temperature overnight. The solvent is removed in vacuo and aqueous sodium bicarbonate added to pH 9, then extracted with ethyl acetate. The organic layer is dried (Na₂SO₄) and concentrated to give 1-Ethyl-2-{[2-(3-fluorophenyl)-pyrazol-3-yl]methyl}-5-[1-(methoxyimino)ethyl]-1H-benzimidazole. ¹H NMR (CDCl₃): δ 7.94 (d, 1H); 7.73 (dd, 1H); 7.73 (dd, 1H); 7.63 (d, 1H); 7.52 (m, 1H); 7.38-7.40 (m, 3H); 7.27 (m, 1H); 6.19 (d, 1H); 4.32 (s, 2H); 3.87-4.10 (m, 5H); 2.30 (s, 3H); 1.22 (t, 3H).

Example 7


Synthesis of 1-Ethyl-2-{[2-(2,5-difluorophenyl)-pyrazol-3-yl]methyl}-5-[5-(4-fluorophenyl)-1,3,4-oxadiazol-2-yl]-1H-benzimidazole



5 A mixture of 1-Ethyl-2-{[2-(2,5-difluorophenyl)-pyrazol-3-yl]methyl}-1H-benzimidazole-5-carboxylate (152 mg) and 4-fluorobenzhydrazide (1.05 eq.) in phosphorous oxychloride (6 mL) is heated at reflux for 1.5 h. The mixture is cooled and concentrated, and then water (5 mL) is added to the residue.
 10 After adjusting to pH >7 with saturated aqueous bicarbonate, the solution is extracted with ethyl acetate (3X), and the combined organic layers are washed with water (2X) then with saturated aqueous sodium chloride, dried (MgSO₄), concentrated, and the residue purified by preparative silica gel chromatography to give
 15 1-Ethyl-2-{[2-(2,5-difluorophenyl)-pyrazol-3-yl]methyl}-5-[5-(4-fluorophenyl)-1,3,4-oxadiazol-2-yl]-1H-benzimidazole. LCMS MH⁺ 501.068.

Example 8


20 Synthesis of 1-Ethyl-2-{[2-(3-fluorophenyl)-1H-imidazol-1-yl]methyl}-5-(1,3,4-oxadiazol-2-yl)-1H-benzimidazole



A mixture of 1-ethyl-2-{[2-(3-fluorophenyl)-1H-imidazol-1-yl]methyl}-1H-benzimidazol-5-hydrazide (177 mg), triethylorthoformate (8 mL) and acetic acid (2 mL) is heated at reflux for 5 h. The reaction is cooled, concentrated, and the residue purified by preparative silica gel chromatography to give 1-Ethyl-2-{[2-(3-fluorophenyl)-1H-imidazol-1-yl]methyl}-5-(1,3,4-oxadiazol-2-yl)-1H-benzimidazole, which was converted to the hydrochloride salt in ethyl acetate. ¹H NMR (d6 DMSO): δ 9.32 (s, 1H), 8.19 (s, 1H), 8.03 (d, 1H), 7.97-7.95 (m, 2H), 7.87 (d, 1H), 7.72 (m, 1H), 7.66-7.60 (m, 2H), 7.54-7.50 (m, 1H), 5.99 (s, 2H), 4.34 (q, 2H), 1.31 (t, 3H), LCMS MH⁺ 389.4

Example 9

Synthesis of 1-Ethyl-2-{[2-(2,5-difluorophenyl)-1H-imidazol-1-yl]methyl}-5-amino-1H-benzimidazole



1. Preparation of 4-Fluoro-3-nitroacetanilide

4-Fluoro-3-nitroaniline (5.2 g) is treated with acetic anhydride (1.1 eq.) in dichloromethane at room temperature for 1 h. The reaction is concentrated, the residue taken up in dichloromethane and washed with aqueous sodium bicarbonate (2X) then with saturated aqueous sodium chloride, dried (MgSO₄), and concentrated to give 4.7 g of crude 4-Fluoro-3-nitroacetanilide.

2. Preparation of 3-Nitro-4-(ethylamino)acetanilide

A mixture of crude 4-Fluoro-3-nitroacetanilide (4.6 g), ethylamine (23 mL of 2M in THF), and potassium carbonate (3.5 g) in DMF (100 mL) is stirred at room temperature for 4 h, then heated at 60°C for 5 h. The reaction is allowed to cool, water (150 mL) is added, and extracted with ethyl acetate (3X). The

combined organic extracts are washed with water (3X), then washed with saturated aqueous sodium chloride, dried (MgSO₄), and concentrated to give crude 3-Nitro-4-(ethylamino)acetanilide.

3. Preparation of N-[3-Amino-4-(ethylamino)phenyl] acetamide

Crude 3-Nitro-4-(ethylamino)acetanilide (4.8 g), 10% Pd/C (0.5 g), methanol (50 mL) and ethyl acetate (200 mL) are placed in a Paar apparatus under hydrogen at 50 psi for 5 h. The mixture is filtered through Celite and concentrated to give 4 g of crude N-[3-Amino-4-(ethylamino)phenyl] acetamide

4. Preparation of N-{2-{[2-(2,5-Difluorophenyl)-1H-imidazol-1-yl]methyl}-1-ethyl -1H-benzimidazol-5-yl} acetamide


Using the procedure described in Example 1 Method A, crude N-[3-amino-4-(ethylamino)phenyl] acetamide is converted to N-{2-{[2-(2,5-Difluorophenyl)-1H-imidazol-1-yl]methyl}-1-ethyl -1H-benzimidazol-5-yl} acetamide.

5. Preparation of 1-Ethyl-2-{[2-(2,5-difluorophenyl)-1H-imidazol-1-yl]methyl} -5-amino-1H-benzimidazole

Crude N-{1-Ethyl-2-{[2-(2,5-difluorophenyl)-1H-imidazol-1-yl]methyl}-1H-benzimidazol-5-yl} acetamide is treated with 10% aqueous HCl (20 mL) in methanol (10 mL) at reflux for 1.5 h. After cooling, the methanol is removed in vacuo, and the aqueous layer washed with ethyl acetate. The aqueous layer is adjusted to pH 10 with 3 N NaOH, then extracted with ethyl acetate (3X), and the combined organic extracts washed with water (2X) then saturated aqueous sodium chloride, dried (MgSO₄), and concentrated to give 1-Ethyl-2-{[2-(2,5-difluorophenyl)-1H-imidazol-1-yl]methyl} -5-amino-1H-benzimidazole. LCMS MH⁺ 354.4.

Example 10


Synthesis of 1-Ethyl-2-{[2-(3-fluorophenyl)-pyrazol-3-yl]methyl}-5-(1,2,4-triazol-1-yl)-1H-benzimidazole



1-Ethyl-2-{[2-(3-fluorophenyl)-pyrazol-3-yl]methyl}-5-amino-1H-benzimidazole (274 mg) is treated with N,N-dimethylformamide azine hydrochloride (1 eq) in toluene (10 mL) and methoxyethanol (10 mL) and refluxed for 6 h. After cooling and concentrating, the residue is treated with water and extracted with dichloromethane (2X). The aqueous layer is adjusted to pH 8 with aqueous sodium bicarbonate and extracted with dichloromethane (2X). The combined organic extracts are washed with saturated aqueous sodium chloride, dried (MgSO₄), concentrated, purified by preparative silica gel chromatography, and triturated with ether to afford 1-Ethyl-2-{[2-(3-fluorophenyl)-pyrazol-3-yl]methyl}-5-(1,2,4-triazol-1-yl)-1H-benzimidazole, which is converted to the hydrochloride salt in ethyl acetate. LCMS MH⁺ 406.2.

Example 11

Synthesis of 1-ethyl-2-{[2-(2,5-difluorophenyl)-1H-imidazol-1-yl]methyl}-5-(1,2,3-triazol-1-yl-4-carboxylate)-1H-benzimidazole



1. Preparation of 1-Ethyl-2-{[2-(2,5-difluorophenyl)-1H-imidazol-1-yl]methyl}-5-azido-1H-benzimidazole


To an ice-cold solution of 1-ethyl-2-{[2-(2,5-difluorophenyl)-1H-imidazol-1-yl]methyl}-5-amino-1H-benzimidazole (466 mg) in acetic acid (8 mL) a solution of sodium nitrite (1.1 eq) in water (4 mL) is added dropwise. After stirring at 0°C for 1 h, a solution of sodium azide (1.3 eq) in water (5 mL) is added dropwise. Stirring is continued for 0.5 h at 0°C, and then at room temperature for 0.5 h. The reaction mixture is concentrated to 1/3 volume, water (10 mL) is added and then mixture is then extracted with ethyl acetate (3X). The combined organic layers are washed with aqueous sodium bicarbonate (2X), then washed with saturated aqueous sodium chloride, dried (MgSO₄), and concentrated to give 641 mg of crude 1-Ethyl-2-{[2-(2,5-difluorophenyl)-1H-imidazol-1-yl]methyl}-5-azido-1H-benzimidazole.

2. Preparation of 1-{2-[2-(2,5-Difluoro-phenyl)-imidazol-1-ylmethyl]-1-ethyl-1H-benzoimidazol-5-yl}-1H-[1,2,3]triazole-4-carboxylic acid ethyl ester

A mixture of crude 1-ethyl-2-{[2-(2,5-difluorophenyl)-1H-imidazol-1-yl]methyl}-5-azido-1H-benzimidazole (143 mg) and ethyl propiolate (1 eq.) in ethanol (10 mL) is heated at reflux for 6 h. The mixture is cooled, concentrated, and the residue purified by preparative silica gel chromatography to give 1-{2-[2-(2,5-Difluoro-phenyl)-imidazol-1-ylmethyl]-1-ethyl-1H-benzoimidazol-5-yl}-1H-[1,2,3]triazole-4-carboxylic acid ethyl ester. ¹H NMR (CDCl₃): δ 8.51s, (1H), 8.05(s, 1H), 7.74(m, 1H), 7.50-7.39(m, 4H), 7.21-7.17(m, 2H), 7.06(s, 1H), 5.54(s, 2H), 4.47(q, 2H), 3.86(q, 2H), 1.43(t, 3H), 1.04(t, 3H), LCMS MH⁺ 460.6.

Example 12


Synthesis of 1-ethyl-2-{[2-(2,5-difluorophenyl)-pyrazol-3-yl]methyl}-5-(1,2,3,4-tetrazol-1-yl)-1H-benzimidazole



1-Ethyl-2-{[2-(2,5-difluorophenyl)-pyrazol-3-yl]methyl}-5-amino-1H-benzimidazole (104 mg) is treated with triethylorthoformate (4 eq) in acetic acid (10 mL) at reflux for 4 h. After concentrating the solution, acetic acid is re-added, sodium azide (4 eq) added, and the mixture heated at 70°C for 3 h. After cooling, water (15 mL) is added and the mixture concentrated. The residue is taken up in ethyl acetate, washed with aqueous sodium bicarbonate (2X) then saturated aqueous sodium chloride, dried (MgSO₄), concentrated, and purified by preparative silica gel chromatography to give 1-ethyl-2-{[2-(2,5-difluorophenyl)-pyrazol-3-yl]methyl}-5-(1,2,3,4-tetrazol-1-yl)-1H-benzimidazole, which is converted to the hydrochloride salt in ethyl acetate. LCMS MH⁺ 395.018 .

Example 13

Synthesis of 1-Ethyl-2-{[2-(2,5-difluorophenyl)-pyrazol-3-yl]methyl}-5-(5-methyl-oxazol-2-yl)-1H-benzimidazole



1. Preparation of N-propargyl 1-ethyl-2-{[2-(2,5-difluorophenyl)-pyrazol-3-yl]methyl} -1H-benzimidazole-5-carboxamide

Oxalyl chloride (2.5 eq of 2M in dichloromethane) is added dropwise to a solution of 1-ethyl-2-{[2-(2,5-difluorophenyl)-pyrazol-3-yl]methyl} -1H-benzimidazole-5-carboxylate (368 mg) in DMF (5 drops) and dichloromethane (30 mL) at 0°C. The mixture is stirred at 0°C for 0.5 h, then at room temperature for 1 h. The solution is concentrated, the residue taken up in DMF (30 ml), excess propargylamine added, and the mixture stirred for 6 h.


Dilute aqueous sodium bicarbonate is added then extracted with dichloromethane (3X), the combined organic layers washed with saturated aqueous sodium chloride, dried (MgSO₄), and concentrated to give 462 mg of crude N-propargyl 1-ethyl-2-{[2-(2,5-difluorophenyl)-pyrazol-3-yl]methyl} -1H-benzimidazole-5-carboxamide.

2. Preparation of 1-Ethyl-2-{[2-(2,5-difluorophenyl)-pyrazol-3-yl]methyl}-5-(5-methyl-oxazol-2-yl)-1H-benzimidazole

A mixture of crude N-propargyl 1-ethyl-2-{[2-(2,5-difluorophenyl)-pyrazol-3-yl]methyl} -1H-benzimidazole-5-carboxamide (450 mg) and mercury(II) acetate (1 eq) is heated at reflux in acetic acid (15 mL) for 6 h. The mixture is concentrated, saturated aqueous potassium carbonate added and extracted with ethyl acetate (3X). The combined organic layers were washed with water then with saturated aqueous sodium chloride, dried (MgSO₄), concentrated, and triturated with ether to give 1-ethyl-2-{[2-(2,5-difluorophenyl)-pyrazol-3-yl]methyl}-5-(5-methyl-oxazol-2-yl)-1H-benzimidazole, which is converted to the hydrochloride salt in ethyl acetate. ¹H NMR (d₆ DMSO): δ 8.16(s,1H), 8.09-9.01m.(2H), 7.78(m,1H), 7.62(m,1H), 7.49-7.45(m,3H), 7.04(m,1H), 6.56(s,1H), 4.70(s,2H), 4.36(q,2H), 2.41(s,3H), 1.21(t,3H). LCMS MH⁺ 420.5.

Example 14

Synthesis of 2-{[2-(3-Fluorophenyl)-1H-imidazol-1-yl]methyl}-1-ethyl-5-(1,2,4-oxadiazol-3-yl)-1H-benzimidazole




1. Preparation of 1-Ethyl-2-{[2-(2,5-difluorophenyl)-1H-imidazol-1-yl]methyl}-1H-benzimidazol-5-carboxamide oxime
1-Ethyl-2-{[2-(3-fluorophenyl)-1H-imidazol-1-yl]methyl}-5-cyano-1H-benzimidazole (340 mg) is added to a solution of hydroxylamine hydrochloride (137 mg, 2eq) and triethylamine (0.3 mL, 2.2 eq) in methanol (3 mL), then heated at reflux for 2.5 h. The reaction is cooled, concentrated, water added, the solid collected and rinsed well with water and dried to give 305 mg of 1-Ethyl-2-{[2-(2,5-difluorophenyl)-1H-imidazol-1-yl]methyl}-1H-benzimidazol-5-carboxamide oxime.

2. Preparation of 1-Ethyl-2-{[2-(2,5-difluorophenyl)-1H-imidazol-1-yl]methyl}-1H-benzimidazol-5-carboxamide oxime
1-Ethyl-2-{[2-(3-fluorophenyl)-1H-imidazol-1-yl]methyl}-1H-benzimidazol-5-carboxamide oxime (161 mg) is treated with triethylorthoformate (3 mL) and boron trifluoride-THF complex (0.1 mL) at 100°C for 2.5 h. 1N HCl (0.5 mL) is added to the hot mixture; the reaction is then cooled and concentrated. After cooling in an ice bath, aqueous sodium hydroxide is added. The solid is collected, rinsed with water, then ether and dried to give 1-Ethyl-2-{[2-(2,5-difluorophenyl)-1H-imidazol-1-yl]methyl}-1H-benzimidazol-5-carboxamide oxime, which is converted to the methanesulfonate salt in acetone. Mp 183-186°C.

Example 15

Synthesis of 4-amino-3-ethylaminobenzonitrile



1. 4-nitro-3-chlorobenzonitrile

To an ice cold solution of 4-amino-3-chlorobenzonitrile (1 g, 6.6 mmol) in concentrated HCl (2.5 mL) plus water (2.5 mL) a chilled solution of sodium nitrite (0.74 g, 1.62 eq) in water (3.6 mL) is added dropwise to maintain the reaction temperature <0°C. After stirring at 0°C for 10 min, the mixture is added portion-wise to an ice cold solution of sodium nitrite (3.29 g, 7.22 eq) and copper(I) oxide (349 mg, 0.37 eq) in water 14.5 mL). Stirring is continued at 0°C for 40 min, and then at room temperature for 0.5 h. The reaction mixture is extracted with dichloromethane (2X), the combined organic layers washed with saturated aqueous sodium chloride, dried (MgSO₄), concentrated, and purified by silica gel chromatography to give 4-nitro-3-chlorobenzonitrile. ¹H NMR (CDCl₃): δ 7.96 (d, J = 8.24 Hz, 1H), 7.88 (d, J = 1.65 Hz, 1H), 7.63 (dd, J = 1.65, 8.24 Hz, 1H).

2. Preparation of 4-Ethylamino-3-chloro benzonitrile

A mixture of 4-nitro-3-chlorobenzonitrile (0.74 g, 4.1 mmol), potassium carbonate (1.68 g, 12.2 mmol), and ethylamine (4 mL, 2M in THF) in DMF (2 mL) is stirred at ambient temperature for 3 h. Additional ethylamine (2.1 mL, 2M in THF) is added, the flask stoppered, and stirring continued an additional 15.5 h. Water (50 mL) is added and extracted 2X with ethyl acetate (50 mL), and the combined organic layers are washed with water, then brine, dried (MgSO₄), and concentrated to give a mixture of 3-Ethylamine-4-nitrobenzonitrile and 4-Ethylamino-3-chloro benzonitrile which was carried forward without purification.

3. Preparation of 4-Amino-3-ethylaminobenzonitrile


To a suspension of crude 4-ethylamino-3-chloro benzonitrile (above) in conc. HCl (4 mL) tin(II) chloride dihydrate (3.16 g)

is added. After stirring at ambient temperature for 1.5 h, the mixture is poured onto ice and the solution made alkaline with 10 N aq. NaOH. The aqueous solution is extracted 2X with dichloromethane, and the combined organic layers washed with
5 brine, dried (MgSO₄), and concentrated. Trituration with ether affords 4-Amino-3-ethylaminobenzonitrile as a white solid. ¹H NMR (CDCl₃): δ 7.00 (dd, J = 1.65, 7.97 Hz, 1H), 6.83 (d, J = 1.65 Hz, 1H), 6.67 (d, J = 7.97 Hz, 1H), 3.13 (q, J = 7.14 Hz, 1.31 (t, J = 7.14 Hz, 3H).

10

Example 16

Synthesis of 2-Fluoro-6-(1H-imidazol-2-yl)-pyridine



1. Preparation of 2-Fluoropyridine-6-carboxaldehyde

Method A

2-Fluoro-6-methylpyridine (14.4 g, 0.13 mol) and tert-butoxybis(dimethylamino)methane (Bredereck's reagent; 34.9 g, 0.20 mol) are heated at 140 °C for 24 h. The reaction is cooled and diluted with THF (100 mL). Sodium periodate (75 g) in water (400 mL) is added at 0-5 °C, and the reaction mixture is then stirred for 24 h at room temperature. The precipitate is filtered through celite, and the filtrate extracted 5X with diethyl ether. The combined ether layers are washed with water,
25 brine, and dried (MgSO₄). Most of the solvent is removed by concentration at 0 °C (by keeping ice in the bath) to provide 2-Fluoropyridine-6-carboxaldehyde.

Method B

2-Fluoropyridine-6-carboxaldehyde can also be prepared as
30 follows: To a solution of diisopropylamine (6.54 mL, 1.2 equiv) in 30 mL of THF at 0 °C a solution of n-butyllithium (17.1 mL,

2.5M in hexanes) is added dropwise. Stirring is continued for 15 minutes at 0 °C, the reaction is then cooled to -78 °C. 2-Fluoro-6-methylpyridine (4.00 mL, 38.9 mmol) is added dropwise to the cold solution. The reaction mixture is stirred at -78 °C for 1 h and then quenched with DMF (4.52 mL, 1.5 equiv). The reaction is maintained at -78 °C for 30 minutes and then warmed to 0 °C. The cold solution is added to a mixture of sodium periodate (24.9 g) in 120 mL of water at 0 °C. The reaction mixture is allowed to gradually warm to room temperature over 1 h and then stirred at room temperature for 24 h. The reaction mixture is filtered through a plug of celite to remove the precipitate and the plug is washed with ether. The organic layer is separated, washed with aqueous sodium bicarbonate (1 x 40 mL), then with 0.25M KH₂PO₄ (1 x 40 mL) and then brine (1 x 40 mL). The organic solution is dried (NaSO₄) and concentrated in vacuo.

2. Preparation of 2-Fluoro-6-(1H-imidazol-2-yl)-pyridine

To a solution of the crude aldehyde from step 1, Method B (above) in methanol (12 mL) aqueous glyoxal (6.21 mL, 40 wt.% in water) is added dropwise. The solution is cooled to 0 °C and aqueous ammonium hydroxide (6.0 mL, 28 wt.% in water) is added. The reaction is allowed to warm to room temperature gradually over about an hour and then stirred another 3 h at room temperature. Most of the methanol is removed in vacuo, the reaction mixture diluted with water (10 mL) and extracted with ethyl acetate (30 mL). The organic layer is washed with brine (20 mL), diluted with hexanes (15 mL), passed through a plug of silica gel (1/4 inch deep x 1 ¼ inch diameter), and the plug washed with more 2:1 ethyl acetate/hexanes (20 mL). The combined eluents are concentrated in vacuo to yield crude 2-Fluoro-6-(1H-imidazol-2-yl)-pyridine.

Example 17

Synthesis of 2-Fluoro-6-(1H-imidazol-2-yl)-pyridine acetic acid

1. Preparation of [2-(6-fluoropyridin-2-yl)-imidazol-1-yl]acetic acid tert-butyl ester


A mixture of 2-Fluoro-6-(1H-imidazol-2-yl)pyridine (410mg; 2.51mmol), tert-butyl bromoacetate (539mg; 2.76mmol), potassium carbonate (520mg; 3.77mmol), and N,N-dimethylformamide (20mL) is stirred at room temperature for 16h. Water (60mL) is added and the mixture extracted with ethyl acetate (3 x 70mL). The organic extracts are washed with water (3 x 40mL) and brine (1 x 40mL), dried (MgSO₄), and concentrated to give [2-(6-Fluoropyridin-2-yl)-imidazol-1-yl]acetic acid tert-butyl ester as an orange oil (630mg). ¹H NMR (CDCl₃) δ 8.12 (dd,1H), 7.82(q,1H), 7.16(s,1H), 6.98(s,1H), 6.81(dd,1H), 5.14(s,2H), 1.44s,(9H), LCMS: 276.1(M⁺).

2. Preparation of 2-Fluoro-6-(1H-imidazol-2-yl)-pyridine acetic acid

A solution of [2-(6-fluoropyridin-2-yl)-imidazol-1-yl]acetic acid tert-butyl ester (630mg; 2.27mmol) and trifluoroacetic acid (4mL) in dichloromethane (4mL) is stirred at room temperature for 5hr. The reaction mixture is concentrated. The residue is treated with 3M HCl in ethyl acetate (10mL) and the suspension stirred at room temperature for 1hr. The precipitate is collected, washed with ethyl acetate, and dried to give 2-Fluoro-6-(1H-imidazol-2-yl)-pyridine acetic acid as a tan solid (581mg), ¹H NMR (d₆DMSO) δ 8.35-8.26(m,2H), 7.83(s,1H), 7.76(s,1H), 7.44-7.41(m,1H), 5.40(s,2H). LCMS: 222.1(MH⁺).

Example 18

Synthesis of 2-{[2-(2,5-di-fluorophenyl)-1H-imidazol-1-yl]methyl}-1-ethyl-1H-imidazo[4,5-b]pyridine



1. Preparation of 3-ethylamino-2-nitro-pyridine

Ice cold ethylamine (0.55 mL) was added to a solution of 3-fluoro-2-nitropyridine (400 mg, 2.8 mmol; prepared according to N. Plé and G. Quéguiner, *J. Heterocyclic Chem*, 1989, 26, 475-476) in 2:1 DMF:THF (6mL) at ambient temperature, and the mixture stirred for 0.75 h. Water was added, then extracted 2X with ethyl acetate. The combined organic layers were washed with water then with saturated aqueous sodium chloride, dried (MgSO₄), and concentrated to give 3-ethylamino-2-nitro-pyridine as a gold solid. ¹H NMR (CDCl₃): δ 7.87 (d, 1H), 7.43 (dd, 1H), 7.35 (d, 1H), 3.38 (pentet, 2H), 1.4 (t, 3H). LCMS: 121.1(MH⁺-NO₂).

2. Preparation of 2-Amino-3-ethylaminopyridine

A mixture of 3-ethylamino-2-nitro-pyridine (400 mg), 10% palladium on carbon (40 mg) and 1:1 ethyl acetate:methanol (30 mL) were placed in a Parr apparatus under hydrogen (50 psi) for 50 min. The mixture was filtered through Celite, concentrated, and the product triturated with hexane to give 2-Amino-3-ethylaminopyridine as a brown solid. More material was recovered from the filtrate. ¹H NMR (CDCl₃): δ 7.59 (d, J = 4.94 Hz, 1H), 6.81 (d, J = 6.32 Hz, 1H), 6.71 (dd, J = 4.94, 7.69 Hz, 1H), 3.12 (q, J = 7.14 Hz, 2H), 1.31 (t, J = 7.14 Hz, 3H).


3. Preparation of 2-{[2-(2,5-Difluorophenyl)-1H-imidazol-1-yl]methyl}-1-ethyl-1H-imidazo[4,5-b]pyridine

2-Amino-3-ethylaminopyridine (60 mg, 0.43 mmol) and methyl 1-carboxymethyl-2-(2,5-difluorophenyl)-imidazole (193 mg, 0.76 mmol) were used according to Example 1 Method B to prepare 2-{[2-(2,5-Difluorophenyl)-1H-imidazol-1-yl]methyl}-1-ethyl-1H-imidazo[4,5-b]pyridine, which is converted to the hydrochloride

salt in ethyl acetate. m.p. 214-216 °C. LCMS MH⁺ 322.2, MH⁻ 320.3.

Example 19

5 Synthesis of 2-{[2-(3-Fluorophenyl)-1H-imidazol-1-yl]methyl}-1-ethyl-3H-imidazo[4,5-c]pyridine



1. Preparation of 3-Chloro-4-nitropyridine N-oxide

Aqueous hydrogen peroxide (30 mL of 30%) is added dropwise to an ice cold mixture of 3-chloropyridine (6.0 g, 53 mmol) in acetic anhydride (30 mL). The mixture is allowed to stir at ambient temperature for 24 h, water is then added and the mixture concentrated. The residue is taken up in concentrated sulfuric acid (10 mL) and fuming sulfuric acid (5 mL), and then cooled to 0°C. Concentrated nitric acid (24 mL) is added slowly, and the ice bath removed. The reaction is then heated at reflux for 2 h, cooled, and poured into ice water. Ammonium bicarbonate is added carefully until pH 8 is achieved, the solution is then extracted with dichloromethane. The organic layer is washed with water, dried (NaSO₄), and concentrated to give 3-Chloro-4-nitropyridine N-oxide as a yellow solid. ¹H NMR (CDCl₃): δ 8.47 (d, J = 4 Hz, H), 8.35 (dd, J = 5, 12 Hz, 1H), 7.73 (ddd, J = 4, 9, 12 Hz, 1H).

2. Preparation of 3-Ethylamino-4-nitro-3-pyridine N-oxide

To an ice cold solution of 3-chloro-4-nitropyridine N-oxide (2.93 g, 16.8 mmol) in ethanol (30 mL) ethylamine (25 mL of 2M in THF) is added. The mixture is allowed to stir overnight at ambient temperature, then concentrated in vacuo to afford crude 3-Ethylamino-4-nitro-3-pyridine N-oxide as a yellow oil. ¹H NMR

(CDCl₃): δ 8.01 (d, J = 7 Hz, H), 7.91 (s, 1H), 7.45 (d, J = 7 Hz, 1H), 3.31 (q, J = 7 Hz, 2H), 1.40 (t, J = 7 Hz, 3H).

3. Preparation of 3-Ethylamino-4-aminopyridine


Crude 3-ethylamino-4-nitro-3-pyridine N-oxide and 10% palladium on carbon (1 g) in methanol (30 mL) are placed in a Parr apparatus under hydrogen (60 psi) for 2 days. Additional 10% palladium on carbon (880 mg) is added and returned under hydrogen (60 psi) for 2 days. The mixture was filtered through Celite, concentrated, and the product triturated with hexane-ether to afford 1.3 g of 3-Ethylamino-4-aminopyridine. ¹H NMR (d₆-DMSO): δ 7.63 (d, J = 6 Hz, 1H), 7.47 (s, 1H), 6.57 (d, J = 6 Hz, 1H), 3.05 (q, J = 7 Hz, 2H), 1.21 (t, J = 7 Hz, 3H).

4. Preparation of 2-{[2-(3-Fluorophenyl)-1H-imidazol-1-yl]methyl}-1-ethyl-3H-imidazo[4,5-c]pyridine

3-Ethylamino-4-aminopyridine (548 mg, 3.99 mmol) and methyl 1-carboxymethyl-2-(3-fluorophenyl)-imidazole (467 mg, 1.99 mmol) were used according to Example 1 Method B to prepare 2-{[2-(3-Fluorophenyl)-1H-imidazol-1-yl]methyl}-1-ethyl-3H-imidazo[4,5-c]pyridine. ¹H NMR (CDCl₃): δ 1.08 (t, J = 7 Hz, 3H), 3.89 (q, J = 7 Hz, 2H), 5.52 (s, 2H), 7.04 (s, 1H), 7.16-7.20 (m, 2H), 7.37-7.50 (m, 3H), 7.68 (d, J = 6 Hz, 1H), 8.47 (d, J = 6 Hz, 1H), 8.78 (s, 1H).

Example 20

Synthesis of 3-fluoro-(1H-imidazol-2-yl)benzene




Saturated ammonium hydroxide solution (30 mL) is slowly added to a solution of 3-fluorobenzaldehyde (12.4 g, 100 mmol) and glyoxal (17.5 mL of 40% wt in water, 120 mmol) in methanol (100 mL) at ambient temperature. After stirring for 24 h, most

of the solvent is removed at reduced pressure. Benzene is added and evaporated to remove residual water. The resulting dark oil is purified by chromatography on silica gel (2% MeOH/CH₂Cl₂) to obtain a tan solid. Trituration with ether/hexane provides 3-fluoro-(1H-imidazol-2-yl)benzene as a white solid. LRMS m/z (M+1) 163.2.

Example 21


Synthesis of 3-chloro-4-fluoro-(1H-imidazol-2-yl)benzene



A mixture of 3-chloro-4-fluoro-benzaldehyde (0.032 mol), glyoxal (40% in water, 0.038 mol) and ammonium hydroxide (28% in water, 0.16 mol) in MeOH (60 mL) is stirred at room temperature overnight. Solvent is removed in vacuo and the residue is partitioned between water and CH₂Cl₂. The organic layer is washed with brine, dried (Na₂SO₄), and concentrated. The residue is purified by column chromatography on silica gel eluting with CH₂Cl₂/MeOH (95/5) to afford 3-chloro-4-fluoro-(1H-imidazol-2-yl)benzene as a yellow solid. ¹H NMR (CDCl₃) δ 7.88 (dd, 1H), 7.70 (m, 1H), 7.19 (t, 1H), 7.17 (s, 2H). LRMS m/z (M+1) 197.0.

Example 22

Synthesis of 2,3,4-trifluoro-(1H-imidazol-2-yl)benzene




Saturated ammonium hydroxide solution (26 mL) is slowly added to a solution of 2,3,4-trifluorobenzaldehyde (5.0 g, 31.2


mmol) and glyoxal (10.75 mL of 40% wt in water, 93.7 mmol) in methanol (100 mL) at ambient temperature. After stirring for 24 h, most of the solvent is removed at reduced pressure. Benzene is added and evaporated to remove residual water. The resulting
5 dark oil is purified by chromatography on silica gel (5% MeOH/CH₂Cl₂) to obtain a tan solid. Trituration with ether/hexane provides 2,3,4-trifluoro-(1*H*-imidazol-2-yl)benzene as a white solid. LRMS *m/z* (*M*+1) 199.10.

10 **Example 23**

Synthesis of 2-(1*H*-imidazol-2-yl)-thiazole



1. Preparation of 1-ethoxymethyl-2-tributylstannanyl-1*H*-imidazole




15 1.6 *M* *n*-BuLi (12.0 mL, 19.2 mmol) is slowly added to a solution of 1-ethoxymethyl-1*H*-imidazole [available via the procedure outlined in Tang, C. C.; Davalian, D.; Huang, P.; Breslow, R. *J. Am. Chem. Soc.* **1978**, *100*, 3918] (2.20 g, 17.4
20 mmol) in THF (30 mL) at -78 °C under N₂. The reaction mixture is stirred at -78 °C for 20 min. whereupon tributyltin chloride (5.7 mL, 20.9 mmol) is slowly added. The reaction mixture is stirred at -78 °C for 10 min. and then warmed to room temperature. After stirring at room temperature for 1.5 h, the reaction mixture is
25 concentrated in vacuo. The residue is triturated with hexanes and filtered, and the filtrate is concentrated in vacuo. The residue is again triturated with hexanes and filtered, and the filtrate concentrated in vacuo. The ¹H NMR of the resulting oil indicates a 2:1 mixture of 1-ethoxymethyl-2-tributylstannanyl-1*H*-

imidazole: 1-ethoxymethyl-1*H*-imidazole. This material is used in the next reaction without further purification. Selected ¹H NMR resonances (400 MHz, CDCl₃) δ 7.21 (s, 1H), 7.14 (s, 1H), 5.24 (s, 2H) ppm.


5

2. Preparation of 2-(1-ethoxymethyl-1*H*-imidazol-2-yl)-thiazole



10 A solution of crude 1-ethoxymethyl-2-tributylstannanyl-1*H*-imidazole (previous experimental), 2-bromothiazole (1.05 mL, 11.6 mmol, 1.0 eq based on integration of ¹H NMR of crude 1-ethoxymethyl-2-tributylstannanyl-1*H*-imidazole), and Pd(PPh₃)₄ (0.67 g, 0.58 mmol) in toluene (20 mL) is stirred at 80 °C for 18 h. After cooling to room temperature, the reaction mixture is poured into saturated aqueous NaHCO₃ and extracted twice with
15 CH₂Cl₂. The combined extracts are dried over Na₂SO₄ and concentrated in vacuo. The residue is purified by flash chromatography on silica gel, eluting with 2:1 hexanes-EtOAc (+ 0.5% Et₃N). Fractions containing product are concentrated and resubjected to flash chromatography on silica gel. Elution with
20 2:1 hexanes-EtOAc (+ 0.5% Et₃N) affords (26%) of 2-(1-ethoxymethyl-1*H*-imidazol-2-yl)-thiazole as a bright yellow oil. ¹H NMR (400 MHz, CDCl₃) δ 7.82 (d, J = 3.2 Hz, 1H), 7.33 (d, J = 3.2 Hz, 1H), 7.20 (d, J = 1.2 Hz, 1H), 7.15 (d, J = 1.2 Hz, 1H), 6.03 (s, 2H), 3.56 (q, J = 7.2 Hz, 2H), 1.17 (t, J = 7.2 Hz, 3H)
25 ppm.

3. Preparation of 2-(1*H*-imidazol-2-yl)-thiazole




Concentrated HCl (10 ml) is added to a solution of 2-(1-ethoxymethyl-1H-imidazol-2-yl)-thiazole (940 mg, 4.49 mmol) in 24 mL of 1:1 EtOH-H₂O at room temperature. The solution is stirred at reflux for 3 h. The reaction mixture is then cooled to 0 °C and made basic by the addition of about 12 mL of 10 N aqueous NaOH. The mixture is back titrated to approximately pH 4 using concentrated HCl. Solid NaHCO₃ is added to the point of saturation and approximately pH 8. The mixture is then extracted twice using a mixture of THF and EtOAc. The combined extracts are dried over Na₂SO₄ and concentrated to an oily solid, which is triturated with a small amount of CH₂Cl₂. The solid is collected by filtration. The filtrate is concentrated, and the oily solid is triturated once more with CH₂Cl₂. The second resultant solid is collected by filtration and combined with the solid first obtained. The product, 2-(1H-imidazol-2-yl)-thiazole, is obtained as a slightly off-white solid. ¹H NMR (400 MHz, DMSO-d₆) δ 13.04 (br, 1H), 7.87 (d, J = 3.2 Hz, 1H), 7.70 (d, J = 3.2 Hz, 1H), 7.14 (br, 2H) ppm.

The compound of this example can be prepared essentially according to the procedures described in Example 21.

Example 24

Synthesis of 2-(1H-Imidazol-2-yl)-pyrimidine hydrochloride




A mixture of 2-Cyanopyrimidine (8.0 g, 76 mmol, prepared according to *Liebigs Ann.Chem.* 2, 1981, 333-341) and Aminoacetaldehyde dimethyl acetal (8 g, 76 mmol) is heated at 100

°C for 4 hours, cooled, and 100 mL of MeOH and 5 mL of concentrated HCl are added. The mixture is heated at reflux with stirring for 30 hours, cooled and evaporated to dryness in vacuo. 50 mL of i-PrOH is added to the residue and the mixture is heated at reflux with stirring for 30 minutes and cooled. The crystals are collected by filtration, washed with ether, and dried to give the title compound. ¹H NMR (DMSO): δ 9.08 (d, 2H), 7.87 (s, 2H), 7.75 (t, 1H). LRMS calcd 146, found 147 (MH+).

Example 25

Synthesis of 2-(1H-Imidazol-2-yl)-thiazole-4-carbonitrile



1) Preparation of 1-Dimethylsulfamoyl-1H-imidazole-2-carbothioic acid methoxymethyl-amide

To a solution of 1-Dimethylsulfamoyl-1-imidazole (2.0 g, 11.4 mmol) in 10 mL of anhydrous THF at -78 °C is added a solution of n-BuLi in Hexane (2.5 M, 4.6 mL) dropwise under N₂. After the mixture is stirred at the same temperature for about one hour, Methoxymethyl isocyanate (1.2 g, 11.4 mmol) is added dropwise at -78 °C. The reaction mixture is stirred at -78 °C for about 6 hours and slowly warmed to room temperature. After quenched with a saturated aqueous NH₄Cl solution (20 mL), the mixture is extracted with Ethyl acetate (20 mL x2). The extract is washed with water, dried and concentrated. The residue is purified on a silica gel column with 3 % MeOH in CH₂Cl₂ as eluent to give the title compound. ¹H NMR (CDCl₃): δ 8.60 (br, 1H), 7.32 (d, 1H), 7.01 (d, 1H), 5.22 (d, 2H), 3.51 (s, 3H), 3.00 (d, 6H). LRMS calcd 278, found 279 (MH+).

2) Preparation of 1H-Imidazole-2-carbothioic acid amide

A solution of 1-Dimethylsulfamoyl-1H-imidazole-2-carbothioic acid methoxymethyl-amide (1.0 g, 3.6 mmol) in Acetic acid (10 mL) and water (2 mL) is stirred at 50 °C overnight.

5 After the volatiles are evaporated in vacuo, the residue is partitioned between Ethyl acetate (40 mL) and aqueous NaHCO₃ (10 ML). The organic layer is separated, washed with water, dried and concentrated in vacuo to a solid (250 mg). ¹H NMR (DMSO): δ 12.7 (s, 1H), 9.65 (s, 1H), 9.42 (s, 1H), 7.24 (s, 1H), 7.15 (s,
10 1H).

3) Preparation of 2-(1H-Imidazole-2-yl)-thiazole-4-carboxylic acid ethyl ester

10 A mixture of 1H-Imidazole-2-carbothioic acid amide (200 mg, 1.57 mmol) and Ethyl bromopyruvate (340 mg, 1.57 mmol) is heated at 75 °C for about 5 hours. After the volatiles are evaporated in vacuo, the residue is partitioned between Ethyl acetate (40 mL) and aqueous NaHCO₃ (10 ML). The organic layer is separated, washed with water, dried and concentrated. The
20 residue is purified on a silica gel column with 3 % MeOH in CH₂Cl₂ as eluent to give 200 mg of the titled compound. ¹H NMR (CDCl₃): δ 8.15 (s, 1H), 7.20 (br, 1H), 7.10 (br, 1H), 4.33 (q, 2H), 1.23 (t, 3H). LRMS calcd 223, found 224 (MH⁺).

25 4) Preparation of 2-(1H-Imidazole-2-yl)-thiazole-4-carboxylic acid amide


A solution of 2-(1H-Imidazole-2-yl)-thiazole-4-carboxylic acid ethyl ester (190 mg) in 5 mL of MeOH is saturated with NH₃ gas, and heated at 60 °C in a sealed tube overnight. After
30 cooled, the reaction mixture is transferred to a round bottle flask and evaporated in vacuo to a solid. LRMS calcd 194, found 195 (MH⁺).

5) Preparation of 2-(1H-Imidazole-2-yl)-thiazole-4-carbonitrile

To a solution of 2-(1H-Imidazole-2-yl)-thiazole-4-carboxylic acid amide (120 mg) in 2 mL of anhydrous pyridine is added 0.25 mL of POCl₃ dropwise at 0 °C. The reaction mixture is stirred at the same temperature for 2 hours, poured into ice water, neutralized with solid NaHCO₃, and extracted with 5% MeOH in CH₂Cl₂. The extract is washed with brine, dried and concentrated to a solid. ¹H NMR (DMSO): δ 13.45 (br, 1H), 8.82 (s, 1H), 7.38 (s, 1H), 7.12 (s, 1H). LRMS calcd 176, found 177 (MH⁺).

Example 26

Synthesis of 2-[2-(6-fluoro-pyridin-2-yl)-imidazol-1-ylmethyl]-1-propyl-1-H-imidazo[4,5-c]pyridine



1. Preparation of (3-nitro-pyridin-4-yl)-propyl-amine

To a solution of 4-chloro-3-nitro-pyridine (18.0g, 114mmol) in EtOH (300ml) is added n-propylamine (46.9ml, 568mmol) slowly at 0°C and the resulting yellow mixture is warmed to room temp. After stirring for 2 h, the reaction is concentrated in vacuo. The residue is purified by flash chromatography (eluent 50% Hex/EtOAc) to give (3-nitro-pyridin-4-yl)-propyl-amine as a yellow solid. ¹H NMR (CDCl₃) δ 9.16 (s, 1H), 8.25 (d, J = 6 Hz, 1H), 8.15 (br s, 1H), 6.68 (d, J = 6 Hz, 1H), 3.29 (q, J = 6.8 Hz, 2H), 1.74 (sextet, J = 7.2 Hz, 2H), 1.02 (t, J = 7.6 Hz, 3H). LRMS calcd 181.19, found 182.5 (MH⁺).

2. Preparation of N-propyl-pyridine-3,4-diamine


A solution of (3-nitro-pyridin-4-yl)-propyl-amine(19.7g, 109mmol) in abs. MeOH(150ml) is hydrogenated at room temp. and 50 psi in the presence of 10% palladium on carbon(1.9g) for 15h. Removal of catalyst, concentration of filtrate, and
5 recrystallization the residue in dichloromethane-hexane yields N-propyl-pyridine-3,4-diamine as a tan solid. ¹H NMR (DMSO-d₆) δ 7.59(s, 1H), 7.55(d, J = 5.4 Hz, 1H), 6.30(d, J = 5.4 Hz, 1H), 5.26(br s, 1H), 4.55(br s, 2H), 3.02(q, J = 6.6 Hz, 2H), 1.56(sextet, J = 7.1 Hz, 2H), 0.92(t, 7.2 Hz, 3H). LRMS calcd
10 151.21, found 152.4 (MH+).

3. Preparation of 2-[2-(6-fluoro-pyridin-2-yl)-imidazol-1-ylmethyl]-1-propyl-1-H-imidazo[4,5-c]pyridine

Trimethylaluminum (11ml of 2.0M in toluene, 22.4mmol) is
15 added dropwise to a solution of N-propyl-pyridine-3,4-diamine(1.70g, 11.2mmol) in dichloromethane(50ml), and the mixture is stirred for 30 minutes at room temperature. To this is added a solution of [2-(6-fluoro-pyridin-2-yl)-imidazol-1-yl]-acetic acid methyl ester (1.32g, 5.61mmol) in dichloromethane
20 (10ml) The mixture is heated at reflux for 18 hours. The resulting brown mixture is poured into ice-water and filtered through celite. The filtrate is extracted with dichloromethane and the combined extracts are washed with saturated brine, dried over anhydrous Na₂SO₄, filtered, concentrated in vacuo, and
25 chromatographed on silica gel to afford 2-[2-(6-fluoro-pyridin-2-yl)-imidazol-1-ylmethyl]-1-propyl-1-H-imidazo[4,5-c]pyridine as a pale yellow solid. ¹H NMR (CDCl₃) δ 9.06(s, 1H), 8.42(d, J = 5.6 Hz, 1H), 8.18(dd, J = 2.1, 7.6 Hz, 1H), 7.88(dd, J = 7.6, 8.4 Hz, 1H), 7.29(d, J = 5.6 Hz, 1H), 7.22(s, 1H), 7.18(s, 1H), 6.88(dd, J = 2.4, 8.4 Hz, 1H), 6.29(s, 2H), 4.26(t, J = 7.6 Hz, 2H),
30 1.68(sextet, J = 7.2 Hz, 2H), 0.84(t, J = 7.2 Hz, 3H). LRMS calcd 336.37, found 337.6 (MH+).

Example 27

Synthesis of 3-Ethyl-2-{[2-(1,3-thiazol-2-yl)-1H-imidazol-1-yl]methyl}-3H-imidazo[4,5-c]pyridine



1. Preparation of 3-Chloro-4-nitro-pyridine-1-oxide

Aqueous 30% H_2O_2 (60 mL) is added dropwise to a magnetically stirred solution of 3-chloro-pyridine (12 g, 105 mmol) in acetic anhydride (60 mL) under cold conditions (0 to 10 °C). The resulting mixture is allowed to warm up to room temperature slowly and then stirred overnight at room temperature. The reaction mixture is quenched with water (50 mL), diluted with toluene and concentrated to obtain the crude N-oxide as an oil in near quantitative yield.

Fuming H_2SO_4 (25 mL) is added dropwise to a solution of crude 3-chloro-pyridine-1-oxide in concentrated H_2SO_4 (25 mL) under cold conditions (0 °C) with stirring. HNO_3 (fuming, 90%, 60 mL) is added carefully to the above mixture with caution to keep the offset of any exotherm under control, and then allowed to warm to room temperature slowly. The resulting mixture is then heated at 120 °C for 4 h with stirring, cooled, poured into ice-cold water, and extracted with CHCl_3 . The combined organic phase is washed successively with saturated aqueous NaHCO_3 , water, brine, dried over Na_2SO_4 , and concentrated in vacuo to afford 3-Chloro-4-nitro-pyridine-1-oxide as a yellow solid. ^1H NMR (300 MHz, CDCl_3): δ 8.31 (d, J = 1.5 Hz, 1H), 8.13 (dd, J = 1.5, 5.4 Hz, 1H), 7.95 (d, J = 7.2 Hz, 1H).

2. Preparation of 3-Ethylamino-4-nitro-pyridine-1-oxide

Anhydrous K_2CO_3 (19 g, 144.5 mmol) is suspended in a solution of 3-Chloro-4-nitro-pyridine-1-oxide (10 g, 57.8 mmol) in anhydrous acetonitrile (100 mL). Excess diethyl amine (2.0 M) in THF is added to the above suspension under cold conditions (ice-bath). After complete addition of diethyl amine, the reaction mixture is allowed to warm to room temperature and left stirring overnight. The reaction mixture is filtered to remove K_2CO_3 and the filtrate evaporated under reduced pressure to remove volatile solvents. The organic residue is subjected to chromatography, eluting with 30% EtOAc-hexanes to afford 3-Ethylamino-4-nitro-pyridine-1-oxide as an orange solid. 1H NMR (300 MHz, $CDCl_3$): δ 8.01 (d, J = 7.5 Hz, 1H), 7.91 (s, 1H), 7.8 (brs, NH), 7.44 (d, J = 7.2 Hz, 1H), 3.30 (t, J = 7.2 Hz, 2H), 1.39 (t, J = 7.2 Hz, 3H), m/z 184 [M+1]

3. Preparation of N^3 -Ethyl-pyridine-3,4-diamine

A solution of 3-ethylamino-4-nitro-pyridine 1-oxide (1.5 g, 8.19 mmol), in methanol (30 mL) is hydrogenated over 10% Pd-C (1.5 g) at 50-60 psi for 48 h. The catalyst is removed by filtration through a pad of celite, and the solvent is evaporated under vacuum to afford N^3 -Ethyl-pyridine-3,4-diamine as a white solid. 1H NMR (300 MHz, $CDCl_3$): δ 7.59 (s, 1H), 7.57 (d, J = 2.4 Hz, 1H), 6.56 (d, J = 5.4 Hz, 1H), 3.20 (t, J = 7.2 Hz, 2H), 1.29 (t, J = 7.2 Hz, 3H), m/z 138 [M+1]

4. Preparation of N -(3-Ethylamino-pyridin-4-yl)-2-(2-thiazol-2-yl-imidazol-1-yl)-acetamide

A solution (2.0 M) of Me_3Al (4.8 mL, 9.6 mmol) in toluene is added to a solution of the N^3 -ethyl-pyridine-3,4-diamine (1.06 g, 7.70 mmol) in anhydrous dichloromethane (40 mL) slowly under nitrogen, and the resulting mixture is stirred at room temperature for 1 h. (2-Thiazol-2-yl-imidazol-1-yl)-acetic acid methyl ester (860 mg, 3.85 mmol) is added to the above reaction

10033065
122101
mixture. The resulting suspension is refluxed overnight and then cooled to room temperature. The reaction mixture is decomposed with a few drops of MeOH. The resulting brown precipitate is suspended in 15% MeOH in CH₂Cl₂ (200 mL), subjected to sonication for 25 min, filtered through a pad of celite, and concentrated to afford a mixture of 3-Ethyl-2-{[2-(1,3-thiazol-2-yl)-1H-imidazol-1-yl]methyl}-3H-imidazo[4,5-c]pyridine and N-(3-ethylamino-pyridin-4-yl)-2-(2-thiazol-2-yl-imidazol-1-yl)-acetamide in a 2:1 ratio which is carried over to the next step without further purification. A portion of the crude residue is chromatographed over silica gel eluting with 7% MeOH-CH₂Cl₂ containing few drops of NH₄OH to obtain an analytical sample of N-(3-ethylamino-pyridin-4-yl)-2-(2-thiazol-2-yl-imidazol-1-yl)-acetamide for characterization purposes. ¹H NMR (300 MHz, CD₃OD): δ 7.95 (s, 1H), 7.87 (d, J = 3.3 Hz, 1H), 7.79 (d, J = 5.4 Hz, 1H), 7.61 (d, J = 3.3 Hz, 1H), 7.42 (d, J = 5.1 Hz, 1H), 7.33 (d, J = 1.5 Hz, 1H), 7.12 (d, J = 1.2 Hz, 1H), 5.51 (s, 2H), 3.20 (t, J = 6.9 Hz, 2H), 1.23 (t, J = 6.9 Hz, 3H); LRMS Calcd 328.38; found 329


20 5. Preparation of 3-Ethyl-2-{[2-(1,3-thiazol-2-yl)-1H-imidazol-1-yl]methyl}-3H-imidazo[4,5-c]pyridine

The mixture (2:1) of the cyclized 3-Ethyl-2-{[2-(1,3-thiazol-2-yl)-1H-imidazol-1-yl]methyl}-3H-imidazo[4,5-c]pyridine and the intermediate N-(3-ethylamino-pyridin-4-yl)-2-(2-thiazol-2-yl-imidazol-1-yl)-acetamide obtained from the above Me₃Al-mediated coupling conditions is dissolved in 30 mL of acetic acid and heated at 115 °C overnight with stirring. The next day, most of the AcOH is evaporated off under reduced pressure. The residue is diluted with 100 mL of CH₂Cl₂, washed with NaHCO₃, water, brine, dried over Na₂SO₄, and concentrated. The crude product is purified by chromatography, eluting with 3% MeOH-CH₂Cl₂ containing few drops of NH₄OH to afford 3-ethyl-2-{[2-(1,3-thiazol-2-yl)-1H-imidazol-1-yl]methyl}-3H-imidazo[4,5-c]pyridine

as an white solid. ^1H NMR (300 MHz, CD_3OD): δ 8.87 (s, 1H), 8.30 (d, J = 5.7 Hz, 1H), 7.75 (d, J = 3.3 Hz, 1H), 7.57 (d, J = 4.8 Hz, 1H), 7.52 (d, J = 3.3 Hz, 1H), 7.39 (d, J = 1.5 Hz, 1H), 7.18 (d, J = 1.5 Hz, 1H), 6.29 (s, 2H), 4.52 (q, J = 7.5 Hz, 2H), 1.44 (t, J = 7.5 Hz, 3H); LRMS Calcd 310.38 found 311; Anal. Calcd for CHNS : C, 58.05, H 4.55, N 27.08. Found C 57.79 ; H, 4.38; N, 27.04.

Example 28

Synthesis of 3-ethyl-6-isopropyl-2- $\{[2-(1,3\text{-thiazol-2-yl})-1H\text{-imidazol-1-yl}]methyl\}$ -3H-imidazo[4,5-c]pyridine and 1-ethyl-6-isopropyl-2- $\{[2-(1,3\text{-thiazol-2-yl})-1H\text{-imidazol-1-yl}]methyl\}$ -1H-imidazo[4,5-c]pyridine



1. Preparation of isobutyrylacetic acid

A mixture of ethyl isobutyrylacetate (20.47 g, 129 mmol) and 1.5 M aq NaOH (250 mL) is stirred at room temperature overnight. The solution is then cooled to 0 $^{\circ}\text{C}$ and acidified to pH 1-2 by the slow addition of conc. HCl (~35 mL). The solution is then saturated with NaCl and extracted thrice with EtOAc and once with CHCl_3 . The combined extracts are dried over Na_2SO_4 and concentrated to 14.46 g (86%) of isobutyrylacetic acid as an oil. ^1H NMR (CDCl_3 , 300 MHz) reveals a mixture of keto and enol forms (~4:1), in favor of the keto form. Keto form: δ 3.57 (s, 2H), 2.73 (sept, J = 6.9 Hz, 1H), 1.16 (d, J = 6.9 Hz, 6H) ppm. Enol form: 11.85 (s, 1H), 5.03 (s, 1H), 2.44 (sept, J = 6.9 Hz, 1H), 1.10 (d, J = 6.9 Hz, 6H) ppm.

2. Preparation of 4-hydroxy-3-isobutyryl-6-isopropyl-pyran-2-one

To a solution of the isobutyrylacetic acid (14.45 g, 111 mmol) in THF (200 mL) at room temperature under N₂ is added CDI (19.8 g, 122 mmol) in one portion. The yellow solution is stirred at room temperature for 20 h and then concentrated in vacuo. The residue is dissolved in CH₂Cl₂ and washed with 10% aq HCl (100 mL) followed by H₂O (50 mL). The aqueous washes are reextracted once with CH₂Cl₂, and the combined extracts are dried over Na₂SO₄ and concentrated to provide 4-hydroxy-3-isobutyryl-6-isopropyl-pyran-2-one as a yellow oil. This material is sufficiently pure to be used without further purification in the next reaction. ¹H NMR (CDCl₃, 300 MHz) δ 17.02 (s, 1H), 5.92 (s, 1H), 3.94 (sept, J = 6.9 Hz, 1H), 2.72 (sept, J = 6.9 Hz, 1H), 1.25 (d, J = 6.9 Hz, 6H), 1.16 (d, J = 6.9 Hz, 6H) ppm.

3. Preparation of 4-hydroxy-6-isopropyl-pyran-2-one

A solution of 4-hydroxy-3-isobutyryl-6-isopropyl-pyran-2-one (10.26 g, 45.8 mmol) in conc. H₂SO₄ (40 mL) is stirred at 130 °C for 15 min. The dark reaction mixture is then cooled to 0 °C, and crushed ice (~200 g) is added with stirring. The resulting solution is extracted thrice with Et₂O, and the combined extracts are dried over Na₂SO₄ and concentrated. The crude material is purified by flash chromatography on silica gel. Gradient elution with 3:2 hexanes-EtOAc, 1:1 EtOAc-hexanes, 2:1 EtOAc-hexanes, and 3:1 EtOAc-hexanes affords 4-hydroxy-6-isopropyl-pyran-2-one as a light yellow solid. ¹H NMR (CDCl₃, 300 MHz) δ 11.2 (br, 1H), 6.00 (d, J = 1.8 Hz, 1H), 5.58 (d, J = 1.8 Hz, 1H), 2.73 (sept, J = 6.9 Hz, 1H), 1.22 (d, J = 6.9 Hz, 1H) ppm.

4. Preparation of 4-hydroxy-6-isopropyl-1H-pyridin-2-one

A mixture of 4-hydroxy-6-isopropyl-pyran-2-one (4.82 g, 31.3 mmol) in conc. NH₄OH (15 mL) is stirred at 100 °C in a sealed tube for 4 h. The solution is then transferred to a recovery flask and concentrated in vacuo. Toluene is used to

azeotropically remove any remaining water. 4-Hydroxy-6-isopropyl-1H-pyridin-2-one is obtained as a tan powder. ¹H NMR (DMSO-d₆, 300 MHz) δ 10.96 (br s, 1H), 10.36 (br s, 1H), 5.59 (d, J = 2.1 Hz, 1H), 5.33 (d, J = 2.1 Hz, 1H), 2.64 (sept, J = 6.9 Hz, 1H), 1.11 (d, J = 6.9 Hz, 6H) ppm.

5. Preparation of 4-hydroxy-6-isopropyl-3-nitro-1H-pyridin-2-one

A solution of 4-hydroxy-6-isopropyl-1H-pyridin-2-one (1.76 g, 11.5 mmol) in 55-60 mL of 50% aq HNO₃ is stirred at 70 °C for 2.5 h. After cooling, the reaction mixture is poured into ice-cold H₂O (~100 mL). The resulting solution is allowed to sit in the refrigerator for 3.5 days. The solid that precipitates is collected by filtration, washed with water, and dried to afford 4-hydroxy-6-isopropyl-3-nitro-1H-pyridin-2-one as yellow crystals. ¹H NMR (DMSO-d₆, 300 MHz) δ 12.30 (br s, 1H), 11.82 (br s, 1H), 5.82 (d, J = 1.2 Hz, 1H), 2.71 (sept, J = 6.9 Hz, 1H), 1.13 (d, J = 6.9 Hz, 6H) ppm.

6. Preparation of 4-chloro-6-isopropyl-3-nitro-1H-pyridin-2-one

To a suspension of 4-hydroxy-6-isopropyl-3-nitro-1H-pyridin-2-one (3.02 g, 15.2 mmol) in CH₃CN (60 mL) at room temperature is added benzyltriethylammonium chloride (13.88 g, 61.0 mmol). The mixture is stirred at room temperature for 5-10 min, and then POCl₃ (6.25 mL, 67.1 mmol) is added. The reaction mixture is stirred at room temperature for 15 min and then at reflux for 1 h. After cooling, the reaction mixture is concentrated in vacuo. The flask is then placed in an ice bath, and the residue treated with H₂O (~60 mL). The mixture is stirred for 4 h while slowly warming to rt. The solid that forms over this period of time is collected by filtration and then washed with H₂O and a small amount of hexanes. 4-Chloro-6-isopropyl-3-nitro-1H-pyridin-2-one is obtained as a yellow solid. ¹H NMR (CDCl₃, 300 MHz) δ 13.13

(br s, 1H), 6.24 (s, 1H), 2.89 (sept, J = 6.9 Hz, 1H), 1.31 (d, J = 6.9 Hz, 6H) ppm.

7. Preparation of 4-amino-6-isopropyl-3-nitro-1H-pyridin-2-one

5 A mixture of 4-chloro-6-isopropyl-3-nitro-1H-pyridin-2-one (700 mg, 3.23 mmol) in 6-7 mL of 7 N NH₃ in MeOH is stirred at 100 °C in a sealed tube for 1.5 h. After cooling, the reaction mixture is concentrated in vacuo. The residue is suspended in H₂O, filtered, washed with H₂O, and dried, yielding 4-amino-6-isopropyl-3-nitro-1H-pyridin-2-one. ¹H NMR (DMSO-d₆, 300 MHz) δ 10.96 (br s, 1H), 8.15 (br s, 2H), 5.70 (d, J = 0.9 Hz, 1H), 2.58 (sept, J = 6.9 Hz, 1H), 1.11 (d, J = 6.9 Hz, 6H) ppm.

8. Preparation of 2-chloro-6-isopropyl-pyridin-4-ylamine

15 A mixture of 4-amino-6-isopropyl-3-nitro-1H-pyridin-2-one (555 mg, 2.81 mmol) in POCl₃ (10 mL) is stirred at reflux under N₂ for 1.5 h. After cooling, the solution is concentrated in vacuo. The reaction flask is then placed in an ice bath, and crushed ice (~20 g) is added to the residue. The mixture is 20 swirled vigorously for several minutes and then stirred at room temperature for 30 min. The mixture is then extracted with EtOAc. The extract is washed with an additional 20 mL of H₂O. The aqueous washes are reextracted once with EtOAc, and the combined extracts are dried over Na₂SO₄ and concentrated, 25 yielding 2-chloro-6-isopropyl-pyridin-4-ylamine as an orange oil. This material is used without further purification. ¹H NMR (DMSO-d₆, 300 MHz) δ 7.27 (br s, 2H), 6.65 (s, 1H), 2.77 (sept, J = 6.9 Hz, 1H), 1.13 (d, J = 6.9 Hz, 6H) ppm.

9. Preparation of 6-isopropyl-pyridine-3,4-diamine hydrochloride

30 A solution of 2-chloro-6-isopropyl-pyridin-4-ylamine (684 mg, 3.17 mmol) in MeOH (10-15 mL) containing 10% Pd/C (~70 mg) is stirred under an atmosphere of H₂ (double stuffed balloon) for 5

h. The reaction mixture is then filtered through a pad of Celite using MeOH, and the filtrate is concentrated in vacuo. The residue is dissolved in a small amount of MeOH and further diluted with toluene. The solution is then concentrated in vacuo. This is repeated once more, affording 6-isopropyl-pyridine-3,4-diamine hydrochloride as an orange-yellow solid. ¹H NMR (DMSO-d₆, 300 MHz) δ 13.26 (br, 1H), 7.45, 7.40 (br m, 3H), 6.59 (s, 1H), 5.48 (br, 2H), 2.94 (sept, J = 6.9 Hz, 1H), 1.19 (d, J = 6.9 Hz, 6H) ppm.

10. Preparation of 6-isopropyl-2-(2-thiazol-2-yl-imidazol-1-ylmethyl)-1H-imidazo[4,5-c]pyridine

To a mixture of 6-isopropyl-pyridine-3,4-diamine hydrochloride (265 mg, 1.41 mmol) in 1,2-dichloroethane (10 mL) at room temperature under N₂ is added 2.0 M AlMe₃ in toluene (2.12 mL, 4.24 mmol). The mixture is stirred at room temperature for 45 min and then treated with a solution of (2-thiazol-2-yl-imidazol-1-yl)-acetic acid methyl ester (252 mg, 1.13 mmol) in 1,2-dichloroethane (2 mL) via cannula, followed by a 1 mL rinse. The reaction mixture is then stirred at reflux for 20 h. After cooling, the reaction mixture is diluted with some MeOH (~2 mL), followed by some water and then saturated aq NaHCO₃. The resulting mixture is stirred vigorously for 15 min. The mixture is then extracted thrice with CHCl₃ containing some (~5%) MeOH. The combined extracts are dried over Na₂SO₄ and concentrated. The crude residue is then triturated with CHCl₃, and the resulting mixture filtered. The solid is washed with a small amount of CHCl₃ and dried, affording N-(4-amino-6-isopropyl-pyridin-3-yl)-2-(2-thiazol-2-yl-imidazol-1-yl)-acetamide as an off-white solid. ¹H NMR (DMSO-d₆, 300 MHz) δ 9.52 (s, 1H), 7.86 (s, 1H), 7.85 (d, J = 3.3 Hz, 1H), 7.75 (d, J = 3.6 Hz, 1H), 7.39 (s, 1H), 7.04 (d, J = 0.9 Hz, 1H), 6.45 (s, 1H), 5.65 (s, 2H), 5.38 (s, 2H), 2.74 (sept, J = 6.9 Hz, 1H), 1.13 (d, J = 6.9 Hz,

6H) ppm. The filtrate is concentrated and further purified by flash chromatography on silica gel. Elution with 20:1 CHCl₃-MeOH followed by 15:1 CHCl₃-MeOH affords 6-isopropyl-2-(2-thiazol-2-yl-imidazol-1-ylmethyl)-1H-imidazo[4,5-c]pyridine (160 mg, 44%).

¹H NMR (CDCl₃, 300 MHz) δ 12.23 (br, 1H), 8.97 (br, 1H), 7.94 (d, J = 2.7 Hz, 1H), 7.43 (d, J = 3.3 Hz, 1H), 7.23 (br, 1H), 7.20 (d, J = 0.9 Hz, 1H), 7.04 (s, 1H), 5.82 (s, 2H), 3.14 (sept, J = 6.9 Hz, 1H), 1.30 (d, J = 6.9 Hz, 6H) ppm.

11. Preparation of 3-ethyl-6-isopropyl-2-{[2-(1,3-thiazol-2-yl)-1H-imidazol-1-yl]methyl}-3H-imidazo[4,5-c]pyridine and 1-ethyl-6-isopropyl-2-{[2-(1,3-thiazol-2-yl)-1H-imidazol-1-yl]methyl}-1H-imidazo[4,5-c]pyridine

To a solution of 6-isopropyl-2-(2-thiazol-2-yl-imidazol-1-ylmethyl)-1H-imidazo[4,5-c]pyridine (160 mg, 0.493 mmol) in DMF (1.0 mL) at room temperature under N₂ is added Cs₂CO₃ (168 mg, 0.518 mmol) followed by iodoethane (0.047 mL, 0.592 mmol). The mixture is stirred at room temperature for 1 h and then diluted with water and extracted twice with CH₂Cl₂. The combined


extracts are dried over K₂CO₃ and concentrated. The residue is purified by preparative TLC, developing with 15:1 CHCl₃-MeOH (+0.5% Et₃N). The less polar (top) band affords 3-ethyl-6-isopropyl-2-{[2-(1,3-thiazol-2-yl)-1H-imidazol-1-yl]methyl}-3H-imidazo[4,5-c]pyridine. ¹H NMR (CDCl₃, 300 MHz) δ 8.70 (d, J = 1.2 Hz, 1H), 7.83 (d, J = 3.0 Hz, 1H), 7.53 (d, J = 0.9 Hz, 1H), 7.39 (d, J = 3.0 Hz, 1H), 7.18 (d, J = 0.9 Hz, 1H), 7.14 (d, J = 0.9 Hz, 1H), 6.32 (s, 2H), 4.32 (q, J = 7.2 Hz, 2H), 3.18 (sept, J = 6.9 Hz, 1H), 1.34 (d, J = 6.9 Hz, 6H), 1.17 (t, J = 7.2 Hz, 3H) ppm. The more polar (bottom) band affords 74 mg (43%) of 1-

ethyl-6-isopropyl-2-{[2-(1,3-thiazol-2-yl)-1H-imidazol-1-yl]methyl}-1H-imidazo[4,5-c]pyridine. ¹H NMR (CDCl₃, 300 MHz) δ 9.01 (s, 1H), 7.83 (d, J = 3.2 Hz, 1H), 7.38 (d, J = 3.2 Hz, 1H), 7.15 (s, 1H), 7.12 (s, 1H), 7.09 (s, 1H), 6.32 (s, 2H), 4.21 (q,

J = 7.2 Hz, 2H), 3.17 (sept, J = 6.8 Hz, 1H), 1.34 (d, J = 6.8 Hz, 6H), 1.10 (t, J = 7.2 Hz, 3H) ppm.

Example 29

5 Synthesis of 1-ethyl-6-isopropyl-2-{[2-(1,3-thiazol-2-yl)-1H-imidazol-1-yl]methyl}-1H-imidazo[4,5-c]pyridine



1. Preparation of 4-ethylamino-6-isopropyl-3-nitro-1H-pyridin-2-one

10 To a mixture of 4-chloro-6-isopropyl-3-nitro-1H-pyridin-2-one (Example 28, step 6) (500 mg, 2.31 mmol) in CH₃CN at room temperature is added EtNH₂•HCl (565 mg, 6.92 mmol) followed by Et₃N (1.29 mL, 9.23 mmol). The reaction mixture is then stirred in a sealed tube at 100 °C for 1.5 h. After cooling, the
15 reaction mixture is diluted with CH₂Cl₂ and washed with H₂O (2 x 25 mL). The aqueous washes are reextracted once with CH₂Cl₂, and the combined extracts are dried over Na₂SO₄ and concentrated to provide 4-ethylamino-6-isopropyl-3-nitro-1H-pyridin-2-one as a yellow-orange solid. The material is sufficiently pure to be
20 used without further purification. ¹H NMR (DMSO-d₆, 400 MHz) δ 11.1 (br, 1H), 8.93 (m, 1H), 5.73 (s, 1H), 3.37 (dq, J = 7.4, 6.8 Hz, 2H), 2.67 (sept, J = 6.8 Hz, 1H), 1.17 (t, J = 7.4 Hz, 3H), 1.15 (d, J = 6.8 Hz, 6H) ppm.

25 2. Preparation of (2-chloro-6-isopropyl-3-nitro-pyridin-4-yl)-ethyl-amine

A mixture of 4-ethylamino-6-isopropyl-3-nitro-1H-pyridin-2-one (520 mg, 2.31 mmol) in POCl₃ is stirred at reflux for 1.5 h. After cooling, the solution is concentrated in vacuo. The

reaction flask is then placed in an ice bath and crushed ice is added to the residue. The mixture is swirled vigorously for a few minutes and then stirred at 0 °C for 1 h. The mixture is then extracted with EtOAc. The extract is washed with additional
5 H₂O (20 mL) and brine (20 mL). The extract is dried over Na₂SO₄ and concentrated to afford (2-chloro-6-isopropyl-3-nitro-pyridin-4-yl)-ethyl-amine as a yellow solid. The material is sufficiently pure to be used without further purification. ¹H NMR (CDCl₃, 400 MHz) δ 6.62 (br, 1H), 6.44 (s, 1H), 3.31 (m, 2H),
10 2.91 (sept, J = 6.8 Hz, 1H), 1.33 (t, J = 7.4 Hz, 3H), 1.26 (d, J = 6.8 Hz, 6H) ppm.

3. Preparation of N⁴-ethyl-6-isopropyl-pyridine-3,4-diamine hydrochloride

15 A solution of (2-chloro-6-isopropyl-3-nitro-pyridin-4-yl)-ethyl-amine (0.56 g, 2.30 mmol) in MeOH (20 mL) containing 10% Pd/C (~50 mg) is stirred under an atmosphere of H₂ for 3.5 h. The reaction mixture is then filtered through a pad of Celite using MeOH. The filtrate is concentrated in vacuo. Toluene is
20 added to the residue and then removed in vacuo. This is repeated once more, yielding N⁴-ethyl-6-isopropyl-pyridine-3,4-diamine hydrochloride as a dark solid. ¹H NMR (CD₃OD, 400 MHz) δ 7.46 (s, 1H), 6.60 (s, 1H), 3.43 (q, J = 7.4 Hz, 2H), 3.03 (sept, J = 6.8 Hz, 1H), 1.35 (t, J = 7.4 Hz, 3H), 1.33 (d, J = 6.8 Hz,
25 6H) ppm.


4. Preparation of 1-ethyl-6-isopropyl-2-{[2-(1,3-thiazol-2-yl)-1H-imidazol-1-yl]methyl}-1H-imidazo[4,5-c]pyridine

To a suspension of N⁴-ethyl-6-isopropyl-pyridine-3,4-diamine
30 hydrochloride (126 mg, 0.582 mmol) in CH₂Cl₂ (5 mL) at room temperature under N₂ is added 2.0 M AlMe₃ in toluene (0.35 mL, 0.728 mmol). The dark solution is then stirred at room temperature for 45 min. Next, a solution of (2-Thiazol-2-yl-

imidazol-1-yl)-acetic acid methyl ester (65 mg, 0.291 mmol) in CH₂Cl₂ (2 mL) is added via cannula followed by a 1 mL rinse. The reaction mixture is stirred at reflux overnight. After cooling, it is treated with MeOH (~0.5 mL) and then diluted with 0.5 N NaOH and additional CH₂Cl₂. The mixture is stirred vigorously for 15 min and then extracted thrice with CH₂Cl₂ containing some (~5%) MeOH. The combined extracts are dried over K₂CO₃ and concentrated to 156 mg of a crude mixture of the desired 1-ethyl-6-isopropyl-2-{[2-(1,3-thiazol-2-yl)-1H-imidazol-1-yl]methyl}-1H-imidazo[4,5-c]pyridine and N-(4-amino-6-isopropyl-pyridin-3-yl)-2-(2-thiazol-2-yl-imidazol-1-yl)-acetamide. This material is then dissolved in AcOH (6 mL), and the resulting solution is stirred at reflux for 5 h. After cooling, the reaction mixture is concentrated in vacuo. The residue is dissolved in CH₂Cl₂ and washed with 0.5 N NaOH. The aqueous layer is reextracted twice with CH₂Cl₂, and the combined extracts are dried over K₂CO₃ and concentrated. The crude material is purified by preparative TLC. Developing with 15:1 CHCl₃ yields 1-ethyl-6-isopropyl-2-{[2-(1,3-thiazol-2-yl)-1H-imidazol-1-yl]methyl}-1H-imidazo[4,5-c]pyridine. ¹H NMR (CDCl₃, 400 MHz) δ 9.01 (s, 1H), 7.83 (d, J = 3.2 Hz, 1H), 7.38 (d, J = 3.2 Hz, 1H), 7.15 (s, 1H), 7.12 (s, 1H), 7.09 (s, 1H), 6.32 (s, 2H), 4.21 (q, J = 7.2 Hz, 2H), 3.17 (sept, J = 6.8 Hz, 1H), 1.34 (d, J = 6.8 Hz, 6H), 1.10 (t, J = 7.2 Hz, 3H) ppm.

25 Example 30

Synthesis of 1-ethyl-2-{[2-(1,3-thiazol-2-yl)-4-(trifluoromethyl)-1H-imidazol-1-yl]methyl}-1H-imidazo[4,5-c]pyridine



1. Preparation of 2-(4-trifluoromethyl-1H-imidazol-2-yl)-thiazole

Analogous to the published procedure (*J. Med. Chem.* **2000**, 43, 2165-2175), 1,1-dibromo-3,3,3-trifluoroacetone (4.47 g, 16.6 mmol) is added to a solution of NaOAc (2.72 g, 33.1 mmol) in H₂O (15 mL). The resulting solution is stirred at 100 °C for 30 min and then allowed to cool to rt. Next, a solution of 2-thiazolecarboxaldehyde (1.5 g, 13.3 mmol) in MeOH (15 mL), containing conc. NH₄OH (5 mL), is added. The reaction mixture is then stirred at room temperature for 20 h. It is then filtered, the solid washed with H₂O, and dried. 2-(4-Trifluoromethyl-1H-imidazol-2-yl)-thiazole is obtained as an off-white fluffy solid. ¹H NMR (DMSO-*d*₆, 400 MHz) δ 13.81 (br, 1H), 7.97 (d, *J* = 3.2 Hz, 1H), 7.94 (s, 1H), 7.87 (d, *J* = 3.2 Hz, 1H) ppm.

2. Preparation of (2-thiazol-2-yl-4-trifluoromethyl-imidazol-1-yl)-acetic acid methyl ester

To a suspension of NaH (109 mg of a 60% dispersion in mineral oil, 2.74 mmol) in THF (10 mL) at 0 °C under N₂ is added a solution of 2-(4-trifluoromethyl-1H-imidazol-2-yl)-thiazole (500 mg, 2.28 mmol) in THF (4 mL) via cannula, followed by a 1 mL rinse. The mixture is stirred at 0 °C for 30 min and then treated with methyl bromoacetate (0.28 mL, 2.97 mmol). The reaction mixture is stirred overnight while slowly warming to rt. It is then diluted with H₂O (10 mL) and the resulting mixture stirred vigorously for 5 min. The mixture is further diluted with some brine and then extracted twice with EtOAc. The

combined extracts are dried over Na₂SO₄ and concentrated. The crude material is purified by flash chromatography on silica gel. Elution with 4:1 hexanes-EtOAc followed by 3:1 hexanes-EtOAc affords unreacted 2-(4-trifluoromethyl-1H-imidazol-2-yl)-thiazole as well as (2-thiazol-2-yl-4-trifluoromethyl-imidazol-1-yl)-acetic acid methyl ester. ¹H NMR (CDCl₃, 400 MHz) δ 7.82 (d, J = 3.2 Hz, 1H), 7.40 (d, J = 3.2 Hz, 1H), 7.35 (s, 1H), 5.41 (s, 2H), 3.77 (s, 3H) ppm.


3. Preparation of 1-ethyl-2-{[2-(1,3-thiazol-2-yl)-4-(trifluoromethyl)-1H-imidazol-1-yl]methyl}-1H-imidazo[4,5-c]pyridine

To a mixture of N⁴-ethyl-pyridine-3,4-diamine (94 mg, 0.687 mmol) in CH₂Cl₂ (5 mL) at room temperature under N₂ is added 2.0 M AlMe₃ in toluene (0.43 mL, 0.858 mmol), and the resulting mixture is stirred at room temperature for 1 h. Next, a solution of (2-thiazol-2-yl-4-trifluoromethyl-imidazol-1-yl)-acetic acid methyl ester (100 mg, 0.343 mmol) in CH₂Cl₂ is added via cannula. The reaction mixture is then stirred at reflux overnight. After cooling to rt, it is treated with MeOH (~1 mL). The mixture is further diluted with 0.5 N aq NaOH (10 mL) and then stirred vigorously for 30 min. The mixture is then extracted thrice with CH₂Cl₂ containing some (~5%) MeOH, and the combined extracts are dried over K₂CO₃ and concentrated. The crude mixture of 1-ethyl-2-{[2-(1,3-thiazol-2-yl)-4-(trifluoromethyl)-1H-imidazol-1-yl]methyl}-1H-imidazo[4,5-c]pyridine and N-(4-amino-pyridin-3-yl)-2-(2-thiazol-2-yl-4-trifluoromethyl-imidazol-1-yl)-acetamide is dissolved in AcOH (6 mL), and the resulting solution stirred at reflux overnight. The reaction mixture is then concentrated, and the residue is dissolved in CH₂Cl₂ and washed with 0.5 N aq NaOH. The aqueous layer is reextracted once with CH₂Cl₂, and the combined extracts are dried over K₂CO₃ and concentrated. The crude residue is purified by preparative TLC. Developing with

15:1 CHCl₃-MeOH affords 1-ethyl-2-{[2-(1,3-thiazol-2-yl)-4-(trifluoromethyl)-1H-imidazol-1-yl]methyl}-1H-imidazo[4,5-c]pyridine. ¹H NMR (CDCl₃, 400 MHz) δ 9.10 (s, 1H), 8.47 (d, J = 6.0 Hz, 1H), 7.88 (d, J = 3.2 Hz, 1H), 7.58 (s, 1H), 7.48 (d, J = 3.2 Hz, 1H), 7.31 (d, J = 5.6 Hz, 1H), 6.37 (s, 2H), 4.31 (q, J = 7.2 Hz, 2H), 1.24 (t, J = 7.2 Hz, 3H) ppm.

Example 31

Synthesis of 3-ethyl-2-{[2-(6-fluoropyridin-2-yl)-1H-imidazol-1-yl]methyl}-6-propyl-3H-imidazo[4,5-c]pyridine



1. Preparation of N-(4-amino-6-propyl-pyridin-3-yl)-acetamide

To a mixture of 6-propyl-pyridine-3,4-diamine hydrochloride (prepared from ethyl butyrylacetate as described in **Example 28**, steps 1-9) (187 mg, 1.00 mmol) in CH₂Cl₂ (4 mL) at 0 °C under N₂ is added pyridine (0.24 mL, 3.00 mmol) followed by DMF (2 mL) (for solubility). Next, Ac₂O (0.104 mL, 1.1 mmol) is added. The reaction mixture is stirred for 3 h while slowly warming to rt. It is then poured into saturated aq NaHCO₃ and extracted thrice with CH₂Cl₂. The combined extracts are dried over K₂CO₃ and concentrated to 114 mg of crude N-(4-amino-6-propyl-pyridin-3-yl)-acetamide, which was used without further purification. ¹H NMR (CDCl₃, 300 MHz) δ 8.73 (br, 1H), 8.00 (s, 1H), 8.43 (s, 1H), 4.63 (br, 2H), 2.54 (m, 2H), 2.13 (s, 3H), 1.63 (sextet, J = 7.2 Hz, 2H), 0.89 (t, J = 7.2 Hz, 3H) ppm. Electrospray MS: m/z 194 [M + 1].

2. Preparation of N³-ethyl-6-propyl-pyridine-3,4-diamine

10038069-122101
A solution of the crude *N*-(4-amino-6-propyl-pyridin-3-yl)-acetamide (114 mg, 0.59 mmol) in THF (2.5 mL) at 0 °C under N₂ is treated with 0.5 M EtNMe₂•AlH₃ in toluene (2.4 mL, 1.18 mmol). The reaction mixture is stirred for 4 h while slowly warming to
5 rt, whereupon it is treated with additional 0.5 M EtNMe₂•AlH₃ in toluene (2.4 mL, 1.18 mmol). After stirring at room temperature overnight, the reaction mixture is cooled to 0 °C and treated with moist Na₂CO₃. The mixture is diluted with CH₂Cl₂ and stirred vigorously for 15 min. It is then filtered through a pad of
10 Celite, and the filtrate concentrated. The crude material is partially purified by preparative thin layer chromatography, developing with 15:1 CHCl₃-MeOH (+1% Et₃N). The band containing the desired *N*³-ethyl-6-propyl-pyridine-3,4-diamine does not separate well from the approximately 10% of the slightly more
15 polar, unreacted *N*-(4-amino-6-propyl-pyridin-3-yl)-acetamide. Both bands are collected, and the mixture is carried into the next reaction. ¹H NMR (CDCl₃, 300 MHz) δ 7.74 (s, 1H), 6.42 (s, 1H), 4.10 (br, 1H), 3.67 (m, 1H), 3.10 (q, J = 7.2 Hz, 2H), 2.56 (m, 2H), 1.67 (sextet, J = 7.2 Hz, 2H), 1.25 (t, J = 7.2 Hz, 3H),
20 0.91 (t, J = 7.2 Hz, 3H) ppm. Electrospray MS: *m/z* 180 [M + 1].


3. Preparation of 3-ethyl-2-{[2-(6-fluoropyridin-2-yl)-1*H*-imidazol-1-yl]methyl}-6-propyl-3*H*-imidazo[4,5-*c*]pyridine

25 A solution of *N*³-ethyl-6-propyl-pyridine-3,4-diamine (82 mg, ~0.45 mmol, contaminated with ~10% of *N*-(4-amino-6-propyl-pyridin-3-yl)-acetamide) in 1,2-dichloroethane (4 mL) at room temperature under N₂ is treated with 2.0 M AlMe₃ (0.69 mL, 1.37 mmol). The resulting solution is stirred at room temperature for 30 min and then treated with a solution of [2-(6-fluoro-pyridin-
30 2-yl)-imidazol-1-yl]-acetic acid methyl ester (97 mg, 0.41 mmol) in 1,2-dichloroethane via cannula, followed by a 1 mL rinse. The reaction mixture is then stirred at reflux overnight. After cooling, it is treated with MeOH (~2 mL) and then further diluted

with saturated aq NaHCO₃ (~10 mL) and some H₂O. The mixture is stirred vigorously for 15 min and then extracted thrice with CHCl₃ containing some (~5%) MeOH. The combined extracts are dried over K₂CO₃ and concentrated. The crude material is purified by preparative thin layer chromatography. Developing with 15:1 CHCl₃-MeOH yields 3-ethyl-2-{[2-(6-fluoropyridin-2-yl)-1H-imidazol-1-yl]methyl}-6-propyl-3H-imidazo[4,5-c]pyridine as an off-white solid. ¹H NMR (CDCl₃, 300 MHz) δ 8.70 (d, J = 0.9 Hz, 1H), 8.17 (dd, J = 8.0, 2.6 Hz, 1H), 7.89 (q, J = 7.8 Hz, 1H), 7.47 (s, 1H), 7.22 (d, J = 0.9 Hz, 1H), 7.18 (d, J = 1.5 Hz, 1H), 6.88 (dd, J = 8.6, 2.9 Hz, 1H), 6.27 (s, 2H), 4.42 (q, J = 7.2 Hz, 2H), 2.86 (m, 2H), 1.78 (sextet, J = 7.2 Hz, 2H), 1.30 (t, J = 7.2 Hz, 3H), 0.97 (t, J = 7.2 Hz, 3H) ppm. Electrospray MS: m/z 365 [M + 1].

Example 32

Synthesis of 1-ethyl-2-{[2-(1,3-thiazol-2-yl)-1H-imidazol-1-yl]methyl}-6-(trifluoromethyl)-1H-imidazo[4,5-c]pyridine



1. Preparation of 5-Nitro-2-trifluoromethyl-pyridin-4-ol

25 mL of fuming H₂SO₄ under cold conditions (0 °C) is added dropwise to a stirred solution of 4-hydroxy-2-trifluoromethylpyridine [Chemistry of Heterocyclic Compounds 1997, 33, 995-996] (4g, 24.5 mmol) in concentrated H₂SO₄ (10 mL). HNO₃ (fuming, 90%, 25 mL) is added carefully to the above mixture with caution to keep the offset of any exotherm under control, and the reaction is allowed to warm to room temperature slowly. After heating at 120 °C for 6 h with stirring, the resulting mixture is cooled to room temperature, poured into ice-cold water

maintaining the pH around 1 with the addition of 10N NaOH solution, and then extracted with CHCl₃. The combined organic phase is washed successively with saturated aqueous NaHCO₃, water, brine, dried over Na₂SO₄, and concentrated in vacuo to afford 5-nitro-2-trifluoromethyl-pyridin-4-ol as an yellow solid. ¹H NMR (300 MHz, CDCl₃): δ 9.33 (s, 1H), 7.50 (s, 1H).

2. Preparation of 4-Chloro-5-nitro-2-trifluoromethyl-pyridine

The crude 5-nitro-2-trifluoromethyl-pyridin-4-ol (3.4 g, 16.3 mmol) was treated with PCl₅ (5.2 g, 24.97 mmol) and POCl₃ (7 mL, 24.49 mmol). The resulting mixture is heated at 80 °C for 18 h, cooled and poured into ice-cold water, and extracted with CH₂Cl₂. The combined organic phase is washed successively with saturated aqueous NaHCO₃, water, brine, dried over Na₂SO₄, and concentrated in vacuo to afford 4-Chloro-5-nitro-2-trifluoromethyl-pyridine as an yellow oil. ¹H NMR (300 MHz, CDCl₃): δ 9.17 (s, 1H), 7.9 (s, 1H).

3. Preparation of Ethyl-(5-nitro-2-trifluoromethyl-pyridin-4-yl)-amine

K₂CO₃ (4.15 g, 30.01 mmol), and 2.0 M solution of EtNH₂ (15 mL, 30.01mmol) in THF are added o a solution of 4-chloro-5-nitro-2-trifluoromethyl-pyridine (3.4 g, 15.0 mmol) in acetonitrile (30 mL) under cold conditions . The reaction is complete after stirring at 0°C for 2h. Usual work up affords ethyl-(5-nitro-2-trifluoromethyl-pyridin-4-yl)-amine as an yellow-orange viscous oil. ¹H NMR (300 MHz, CDCl₃): δ 9.24 (s, 1H), 7.09 (s, 1H), 3.45 (t, J = 7.2 Hz, 2H), 1.42 (t, J = 7.2 Hz, 3H).

4. Preparation of N⁴-Ethyl-6-trifluoromethyl-pyridine-3,4-diamine

Ethyl-(5-nitro-2-trifluoromethyl-pyridin-4-yl)-amine is hydrogenated under usual conditions to afford *N*⁴-ethyl-6-trifluoromethyl-pyridine-3,4-diamine. ¹H NMR (300 MHz, CD₃OD): δ 7.71 (s, 1H), 6.75 (s, 1H), 3.28 (t, *J* = 6.9 Hz, 2H), 1.31 (t, *J* = 6.9 Hz, 3H).

5. Preparation of 2-Chloromethyl-1-ethyl-6-trifluoromethyl-1*H*-imidazo[4,5-*c*]pyridine

The HCl salt of ethyl 2-chloro-acetimidate (3.90 g, 24.58 mmol) is added to a solution of *N*⁴-ethyl-6-trifluoromethyl-pyridine-3,4-diamine (1.44 g, 7.02 mmol) in EtOH (20 mL) and refluxed for 3 h. The reaction mixture is cooled to room temperature and concentrated under reduced pressure. The residue is diluted with CH₂Cl₂, washed with NaHCO₃ solution, dried over Na₂SO₄, and concentrated to obtain 2-chloromethyl-1-ethyl-6-trifluoromethyl-1*H*-imidazo[4,5-*c*]pyridine. ¹H NMR (300 MHz, CDCl₃): δ 9.14 (s, 1H), 7.66 (s, 1H), 7.26 (s, 1H), 4.87 (s, 2H), 4.40 (t, *J* = 7.5 Hz, 2H), 1.58 (t, *J* = 7.5 Hz, 3H).


6. Preparation of 1-ethyl-2-{[2-(1,3-thiazol-2-yl)-1*H*-imidazol-1-yl]methyl}-6-(trifluoromethyl)-1*H*-imidazo[4,5-*c*]pyridine

A solution of 2-Chloromethyl-1-ethyl-6-trifluoromethyl-1*H*-imidazo[4,5-*c*]pyridine (1 g, 3.8 mmol) in anhydrous DMF (15 mL) were added K₂CO₃ (2.68 g, 19 mmol), and 2-(1*H*-imidazol-2-yl)-thiazole (555mg, 3.8 mmol) and stirred at room temperature for 48 h. The crude residue after usual work up was purified eluting with 5% MeOH-CH₂Cl₂ containing few drops of NH₄OH to afford 1-ethyl-2-{[2-(1,3-thiazol-2-yl)-1*H*-imidazol-1-yl]methyl}-6-(trifluoromethyl)-1*H*-imidazo[4,5-*c*]pyridine (800 mg, 80 %); ¹H NMR (CD₃OD): δ 8.83 (s, 1H), 8.14 (s, 1H), 7.71 (d, *J* = 2.7 Hz, 1H), 7.53 (d, *J* = 2.7 Hz, 1H), 7.45 (d, *J* = 0.9 Hz, 1H), 7.19 (d, *J* = 0.9 Hz, 1H), 6.30 (s, 2H), 4.54 (q, *J* = 5.4 Hz, 2H), 1.46 (t, *J* = 5.4 Hz, 3H); *m/z* 379 [M+1]

Example 33

Synthesis of 3-ethyl-2-{[2-(1,3-thiazol-2-yl)-1H-imidazol-1-yl]methyl}-6-(trifluoromethyl)-3H-imidazo[4,5-c]pyridine

5



1. Preparation of 5-Nitro-2-trifluoromethyl-pyridin-4-ylamine

Ammonia gas is bubbled through a solution of 2-trifluoromethyl-4-chloro-5-nitro-pyridine (4.12 g, 18.23 mmol) in anhydrous THF at room temperature for 3 h. Removal of solvent under reduced pressure affords 5-nitro-2-trifluoromethyl-pyridin-4-ylamine (3.6 g, 96%); ¹H NMR (300 MHz, CDCl₃): δ 7.15 (s, 1H), 9.15 (s, 1H).

2. Preparation of 6-Trifluoromethyl-pyridine-3,4-diamine

Hydrogenation as described in the previous examples provides 6-trifluoromethyl-pyridine-3,4-diamine. ¹H NMR (300 MHz, CDCl₃): δ 6.95 (s, 1H), 8.01 (s, 1H).

3. Preparation of N³-ethyl-6-trifluoromethyl-pyridine-3,4-diamine

Cesium carbonate (3.1 g, 26.88 mmol) and ethyl iodide (1.76 g, 10.75 mmol) is added to a solution of 6-trifluoromethyl-pyridine-3,4-diamine (1.05 g, 8.96 mmol) in anhydrous DMF (12 mL). The mixture is heated in a sealed tube for 18 h, then cooled to room temperature, and diluted with EtOAc and water. Usual work up and purification by column chromatography, eluting with EtOAc-hexanes provides pure N³-Ethyl-6-trifluoromethyl-pyridine-3,4-

diamine. ^1H NMR (300 MHz, CDCl_3): δ 7.95 (s, 1H); 6.95 (s, 1H), 3.21 (q, J = 5.4 Hz, 2H), 1.35 (t, J = 5.4 Hz, 3H).

4. Preparation of 2-Chloromethyl-3-ethyl-6-trifluoromethyl-3H-imidazo[4,5-c]pyridine


2-Chloromethyl-3-ethyl-6-trifluoromethyl-3H-imidazo[4,5-c]pyridine is made from N^3 -Ethyl 6-Trifluoromethyl-pyridine-3,4-diamine following the procedure described previously for the preparation of 2-Chloromethyl-1-ethyl-6-trifluoromethyl-1H-imidazo[4,5-c]pyridine. ^1H NMR (300 MHz, CDCl_3): δ 8.89 (s, 1H), 8.04 (s, 1H), 4.87 (s, 2H), 4.45 (t, J = 7.2 Hz, 2H), 1.59 (t, J = 7.2 Hz, 3H)

5. Preparation of 3-ethyl-2-([2-(1,3-thiazol-2-yl)-1H-imidazol-1-yl]methyl)-6-(trifluoromethyl)-3H-imidazo[4,5-c]pyridine

As described previously for the preparation of 1-Ethyl-2-(2-thiazol-2-yl-imidazol-1-ylmethyl)-6-trifluoromethyl-1H-imidazo[4,5-c]pyridine, nucleophilic displacement of the 2-chloromethyl-3-ethyl-6-trifluoromethyl-3H-imidazo[4,5-c]pyridine with the same 2-(1H-imidazol-2-yl)-thiazole followed by usual work provides 3-ethyl-2-([2-(1,3-thiazol-2-yl)-1H-imidazol-1-yl]methyl)-6-(trifluoromethyl)-3H-imidazo[4,5-c]pyridine. ^1H NMR (CD_3OD) δ 9.02 (s, 1H), 7.89 (s, 1H), 7.70 (d, J = 3 Hz, 1H), 7.52 (d, J = 3.3 Hz, 1H), 7.46 (d, J = 1.5 Hz, 1H), 7.19 (d, J = 1.5 Hz, 1H), 6.29 (s, 2H), 4.56 (q, J = 7.5 Hz, 2H), 1.52 (t, J = 7.5 Hz, 3H) m/z 379 $[\text{M}+1]$

Example 34

Synthesis of 3-ethyl-2-[(2-pyrimidin-2-yl-1H-imidazol-1-yl)methyl]-3H-imidazo[4,5-c]pyridine-4-carbonitrile



1. Preparation of 3-Ethylamino-4-nitro-pyridine-2-carbonitrile

TMSCN (1.1 mL, 8.33 mmol) and dimethyl amino carbamoyl
 5 chloride (0.76 mL, 8.33 mmol) are added to a solution of 3-ethylamino-4-nitro-pyridine-1-oxide (1 g, 6.94 mmol) in anhydrous DMF (10 mL) were added. The reaction mixture is stirred overnight at 80 °C, cooled to room temperature, and diluted with CH₂Cl₂ and water. The combined organic layer is washed with water, brine,
 10 dried over Na₂SO₄, and concentrated. The residue is chromatographed eluting with 40% EtOAc-hexanes to afford 3-ethylamino-4-nitro-pyridine-2-carbonitrile as a yellow solid. ¹H NMR (300 MHz, CD₃OD): δ 7.82 (d, J = 4.5 Hz, 1H), 7.50 (d, J = 4.5 Hz, 1H), 3.70 (q, J = 7.2 Hz, 2H), 1.39 (t, J = 7.2 Hz, 3H).

2. Preparation of 4-Amino-3-ethylamino-pyridine-2-carbonitrile

Hydrogenation of 3-Ethylamino-4-nitro-pyridine-2-carbonitrile under conditions previously described provides 4-Amino-3-ethylamino-pyridine-2-carbonitrile. ¹H NMR (300 MHz, CD₃OD): δ
 20 7.75 (d, J = 5.4 Hz, 1H), 6.66 (d, J = 5.4 Hz, 1H), 3.35 (q, J = 7.2 Hz, 2H), 1.29 (t, J = 7.2 Hz, 3H).

3. Preparation of 2-Chloromethyl-3-ethyl-3H-imidazo[4,5-c]pyridine-4-carbonitrile


25 2-Chloromethyl-3-ethyl-3H-imidazo[4,5-c]pyridine-4-carbonitrile is prepared from the reaction of 4-amino-3-ethylamino-pyridine-2-carbonitrile and 2-chloro acetimidate following the previously described procedure. ¹H NMR (300 MHz, CDCl₃): δ 8.55 (d, J = 5.4 Hz, 1H), 7.84 (d, J = 5.4 Hz, 1H),
 30 4.98 (s, 2H), 3.41 (q, J = 7.5 Hz, 2H), 1.23 (t, J = 7.5 Hz, 3H).

4. Preparation of 3-ethyl-2-[(2-pyrimidin-2-yl-1H-imidazol-1-yl)methyl]-3H-imidazo[4,5-c]pyridine-4-carbonitrile

3-ethyl-2-[(2-pyrimidin-2-yl-1H-imidazol-1-yl)methyl]-3H-imidazo[4,5-c]pyridine-4-carbonitrile is obtained from 2-chloromethyl-3-ethyl-3H-imidazo[4,5-c]pyridine-4-carbonitrile by nucleophilic displacement with 2-(1H-Imidazol-2-yl)-pyrimidine, as described previously, followed by usual work up. ¹H NMR (300 MHz, CD₃OD): δ 8.70 (d, *J* = 5.1 Hz, 2H), 8.41 (d, *J* = 5.4 Hz, 1H), 7.73 (d, *J* = 5.1 Hz, 1H), 7.52 (d, *J* = 1.2 Hz, 1H), 7.31 (d, *J* = 1.2 Hz, 1H), 7.27 (t, *J* = 4.5 Hz, 1H), 6.35 (s, 2H), 4.75 (q, *J* = 7.5 Hz, 2H), 1.63 (t, *J* = 7.2 Hz, 3H); *m/z* 331 [M+1].

Example 35

Synthesis of 1-(2-{[2-(6-fluoropyridin-2-yl)-1H-imidazol-1-yl]methyl}-1-propyl-1H-imidazo[4,5-c]pyridin-4-yl)ethanone




To the mixture of H₂O (5 mL) and CH₂Cl₂ (5 mL) is added 1-Propyl-2-{[2-(2-fluoropyridin-6-yl)-1H-imidazol-1-yl]methyl}-5-aza-1H-benzimidazole (168 mg, 0.5 mmol), pyruvic acid (132 mg, 1.5 mmol), silver nitrate (7 mg, 0.04 mmol), (NH₄)₂S₂O₈ (342 mg, 1.5 mmol), and sulfuric acid (98%, 100 mg, 1.0 mmol). The mixture is heated to 40 °C for 2 hr, then cooled to room temperature. The aqueous solution is neutralized to pH 8 with saturated sodium bicarbonate solution and extracted with CH₂Cl₂. The combined organic layers are dried over Na₂SO₄ and solvent is removed to give a brown solid. Purification by preparative TLC provides '1-

(2-{[2-(6-fluoropyridin-2-yl)-1H-imidazol-1-yl]methyl}-1-propyl-1H-imidazo[4,5-c]pyridin-4-yl)ethanone as white solid. ¹H NMR (CDCl₃) δ 8.48 (d, 1H), 8.12 (dd, 1H), 7.88 (q, 1H), 7.44 (d, 1H), 7.23 (d, 1H), 7.11 (d, 1H), 6.88 (dd, 1H), 6.40 (s, 2H), 4.22 (q, 2H), 2.84 (s, 3H), 1.58 (m, 2H), 0.74 (t, 3H).

Example 36

Synthesis of 1-(3-ethyl-2-{[2-(1,3-thiazol-2-yl)-1H-imidazol-1-yl]methyl}-3H-imidazo[4,5-c]pyridin-6-yl)ethanone



1. Preparation of 2,5-Dichloro-pyridine 1-oxide

A mixture of 2,5-Dichloropyridine (30g, 0.2 mmol) and 3-Chloroperoxybenzoic acid (70%, 50g, 0.2 mmol) in 500 mL of Dichloroethane is heated at 50 °C overnight. After cooling to -30 °C, NH₃ gas is bubbled into the reaction mixture for 5 minutes. The mixture is filtered, the solid is washed with dichloroethane (50 mL), and the filtrate is concentrated to a solid. The solid is mixed with 200 mL of Hexane and heated at reflux for 30 minutes. After cooling, the precipitate is collected by filtration, and dried to give the titled compound as a tan solid. ¹H NMR (CDCl₃): δ 8.40 (d, 1H), 7.44 (d, 1H), 7.22 (t, 1H).

2. Preparation of 2,5-dichloro-4-nitro-pyridine 1-oxide

At 0°C, 30% oleum (13 mL) is added slowly to fuming HNO₃ (22 mL) and the resulting mixture is added slowly to the solution of

2,5-dichloro-pyridine 1-oxide (5.2 g, 31 mmol) in concentrated H₂SO₄. After addition, the reaction mixture is warmed to room temperature over 1 hr, then heated to 80 °C for 2 hr. After cooling to room temperature, the mixture is poured into ice (120 g) and precipitate is formed gradually. The solid is collected by filtration and washed with water. The product is obtained as yellow solid. ¹H NMR (CDCl₃) δ 8.46 (s, 1H), 8.25 (s, 1H).

3. Preparation of (6-Chloro-4-nitro-1-oxy-pyridin-3-yl)-ethylamine

To a solution of 2,5-Dichloro-4-nitro-pyridine 1-oxide (7 g, 33mmol) in 100 mL of THF is added a solution of Ethylamine (2M, 50 mL) dropwise at °C. The reaction mixture is warmed to room temperature and stirred overnight. The volatiles are evaporated in vacuo. The residue is purified on a silica gel column with 1% CH₂Cl₂ as eluent to give the title compound as an orange solid. ¹H NMR (CDCl₃): δ 8.24 (s, 1H), 8.06 (s, 1H), 7.62 (br, 1H), 3.25 (q, 2H), 1.38 (t, 3H).

4. Preparation of 1-(5-ethylamino-4-nitro-pyridin-2-yl)-ethanone

(6-Chloro-4-nitro-1-oxy-pyridin-3-yl)ethyl-amine (700 mg, 3.2 mmol), tributyl-(1-ethoxy-vinyl)-stannane (1.75 g, 4.9 mmol), PdCl₂(PPh₃)₂ (350 mg, 0.5 mmol), and toluene (35 mL) are added to a sealed tube; the mixture is degassed for 30 min. The reaction is heated to 75 °C for 16 hr and then cooled to room temperature. 6N HCl solution (20 mL) is added to the mixture and the reaction is stirred at room temperature for 1 hr. The reaction mixture is filtered through celite and the aqueous layer is neutralized to basic with saturated NaHCO₃. The aqueous layer is extracted with ethyl acetate and the combined organic layers are dried over Na₂SO₄. The solvent is removed in vacuo. Purification by flash column provides the product as orange solid. ¹H NMR

(CDCl₃) δ 8.70 (s, 1H), 8.48 (s, 1H), 7.96 (br, 1H), 3.55 (m, 2H), 2.65 (s, 3H), 0.92 (t, 3H). LRMS 210.1 (MH⁺).

5. Preparation of 1-(4-amino-5-ethylamino-pyridin-2-yl)-ethanone


5 A mixture of 1-(5-ethylamino-4-nitro-pyridin-2-yl)-ethanone (310 mg, 1.5 mmol) and 10% Pd/C (50 mg) in ethanol (10 mL) is stirred under hydrogen for 1 hr. The mixture is filtered through Celite and concentrated in vacuo to give the product as yellow solid. ¹H NMR (CD₃OD) δ 7.61 (s, 1H), 7.30 (s, 1H), 3.26 (q, 2H),
10 2.53 (s, 3H), 0.90 (t, 3H). LRMS 180.5 (MH⁺).

6. Preparation of 1-(3-ethyl-2-{[2-(1,3-thiazol-2-yl)-1H-imidazol-1-yl]methyl}-3H-imidazo[4,5-c]pyridin-6-yl)ethanone

A mixture of 1-(4-amino-5-ethylamino-pyridin-2-yl)-ethanone
15 (54 mg, 0.3 mmol), (2-thiazol-2-yl-imidazol-1-yl)-acetic acid (61 mg, 0.3 mmol), EDCI (60 mg, 0.3 mmol), and pyridine (2 mL) is stirred at room temperature for 72 hr and then poured into water. The aqueous layer is extracted with ethyl acetate and the combined organic layers are dried over Na₂SO₄. Solvent is
20 evaporated in vacuo and the residue is taken up in acetic acid (5 mL). The mixture is heated to reflux for 16 hr and saturated NaHCO₃ solution is added to neutralize the reaction. The aqueous solution is extracted with ethyl acetate and the combined organic layers are dried over Na₂SO₄. Solvent is removed in vacuo and
25 purification by preparative TLC provides the product as white solid. ¹H NMR (CDCl₃) δ 8.79 (d, 2H), 8.47 (d, 1H), 7.83 (d, 1H), 7.40 (d, 1H), 7.19 (d, 1H), 7.15 (d, 1H), 6.39 (s, 2H), 4.02 (q, 2H), 2.77 (s, 3H), 1.86 (t, 3H). LRMS 353.2 (MH⁺).

30 **Example 37**

Synthesis of 1-Ethyl-2-(2-thiazol-2-yl-2H-pyrazol-3-ylmethyl)-1H-imidazo[4,5-c]pyridine



1. Preparation of (2-Thiazol-2-yl-2H-pyrazol-3-yl)-acetic acid ethyl ester

A mixture of thiazol-2-ylhydrazine (2.2 g, 19.1 mmol) and 3,5,5-triethoxy-pent-2-enoic acid ethyl ester (70% purity, 7.1 g, 19.1 mmol) is refluxed in acetic acid overnight. Acetic acid is removed. To the residue is added NaHCO₃ (aq.) (40 mL) and ethyl acetate (100 mL). The insoluble material is filtered. The organic layer is dried and solvent removed. The crude is subjected to column separation (hexane/ethyl acetate 3:1) to give (2-thiazol-2-yl-2H-pyrazol-3-yl)-acetic acid ethyl ester as an oil.

2. Preparation of 1-Ethyl-2-(2-thiazol-2-yl-2H-pyrazol-3-ylmethyl)-1H-imidazo[4,5-c]pyridine


Trimethylaluminum (2M in toluene) (1.4 mL, 2.85 mmol) is added dropwise under N₂ to a solution of N⁴-Ethyl-pyridine-3,4-diamine (156 mg, 1.14 mmol) in DCM (5 mL). The mixture is stirred at room temperature for 1 hour. A solution of thiazol-2-yl-2H-pyrazol-3-yl)-acetic acid ethyl ester (270 mg, 1.14 mmol) in DCM (2 mL) is added. The mixture is refluxed for 25 hours. On cooling, the reaction is quenched with water added dropwise and DCM (20 mL) is added. The organic layer is separated and the aqueous layer is extracted with DCM (3 x 30 mL). The combined organic layers are dried and solvent removed. Acetic acid (10 mL) is added to the residue. The solution is refluxed for 3.5 h. Acetic acid is removed in vacuo and the residue is purified by PTLC (10% methanol in DCM) to give 1-Ethyl-2-(2-thiazol-2-yl-2H-pyrazol-3-ylmethyl)-1H-imidazo[4,5-c]pyridine as a solid.

¹H NMR (CDCl₃) δ 9.00 (s, 1H), 8.43 (d, 1H), 7.63 (d, 1H), 7.43 (d, 1H),

7.15 (d, 1H), 7.06 (d, 1H), 6.25 (d, 1H), 4.99 (s, 2H), 4.25 (q, 2H), 1.36 (t, 3H).

Example 38

5 Synthesis of 1-Ethyl-2-[1-(2-thiazol-2-yl-2H-pyrazol-3-yl)-ethyl]-1H-imidazo[4,5-c]pyridine



10 1. Preparation of 2-(2-Thiazol-2-yl-2H-pyrazol-3-yl)-propionic acid ethyl ester

15 A mixture of (2-thiazol-2-yl-2H-pyrazol-3-yl)-acetic acid ethyl ester (100 mg, 0.42 mmol), methyl iodide (66 mg, 0.46 mmol) and cesium carbonate (151 mg, 0.46 mmol) in DMF (6 mL) is stirred at room temperature overnight. The reaction is quenched with
20 NH₄Cl (aq.) (6 mL) and ethyl acetate (30 mL). The organic layer is separated and the aqueous layer is extracted with ethyl acetate (3 x 30 mL). The combined organic layers are dried and the solvent removed. The crude is purified by PTLC (hexane/ethyl acetate 3:1) to give 2-(2-thiazol-2-yl-2H-pyrazol-3-yl)-propionic acid ethyl ester as an oil.


2. Preparation of 1-Ethyl-2-[1-(2-thiazol-2-yl-2H-pyrazol-3-yl)-ethyl]-1H-imidazo[4,5-c]pyridine

25 Trimethylaluminum (2M in toluene) (0.5 mL, 1 mmol) is added dropwise under N₂ to a solution of N⁴-Ethyl-pyridine-3,4-diamine (44 mg, 0.32 mmol) in DCM (5 mL). The mixture is stirred at room temperature for 1 hour. A solution of thiazol-2-yl-2H-pyrazol-3-yl)-propionic acid ethyl ester in DCM (2 mL) is added. The mixture is relaxed for 3 days. On cooling, the reaction is
30 quenched with water added dropwise and DCM (20 mL) is added. The organic layer is separated and the aqueous layer is extracted

with DCM (3 x 30 mL). The combined organic layers are dried and the solvent removed. The residue is purified by PTLC (10% methanol in DCM) to give 1-ethyl-2-[1-(2-thiazol-2-yl-2H-pyrazol-3-yl)-ethyl]-1H-imidazo[4,5-c]pyridine as a solid. ¹H NMR (CDCl₃) δ 9.05 (s, 1H), 8.39 (d, 1H), 7.59 (d, 1H), 7.49 (d, 1H), 7.25 (d, 1H), 7.09 (d, 1H), 6.28 (d, 1H), 5.85 (q, 1H), 4.19 (q, 2H), 1.85 (d, 3H), 1.29 (t, 3H).

Example 39

Synthesis of 2-[2,4']Bithiazolyl-5'-ylmethyl-1-ethyl-1H-imidazo[4,5-c]pyridine



1. Preparation of 4-Oxo-4-thiazol-2-yl-butyric acid ethyl ester

To a stirred solution of 2-trimethylsilyl-thiazole (2.6 g, 16.53 mmol) in DCM (30 mL) is added a solution of 3-chlorocarbonyl-propionic acid ethyl ester (5.4 g, 32.81 mmol) in DCM (10 mL). The mixture is stirred overnight at room temperature. To the mixture is added 5% NaHCO₃ (30 mL) and mixture is stirred at room temperature for 20 minutes. The organic layer is separated and the aqueous layer is extracted with DCM (2 x 50 mL). The combined organic layers are dried and the solvent removed. The residue is separated by column (methylene chloride) to give 4-oxo-4-thiazol-2-yl-butyric acid ethyl ester.

2. Preparation of 3-Bromo-4-oxo-4-thiazol-2-yl-butyric acid ethyl ester

Phenyltrimethylammonium tribromide (3.4 g, 8.91 mmol) in DCM (20 mL) is added dropwise to a solution of 4-oxo-4-thiazol-2-yl-butyric acid ethyl ester (1.9 g, 8.91 mmol) in DCM (30 mL). The mixture is stirred overnight at room temperature. 5% NaHCO₃ (40

mL) is added to the mixture. The organic layer is separated and the aqueous layer is extracted with DCM (2 x 40 mL). The combined organic layers are dried and solvent removed. The residue is purified by column (hexanes/ethyl acetate 3:1) to give 3-bromo-4-oxo-4-thiazol-2-yl-butyric acid ethyl ester.

3. Preparation of [2,4']Bithiazolyl-5'-yl-acetic acid ethyl ester

To a solution of 3-bromo-4-oxo-4-thiazol-2-yl-butyric acid ethyl ester (2.2 g, 7.53 mmol) in 1,4-dioxane (40 mL) is added thioformamide (freshly made from formamide and P₂S₅ in 1,4-dioxane) (22.59 mmol). The mixture is refluxed overnight. The solvent is removed and NaHCO₃ (aq.) (40 mL) and DCM (100 mL) are added to the residue. The organic layer is separated and the aqueous layer is extracted with DCM (2 x 40 mL). The combined organic layers are dried and solvent removed. The crude is purified by column (1% methanol in DCM) to give [2,4']bithiazolyl-5'-yl-acetic acid ethyl ester.


4. Preparation of 2-[2,4']Bithiazolyl-5'-ylmethyl-1-ethyl-1H-imidazo[4,5-c]pyridine

Trimethylaluminum (2M in toluene) (2.2 mL, 4.4 mmol) is added dropwise under N₂ to a solution of N⁴-Ethyl-pyridine-3,4-diamine (237 mg, 1.73 mmol) in DCM (5 mL). The mixture is stirred at room temperature for 1 hour. A solution of [2,4']bithiazolyl-5'-yl-acetic acid ethyl ester (440 mg, 1.73 mmol) in DCM (2 mL) is added. The mixture is relaxed for 3 days. On cooling, the reaction is quenched with water added dropwise and DCM (40 mL) is added. The organic layer is separated and the aqueous layer is extracted with DCM (3 x 30 mL). The combined organic layers are dried and solvent removed. The residue is purified by PTLC (10% methanol in DCM) to give 2-[2,4']bithiazolyl-5'-ylmethyl-1-ethyl-1H-imidazo[4,5-c]pyridine as a solid. ¹H NMR (CDCl₃) δ 9.05 (s,

1H), 8.76 (s, 1H), 8.41 (d, 1H), 7.88 (d, 1H), 7.41 (d, 1H), 7.25 (d, 1H), 5.35 (s, 2H), 4.26 (q, 2H), 1.25 (t, 3H).

Example 40

5 Synthesis of 5-Bromo-1-ethyl-2-[3-(3-fluoro-phenyl)-isoxazol-4-ylmethyl]-1H-benzoimidazole



1. Preparation of 3-(5-Bromo-1-ethyl-1H-benzoimidazol-2-yl)-1-(3-fluoro-phenyl)-propan-1-one

To a solution of 1-(3-fluoro-phenyl)-ethanone (0.55 g, 4.0 mmol) in THF (30 mL) at -78°C under N₂ is added LDA (2M in hexanes, 2.0 mL, 4.0 mmol). After 10 minutes, 5-bromo-2-chloromethyl-1-ethyl-1H-benzoimidazole (1 g, 3.65 mmol) in THF (5 mL) is added. The mixture is stirred at this temperature for an additional 1 hour and is gradually warmed to room temperature. The reaction is then quenched with saturated NH₄Cl. The mixture is extracted with ethyl acetate (3 x 50 mL). Upon drying, the organic solvent is removed to give 3-(5-Bromo-1-ethyl-1H-benzoimidazol-2-yl)-1-(3-fluoro-phenyl)-propan-1-one as a yellow oil.


2. Preparation of 5-Bromo-1-ethyl-2-[3-(3-fluoro-phenyl)-isoxazol-4-ylmethyl]-1H-benzoimidazole

A mixture of 3-(5-Bromo-1-ethyl-1H-benzoimidazol-2-yl)-1-(3-fluoro-phenyl)-propan-1-one (0.1 g, 0.27 mmol) and tris(dimethylamino)methane (0.077g, 0.54 mmol) is heated at 60 °C in a sealed tube for 6 hours. The volatile material is removed in vacuo. To the residue is added EtOH (5 mL) and hydroxylamine (1.1 mmol). The mixture is heated at 120 °C for 2 hours. The solvent is removed. To the residue is added NaHCO₃ (aq.) (10 mL) and DCM

(30 mL). The organic layer is separated and the aqueous layer is extracted with DCM (2 x 15 mL). The combined organic layers are dried and the solvent removed. The crude is purified by PTLC (ethyl acetate/Hexanes 1:1) to give 5-Bromo-1-ethyl-2-[3-(3-fluoro-phenyl)-isoxazol-4-ylmethyl]-1H-benzoimidazole as a solid. ¹H NMR (CDCl₃) δ 8.40 (s, 1H), 7.88 (s, 1H), 7.31-7.52 (m, 4H), 7.18-7.22 (m, 2H), 4.14 (s, 2H), 4.04 (q, 2H), 1.18 (t, 3H).

Example 41

Synthesis of 5-Bromo-1-ethyl-2-[3-(3-fluoro-phenyl)-1H-pyrazol-4-ylmethyl]-1H-benzoimidazole





A mixture of 3-(5-Bromo-1-ethyl-1H-benzimidazol-2-yl)-1-(3-fluoro-phenyl)-propan-1-one (0.1 g, 0.27 mmol) and tris(dimethylamino)methane (0.077g, 0.54 mmol) is heated at 60 °C in a sealed tube for 6 hours. The volatile material is removed in vacuo. EtOH (5 mL) and hydrazine acetate (1.1 mmol) are added to the residue. The mixture is heated at 120 °C for 2 hours. The solvent is removed. NaHCO₃ (aq.) (10 mL) and DCM (30 mL) are added to the residue. The organic layer is separated and the aqueous layer is extracted with DCM (2 x 15 mL). The combined organic layers are dried and the solvent removed. The crude is purified by PTLC (ethyl acetate/Hexanes 1:1) to give 5-bromo-1-ethyl-2-[3-(3-fluoro-phenyl)-1H-pyrazol-4-ylmethyl]-1H-benzoimidazole as a solid. LRMS: calcd 399.3, found [M+1] 401.0.

Example 42

Synthesis of 5-Bromo-1-ethyl-2-[3-(3-fluoro-phenyl)-1-methyl-1H-pyrazol-4-ylmethyl]-1H-benzoimidazole and 5-Bromo-1-ethyl-2-[5-

(3-fluoro-phenyl)-1-methyl-1H-pyrazol-4-ylmethyl]-1H-
benzoimidazole




5

A mixture of 3-(5-Bromo-1-ethyl-1H-benzoimidazol-2-yl)-1-(3-fluoro-phenyl)-propan-1-one (0.1 g, 0.27 mmol) and tris(dimethylamino)methane (0.077g, 0.54 mmol) is heated at 60 °C in a sealed tube for 6 hours. The volatile material is removed in vacuo. EtOH (5 mL) and methyl hydrazine (1.1 mmol) are added to the residue. The mixture is heated at 120 °C for 2 hours. The solvent is removed. NaHCO₃ (aq.) (10 mL) and DCM (30 mL) are added to the residue. The organic layer is separated and the aqueous layer is extracted with DCM (2 x 15 mL). The combined organic layers are dried and the solvent removed. The crude is purified by PTLC (ethyl acetate/Hexanes 1:1) to give 5-bromo-1-ethyl-2-[3-(3-fluoro-phenyl)-1-methyl-1H-pyrazol-4-ylmethyl]-1H-benzoimidazole and 5-bromo-1-ethyl-2-[5-(3-fluoro-phenyl)-1-methyl-1H-pyrazol-4-ylmethyl]-1H-benzoimidazole. 5-bromo-1-ethyl-2-[3-(3-fluoro-phenyl)-1-methyl-1H-pyrazol-4-ylmethyl]-1H-benzoimidazole: ¹H NMR (CDCl₃) δ 7.88 (s, 1H), 7.27-7.42 (m, 4H), 7.00-7.20 (m, 3H), 4.21 (s, 2H), 3.95 (t, 2H), 3.83 (s, 3H), 1.15 (t, 3H). 5-bromo-1-ethyl-2-[5-(3-fluoro-phenyl)-1-methyl-1H-pyrazol-4-ylmethyl]-1H-benzoimidazole: ¹H NMR (CDCl₃) δ 7.81 (s, 1H), 7.38-7.45 (m, 2H), 7.32 (dd, 1H), 7.00-7.18 (m, 4H), 4.00 (s, 2H), 3.90 (q, 2H), 3.78 (s, 3H), 1.15 (t, 3H).

Example 43

Synthesis of 5-Bromo-1-ethyl-2-[4-(3-fluoro-phenyl)-thiazol-5-ylmethyl]-1H-benzoimidazole




5

10
15
20
A mixture of 3-(5-bromo-1-ethyl-1H-benzoimidazol-2-yl)-1-(3-fluoro-phenyl)-propan-1-one (0.2 g, 0.53 mmol) and bromine (0.1 g, 0.62 mmol) is refluxed in 1,4-dioxane for 2 hours. Thioformamide (freshly made from formamide and P₂S₅ in 1,4-dioxane) (1.3 mmol) are added to the solution. The mixture is refluxed overnight. The solvent is removed and NaHCO₃ (aq.) (15 mL) and DCM (40 mL) are added to the residue. The organic layer is separated and the aqueous layer is extracted with DCM (2 x 30 mL). The combined organic layers are dried and solvent removed. The crude is purified by PTLC (5% methanol in DCM) to give 5-bromo-1-ethyl-2-[4-(3-fluoro-phenyl)-thiazol-5-ylmethyl]-1H-benzoimidazole. ¹H NMR (CDCl₃) δ 8.78 (s, 1H), 7.91 (s, 1H), 7.38-7.48 (m, 4H), 7.11-7.19 (m, 2H), 4.59 (s, 2H), 3.90 (q, 2H), 1.12 (t, 3H).

Example 44

Synthesis of 1-[3-Ethyl-2-(2-thiazol-2-yl-imidazol-1-ylmethyl)-3H-benzoimidazol-5-yl]-ethanone



25

1. Preparation of (5-Chloro-2-nitro-phenyl)-ethyl-amine

A mixture of 4-chloro-2-fluoronitrobenzene (5 g, 28.48 mmol), ethylamine (2M in THF, 24 mL, 48 mmol), and K₂CO₃ (7.9 g, 57 mmol) in acetonitrile (30 mL) is stirred at room temperature overnight. Solvent is removed. Ethyl acetate (80 mL) and brine (40 mL) are added to the residue. The organic layer is separated and the aqueous layer is extracted with ethyl acetate (2 x 50 mL). The combined organic layers are dried and solvent removed to give 5-chloro-2-nitro-phenyl)-ethyl-amine.

10 2. Preparation of 1-(3-Ethylamino-4-nitro-phenyl)-ethanone

A mixture of 5-chloro-2-nitro-phenyl)-ethyl-amine (1 g, 4.98 mmol), tributyl-(1-ethoxyvinyl)stannane (2.7 g, 7.48 mmol) and dichlorobis(triphenylphosphine)palladium (II) (175 mg, 0.25 mmol) in toluene (80 mL) is heated at 100°C in a sealed tube for 5 hours. On cooling, water (30 mL) and HCl (con. 30 mL) are added and the mixture is stirred at room temperature for 2 hours. The mixture is neutralized with NaOH (2 N) and extracted with DCM (3 x 50 mL). The combined organic layers are dried and solvent removed. The residue is purified by column (hexanes / ethyl acetate 3:1) to give 1-(3-ethylamino-4-nitro-phenyl)-ethanone.

3. Preparation of 1-(4-Amino-3-ethylamino-phenyl)-ethanone

10% Pd/C (100 mg) is added to a Parr bottle containing 1-(3-ethylamino-4-nitro-phenyl)-ethanone (0.98 g, 4.7 mmol) in ethanol (40 mL). The Parr bottle is sealed in a mechanical shaker, evacuated, and then purged with nitrogen followed by hydrogen. The system is pressurized to 50 PSI of hydrogen at room temperature and mechanical shaking engaged. After 2 hours, shaking is stopped, and the system purged with nitrogen prior to opening the vessel. The reaction mixture is filtered through celite, concentrated in vacuo, to give 1-(4-amino-3-ethylamino-phenyl)-ethanone.

4. Preparation of 1-(2-Chloromethyl-3-ethyl-3H-benzoimidazol-5-yl)-ethanone


A mixture of 1-(4-amino-3-ethylamino-phenyl)-ethanone (300 mg, 1.69 mmol) and 2-chloroacetamidine hydrochloride (804 mg, 5.1 mmol) in acetonitrile (20 mL) is refluxed for 3 hours. Solvent is removed and the residue is treated with NaHCO₃ (5%, 20 mL) and DCM (50 mL). The organic layer is separated and the aqueous layer is extracted with DCM (2 x 40 mL). The combined organic layers are dried and solvent removed to give 1-(2-Chloromethyl-3-ethyl-3H-benzoimidazol-5-yl)-ethanone.

5. Preparation of 1-[3-Ethyl-2-(2-thiazol-2-yl-imidazol-1-ylmethyl)-3H-benzoimidazol-5-yl]-ethanone

A mixture of 1-(2-Chloromethyl-3-ethyl-3H-benzoimidazol-5-yl)-ethanone (150 mg, 0.63 mmol), 2-(1H-Imidazol-2-yl)-thiazole (95 mg, 0.63 mmol) and K₂CO₃ (262 mg, 1.9 mmol) in DMF (10 mL) is stirred at room temperature for 48 hours. Brine (6 mL) and DCM (20 mL) are added. The organic layer is separated and the aqueous layer is extracted with DCM (3 x 29 mL). The combined organic layers are dried and solvent removed. The residue is purified by PTLC (10% methanol in DCM) to give 1-[3-Ethyl-2-(2-thiazol-2-yl-imidazol-1-ylmethyl)-3H-benzoimidazol-5-yl]-ethanone as a solid. ¹H NMR (CDCl₃) δ 8.05 (d, 1H), 7.92 (dd, 1H), 7.87 (dd, 1H), 7.81 (dd, 1H), 7.40 (d, 1H), 7.20 (d, 1H), 7.14 (d, 1H), 6.17 (s, 2H), 4.33 (q, 2H), 2.67 (s, 3H), 1.13 (t, 3H).

Example 45

Synthesis of 1-Ethyl-2-[2-(1H-pyrazol-3-yl)-imidazol-1-ylmethyl]-1H-benzoimidazole-5-carbonitrile



1. Preparation of 3-(1H-Imidazol-2-yl)-pyrazole-1-carboxylic acid tert-butyl ester


A mixture of 1H-Pyrazole-3-carbaldehyde (1 g, 10.4 mmol),
5 di-tert-butyl dicarbonate (11.4 mmol) and N,N-dimethylpyridine
(20 mg) in DCM (15 mL) and methanol (5 mL) is stirred at room
temperature overnight. Solvent is removed. Glyoxal (6 mL) and
ammonium hydroxid (8 mL) are added to the residue and the mixture
is stirred at room temperature for 1.5 hours. Acetic acid is
10 added dropwise o the mixture to pH ~ 7. Solvent is removed and
DCM (50 mL) and brine (30 mL) are added to the residue. The
organic layer is separated and the aqueous layer is extracted
with DCM (2 x 40 mL). The combined organic layers are dried and
solvent removed to give 3-(1H-imidazol-2-yl)-pyrazole-1-
5 carboxylic acid tert-butyl ester.

2. Preparation of 1-Ethyl-2-[2-(1H-pyrazol-3-yl)-imidazol-1-ylmethyl]-1H-benzoimidazole-5-carbonitrile

20 As described previously, nucleophilic displacement of 2-
Chloromethyl-1-ethyl-1H-benzoimidazole-5-carbonitrile with 3-(1H-
Imidazol-2-yl)-pyrazole-1-carboxylic acid tert-butyl ester
followed by usual work-up provides 1-Ethyl-2-[2-(1H-pyrazol-3-
yl)-imidazol-1-ylmethyl]-1H-benzoimidazole-5-carbonitrile; ¹H NMR
25 (CD₃OD) δ 7.97 (s, 1H), 7.53 (brs, 1H), 7.46 (d, J = 8.7 Hz, 1H),
7.35 (d, J = 8.1 Hz, 1H), 6.97 (d, J = 6.3 Hz, 2H), 6.79 (br s,
1H), 6.10 (s, 2H), 4.22 (q, J = 7.2 Hz, 2H), 0.88 (t, J = 7.2 Hz,
3H); m/z 318 [M+1].

30 **Example 46**

Synthesis of 4-(1-ethyl-2-{[2-(1,3-thiazol-2-yl)-1H-imidazol-1-ylmethyl]-1H-benzimidazol-5-yl)-2-methylbutan-2-ol



1. Preparation of 4-(1-ethyl-2-([2-(1,3-thiazol-2-yl)-1H-imidazol-1-yl]methyl))-1H-benzimidazol-5-yl)-2-methyl-1-but-3-yn-2-ol

Pd (PPh₃)₄ (15 mg, 0.013 mmol), CuI (5 mg, 0.026 mmol), and 2-methyl-but-3-yn-2-ol (250 μ L, 2.6 mmol) are added to the solution of 1-ethyl-5-bromo-2-([2-(1,3-thiazol-2-yl)-1H-imidazol-1-yl]methyl)-1H-benzimidazole (100 mg, 0.26 mmol) in *i*-Pr₂NH (5 mL). The resulting mixture is heated to 100 °C for 20 hr, and then diluted with water. The aqueous layer is extracted with ethyl acetate. The combined organic layers are dried over Na₂SO₄ and solvent is removed to give a brown oil. Purification by preparative TLC provides 40 mg of product as yellow solid. LCMS 392.37 (MH⁺).


2. Preparation of 4-(1-ethyl-2-([2-(1,3-thiazol-2-yl)-1H-imidazol-1-yl]methyl))-1H-benzimidazol-5-yl)-2-methylbutan-2-ol

To a Parr bottle containing 4-(1-ethyl-2-([2-(1,3-thiazol-2-yl)-1H-imidazol-1-yl]methyl))-1H-benzimidazol-5-yl)-2-methyl-1-but-3-yn-2-ol (40 mg, 0.1 mmol) in ethanol (10 mL) is added 10% Pd/C (50 mg). The Parr bottle is sealed in a mechanical shaker, evacuated, and then purged with nitrogen followed by hydrogen. The system is pressurized to 40 PSI of hydrogen at room temperature and mechanical shaking engaged. After 2 hours, shaking is stopped, and the system purged with nitrogen prior to opening the vessel. The reaction mixture is filtered through

celite and concentrated in vacuo. The product is obtained as white solid. ¹H NMR (CDCl₃) δ 7.85 (d, 1H), 7.60 (s, 1H), 7.38 (d, 1H), 7.22 (m, 1H), 7.15 (m, 2H), 7.09 (s, 1H), 6.31 (s, 2H), 4.17 (q, 2H), 2.80 (m, 2H), 1.81 (m, 2H), 1.22 (d, 6H), 1.01 (t, 3H). LCMS 396.38 (MH⁺).

Example 47


Synthesis of 1-(1-ethyl-2-{[2-(1,3-thiazol-2-yl)-1H-imidazol-1-yl]methyl}-1H-benzimidazol-5-yl)-4-hydroxypentan-1-one



To the bi-layer system of 3 N HCl (5 mL) and CH₂Cl₂ is added 5-(1-ethyl-2-{[2-(1,3-thiazol-2-yl)-1H-imidazol-1-yl]methyl}-1H-benzimidazol-5-yl)-pent-4-yn-2-ol (107 mg, 0.27 mmol), PdCl₂ (5 mg, 0.028 mmol), and *n*-Bu₄NCl (17 mg, 0.056 mmol). The resulting mixture is stirred at room temperature for 72 hr, and then poured into saturated NaHCO₃ solution. The aqueous layer is extracted with CH₂Cl₂ and the combined organic layers are dried over Na₂SO₄. Solvent is removed and purification by preparative TLC provides the product as white solid. ¹H NMR (CDCl₃) δ 8.44 (d, 1H), 7.80 (dd, 1H), 7.85 (d, 1H), 7.40 (d, 1H), 7.36 (d, 1H), 7.18 (d, 1H), 7.14 (d, 1H), 6.36 (s, 2H), 4.27 (q, 2H), 3.86 (m, 1H), 3.19 (t, 2H), 1.93 (m, 2H), 1.25 (d, 3H), 1.09 (t, 3H). LCMS 410.2 (MH⁺).

Example 48

Synthesis of 8-{[2-(6-Fluoropyridin-2-yl)-1H-imidazol-1-yl]methyl}-2,9-dimethyl-9H-purine



1) Preparation of 6-Amino-4-chloro-2,N-dimethyl-5-nitropyrimidine

A solution of Methylamine (15.4 mL 1M in THF, diluted with 10 mL of hexane) is added dropwise to a solution of 4,6-Dichloro-2-methyl-5-nitropyrimidine (3.2 g, 15.4 mmol) in Hexane (30 mL). After addition, the reaction mixture is stirred at room temperature for one hour, and concentrated in vacuo to a solid. After the solid is dissolved in 50 mL of CH₂Cl₂, the resultant solution is washed with 0.1 N HCl (20 mL) and water (20 mL), dried over Na₂SO₄ and concentrated to a yellow solid. ¹H NMR (CDCl₃): δ 7.7 (bs, 1H), 3.10 (d, 3H), 2.55 (s, 3H).

2) Preparation of 2,N-Dimethyl-4,5-diaminopyrimidine

A mixture of 6-Amino-4-chloro-2,N-dimethyl-5-nitropyrimidine (1.0 g, 0.5 mmol), NaOH (40 mg, 1 mmol) and 1.0 g 10% Pd/C in 50 mL of 2% aqueous THF is hydrogenated at 50 PSI of hydrogen at room temperature overnight. The reaction mixture is filtered through celite, concentrated in vacuo, and the obtained solid is purified on a silica gel column eluting with 10/1/0.1 CH₂Cl₂/MeOH/NH₄OH to give the title compound. ¹H NMR (CDCl₃) δ 7.64(s, 1H), 4.95 (bs, 1H), 3.02 (d, 3H), 2.45 (s, 3H), 1.90 (bs, 2H).

3) Preparation of 8-chloromethyl-2,9-dimethyl-9H-purine

A solution of 2,N-Dimethyl-4,5-diaminopyrimidine (180 mg, 1.3 mmol) and ethyl chloroacetimidate hydrochloride (310 g, 2.0

mmol) in dichloroethane (10 mL) is heated at reflux for 17 h and cooled. The mixture is washed with aqueous NaHCO₃, water, dried and concentrated to give 8-Chloromethyl-2,9-dimethyl-9H-purine.

5 4) Preparation 8-{[2-(6-fluoropyridin-2-yl)-1H-imidazol-1-yl]methyl}-2,9-dimethyl-9H-purine

10 A mixture of 8-Chloromethyl-2,9-dimethyl-9H-purine (40 mg, 0.2 mmol), 2-Fluoro-6-(1H-imidazol-2-yl)-pyridine (40 mg, 0.25 mmol) and K₂CO₃ (55 mg, 0.4 mmol) in 2 mL of DMF is stirred at room temperature for four hours. The mixture is diluted with water (10 mL), and extracted with ethyl acetate three times. The combined extract is washed with brine, dried and concentrated. The residue is purified by preparative thin layer chromatography to give the title compound. ¹H NMR(CDCl₃): δ 8.92 (s, 1H), 8.18 (dd, 1H), 7.88 (q, 1H), 7.25 (s, 1H), 7.20 (s, 1H), 7.11 (s, 1H), 6.83 (dd, 1H), 6.22 (s, 2H), 3.89 (s, 3H), 2.8 (s, 3H). LRMS calcd 323, found 324 (MH⁺).

15 **Example 49**

20 The following compounds are prepared essentially according to the procedures in the previous examples.

a) 3-Methyl-2-(2-oxazol-2-yl-imidazol-1-ylmethyl)-3H-imidazo[4,5-c]pyridine

25 b) 3-Ethyl-2-(2-[1,3,4]oxadiazol-2-yl-imidazol-1-ylmethyl)-3H-imidazo[4,5-c]pyridine

c) 3-Ethyl-2-[2-(3-methyl-pyridin-2-yl)-imidazol-1-ylmethyl]-3H-imidazo[4,5-c]pyridine

30 **Example 50**

The compounds listed in tables 1 - 8 are prepared essentially according to the procedures set forth above in Schemes I-X and the preceding examples.

5 In tables 1-5 the designations X_1 , X_2 , W_1 , X_5 , W_6 , etc., on the substituents W_1 , R_5 , etc., indicate the point of attachment of the substituent to the parent structural formula. For example, R_5 in compound number (hereinafter "Comp. #") 112 is an ethyl group; R_5 in compound 134 is a cyclopropylmethyl group; and
10 W_1 in compound 132 is a 3-chlorophenyl group.

LC-MS data is provided for a number of the compounds in tables 1-5. The following HPLC method was used to obtain this data: YMC-pack pro C_{18} column, 33 x 4.6 mm(L x ID), 5 μ m particle size. 3 min gradient from 5% to 95% B with 0.5 min hold
15 at 95% B. Solvent A: 95% H_2O -5%MeOH-0.05%TFA; Solvent B: 95%MeOH-5% H_2O -0.05%TFA). Flow rate = 2.0 ml/min. Injection volume = 1 μ l. MS (ES^+): m/e 360 $[MH]^+$. The LC data is given as HPLC retention times.

10038069 122101






























TABLE 1						
Cmp. #	R5	W1	MW	RT	Mass	GC/FID or LC/MS
100			302.3792	1.60	303.1	LC/MS
101			316.406	1.80	317.1	LC/MS
102			316.406	1.80	317.1	LC/MS
103			320.3693	5.06		GC/FID
104			334.3961	5.07		GC/FID
105			334.3961	5.17		GC/FID
106			306.3425	5.12		GC/FID
107			320.3693	5.11		GC/FID
108			334.3961	5.10		GC/FID
109			334.3961	5.20		GC/FID
110			334.3961	5.21		GC/FID
111			324.3326	4.95		GC/FID
112			338.3594	4.95		GC/FID
113			352.3862	4.95		GC/FID






























TABLE 1						
Cmp. #	R5	W1	MW	RT	Mass	GC/FID or LC/MS
114			352.3862	5.05		GC/FID
115			354.8144	5.43		GC/FID
116			368.8412	5.42		GC/FID
117			368.8412	5.51		GC/FID
118			308.4074	5.27		GC/FID
119			322.4342	5.26		GC/FID
120			322.4342	5.37		GC/FID
121			356.3495	4.99		GC/FID
122			370.3763	4.98		GC/FID
123			384.4031	4.98		GC/FID
124			384.4031	5.07		GC/FID
125			367.2485	5.68		GC/FID
126			381.2753	5.67		GC/FID
127			395.3021	5.65		GC/FID


































TABLE 1						
Cmp. #	R5	W1	MW	RT	Mass	GC/FID or LC/MS
128			395.3021	5.74		GC/FID
129			322.7975	1.67	323.1	LC/MS
130			336.8243	1.87	337.3	LC/MS
131			350.8511	2.00	351.1	LC/MS
132			350.8511	2.07	351.3	LC/MS
133			330.4328	2.00	331.3	LC/MS
134			328.417	1.87	329.2	LC/MS
135			348.4229	5.15		GC/FID
136			346.4071	5.28		GC/FID
137			348.4229	5.18		GC/FID
138			362.4497	5.32		GC/FID
139			346.4071	5.31		GC/FID
140			346.4071	5.31		GC/FID
141			366.413	5.04		GC/FID
142			380.4398	5.18		GC/FID
143			364.3972	5.16		GC/FID


























TABLE 1						
Cmp. #	R5	W1	MW	RT	Mass	GC/FID or LC/MS
144			382.868	5.47		GC/FID
145			380.8522	5.59		GC/FID
146			336.461	5.33		GC/FID
147			334.4452	5.46		GC/FID
148			398.4299	5.04		GC/FID
149			396.4141	5.17		GC/FID
150			409.3289	5.68		GC/FID
151			423.3557	5.80		GC/FID
152			407.3131	5.81		GC/FID
153			364.8779	2.20	365.2	LC/MS
154			378.9047	2.40	379.3	LC/MS
155			362.8621	2.07	363.3	LC/MS























TABLE 2					
					
Cmp. #	R2	R5	W6	RT	GC/FID or LC/MS
156				9.18	GC/FID
157				8.67	GC/FID
158				10.56	GC/FID
159				9.39	GC/FID
160				8.87	GC/FID
161				10.91	GC/FID
162				9.12	GC/FID























TABLE 2					
					
Cmp. #	R2	R5	W6	RT	GC/FID or LC/MS
163				8.61	LC/MS
164				10.46	GC/FID
165				8.98	GC/FID
166				8.49	GC/FID
167				10.28	GC/FID
168				9.77	GC/FID
169				9.17	GC/FID























TABLE 2					
					
Cmp. #	R2	R5	W6	RT	GC/FID or LC/MS
170				11.41	LC/MS
171				8.75	GC/FID
172				8.31	GC/FID
173				9.97	GC/FID
174				9.04	GC/FID
175				8.55	GC/FID
176				10.36	GC/FID
























TABLE 2					
					
Cmp. #	R2	R5	W6	RT	GC/FID or LC/MS
177				9.81	GC/FID
178				9.21	GC/FID
179				11.49	GC/FID
180				9.64	GC/FID
181				9.05	GC/FID
182				11.22	LC/MS
183				10.86	GC/FID

TABLE 3



Cpd. #	R2	R5	W1	RT	GC/FID or LC/MS
189				6.17	GC/FID
190				6.02	GC/FID
191				6.32	GC/FID
192				6.15	GC/FID
193				6.37	GC/FID
194				6.20	GC/FID
195				6.40	GC/FID
196				1.32	LC/MS
197				6.52	GC/FID
198				6.36	GC/FID
































TABLE 3					
					
Cpd. #	R2	R5	W1	RT	GC/FID or LC/MS
199				6.77	GC/FID
200				6.60	GC/FID
201				6.37	GC/FID
202				6.22	GC/FID
203				1.40	LC/MS
204				6.50	GC/FID
205				6.66	GC/FID
206				6.51	GC/FID
207				7.15	GC/FID
208				6.91	GC/FID

TABLE 3	
---------	--

TABLE 3					
Cpd. #	R2	R5	W1	RT	GC/FID or LC/MS
209				7.13	GC/FID
210				6.91	GC/FID
211				5.64	GC/FID
212				5.47	GC/FID
213				5.92	GC/FID
214				5.75	GC/FID
215				1.13	LC/MS
216				5.71	GC/FID
217				5.91	GC/FID
218				5.74	GC/FID


































TABLE 3					
					
Cpd. #	R2	R5	W1	RT	GC/FID or LC/MS
219				6.05	GC/FID
220				5.89	GC/FID
221				5.99	GC/FID
222				5.82	GC/FID
223				6.11	GC/FID
224				5.96	GC/FID
225				6.09	GC/FID
226				5.94	GC/FID
227				6.11	GC/FID
228				5.96	GC/FID

TABLE 3




Cpd. #	R2	R5	W1	RT	GC/FID or LC/MS
229		$\text{H}_3\text{C}-\text{CH}_2-\text{X}_5$		6.33	GC/FID
230		$\text{H}_3\text{C}-\text{CH}_2-\text{X}_5$		6.18	GC/FID
231		$\text{H}_3\text{C}-\text{CH}_2-\text{X}_5$		6.33	GC/FID
232		$\text{H}_3\text{C}-\text{CH}_2-\text{X}_5$		6.19	GC/FID
233		$\text{H}_3\text{C}-\text{CH}_2-\text{X}_5$		6.21	GC/FID
234		$\text{H}_3\text{C}-\text{CH}_2-\text{X}_5$		6.07	GC/FID
235		$\text{H}_3\text{C}-\text{CH}_2-\text{X}_5$		5.81	GC/FID
236		$\text{H}_3\text{C}-\text{CH}_2-\text{X}_5$		5.64	GC/FID
237		$\text{H}_3\text{C}-\text{CH}_2-\text{X}_5$		6.29	GC/FID
238		$\text{H}_3\text{C}-\text{CH}_2-\text{X}_5$		6.13	GC/FID

TABLE 3




Cpd. #	R2	R5	W1	RT	GC/FID or LC/MS
239				5.97	GC/FID
240				5.82	GC/FID
241				6.51	GC/FID
242				6.36	GC/FID
243				6.42	GC/FID
244				6.26	GC/FID
245				6.52	GC/FID
246				6.38	GC/FID
247				6.34	GC/FID
248				6.19	GC/FID

TABLE 3



Cpd. #	R2	R5	W1	RT	GC/FID or LC/MS
249				6.45	GC/FID
250				6.32	GC/FID
251				6.71	GC/FID
252				2.18	LC/MS
253				6.28	GC/FID
254				6.12	GC/FID
255				7.38	GC/FID
256				7.11	GC/FID
257				1.48	LC/MS
258				7.95	GC/FID

TABLE 3



Cpd. #	R2	R5	W1	RT	GC/FID or LC/MS
259				6.34	GC/FID
260				6.18	GC/FID
261				6.47	GC/FID
262				6.32	GC/FID
263				6.10	GC/FID
264				5.95	GC/FID
265				6.15	GC/FID
266				6.00	GC/FID
267				7.43	GC/FID
268				7.15	GC/FID




























































































TABLE 3					
					
Cpd. #	R2	R5	W1	RT	GC/FID or LC/MS
269				7.74	GC/FID
270				7.41	GC/FID
271				6.65	GC/FID
272				6.50	GC/FID
273				6.23	GC/FID
274				6.08	GC/FID
275				6.30	GC/FID
276				6.15	GC/FID

TABLE 4						
						
Cmp. #	R2	R5	W1	RT	Mass	GC/FID or LC/MS
277		$\text{H}_3\text{C}-\text{CH}_2-\text{X}_5$		7.12		GC/FID
278		$\text{H}_3\text{C}-\text{CH}_2-\text{X}_5$		6.77		GC/FID
279		$\text{H}_3\text{C}-\text{CH}_2-\text{X}_5$		7.17		GC/FID
280		$\text{H}_3\text{C}-\text{CH}_2-\text{X}_5$		6.82		GC/FID
281		$\text{H}_3\text{C}-\text{CH}_2-\text{X}_5$		7.10		GC/FID
282		$\text{H}_3\text{C}-\text{CH}_2-\text{X}_5$		6.77		GC/FID
283		$\text{H}_3\text{C}-\text{CH}_2-\text{X}_5$		7.30		GC/FID
284		$\text{H}_3\text{C}-\text{CH}_2-\text{X}_5$		6.92		GC/FID
285		$\text{H}_3\text{C}-\text{CH}_2-\text{X}_5$		7.53		GC/FID

<div> <div>TABLE 4</div> <div> </div> </div>						
Cmp. #	R2	R5	W1	RT	Mass	GC/FID or LC/MS
286				7.07		GC/FID
287				7.79		GC/FID
288				7.28		GC/FID
289				7.29		GC/FID
290				6.92		GC/FID
291				7.86		GC/FID
292				7.34		GC/FID
293				7.77		GC/FID
294				7.29		GC/FID

<div> <div>TABLE 4</div> <div> </div> </div>						
Cmp. #	R2	R5	W1	RT	Mass	GC/FID or LC/MS
295		$\text{H}_3\text{C}-\text{CH}_2-\text{X}_5$		1.44	472.4	LC/MS
296		$\text{H}_3\text{C}-\text{CH}_2-\text{X}_5$		1.56	508.4	LC/MS
297		$\text{H}_3\text{C}-\text{CH}_2-\text{X}_5$		2.02	468.5	LC/MS
298		$\text{H}_3\text{C}-\text{CH}_2-\text{X}_5$		7.84		GC/FID
299		$\text{H}_3\text{C}-\text{CH}_2-\text{X}_5$		6.41		GC/FID
300		$\text{H}_3\text{C}-\text{CH}_2-\text{X}_5$		1.32	410.3	LC/MS
301		$\text{H}_3\text{C}-\text{CH}_2-\text{X}_5$		6.59		GC/FID
302		$\text{H}_3\text{C}-\text{CH}_2-\text{X}_5$		6.35		GC/FID
303		$\text{H}_3\text{C}-\text{CH}_2-\text{X}_5$		6.54		GC/FID

TABLE 4						
						
Cmp. #	R2	R5	W1	RT	Mass	GC/FID or LC/MS
304				6.29		GC/FID
305				6.61		GC/FID
306				6.37		GC/FID
307				6.76		GC/FID
308				6.50		GC/FID
309				6.58		GC/FID
310				6.34		GC/FID
311				6.72		GC/FID
312				6.47		GC/FID

<div> <div>TABLE 4</div> <div>  </div> </div> <div></div> <div></div> <div></div> <div></div> <div></div> <div></div>						
Cmp. #	R2	R5	W1	RT	Mass	GC/FID or LC/MS
322		$\text{H}_3\text{C}-\text{CH}_2-\text{X}_5$		6.67		GC/FID
323		$\text{H}_3\text{C}-\text{CH}_2-\text{X}_5$		6.49		GC/FID
324		$\text{H}_3\text{C}-\text{CH}_2-\text{X}_5$		6.26		GC/FID
325		$\text{H}_3\text{C}-\text{CH}_2-\text{X}_5$		7.13		GC/FID
326		$\text{H}_3\text{C}-\text{CH}_2-\text{X}_5$		6.77		GC/FID
327		$\text{H}_3\text{C}-\text{CH}_2-\text{X}_5$		6.65		GC/FID
328		$\text{H}_3\text{C}-\text{CH}_2-\text{X}_5$		6.42		GC/FID
329		$\text{H}_3\text{C}-\text{CH}_2-\text{X}_5$		7.39		GC/FID
330		$\text{H}_3\text{C}-\text{CH}_2-\text{X}_5$		7.00		GC/FID






















<div> TABLE 4  </div>						
Cmp. #	R2	R5	W1	RT	Mass	GC/FID or LC/MS
331		$\text{H}_3\text{C}-\text{CH}_2-\text{X}_5$		7.26		GC/FID
332		$\text{H}_3\text{C}-\text{CH}_2-\text{X}_5$		6.90		GC/FID
333		$\text{H}_3\text{C}-\text{CH}_2-\text{X}_5$		1.99	456.5	LC/MS
334		$\text{H}_3\text{C}-\text{CH}_2-\text{X}_5$		2.06	492.5	LC/MS
335		$\text{H}_3\text{C}-\text{CH}_2-\text{X}_5$		7.00		GC/FID
336		$\text{H}_3\text{C}-\text{CH}_2-\text{X}_5$		6.71		GC/FID
337		$\text{H}_3\text{C}-\text{CH}_2-\text{X}_5$		7.31		GC/FID
338		$\text{H}_3\text{C}-\text{CH}_2-\text{X}_5$		2.45	545.6	LC/MS
339		$\text{H}_3\text{C}-\text{CH}_2-\text{X}_5$		7.86		GC/FID

TABLE 4



Cmp. #	R2	R5	W1	RT	Mass	GC/FID or LC/MS
349		$\text{H}_3\text{C}-\text{CH}_2-\text{X}_5$		7.44		GC/FID
350		$\text{H}_3\text{C}-\text{CH}_2-\text{X}_5$		7.04		GC/FID
351		$\text{H}_3\text{C}-\text{CH}_2-\text{X}_5$		6.83		GC/FID
352		$\text{H}_3\text{C}-\text{CH}_2-\text{X}_5$		6.56		GC/FID
353		$\text{H}_3\text{C}-\text{CH}_2-\text{X}_5$		6.84		GC/FID
354		$\text{H}_3\text{C}-\text{CH}_2-\text{X}_5$		6.58		GC/FID
355		$\text{H}_3\text{C}-\text{CH}_2-\text{X}_5$		9.63		GC/FID
356		$\text{H}_3\text{C}-\text{CH}_2-\text{X}_5$		8.64		GC/FID
357		$\text{H}_3\text{C}-\text{CH}_2-\text{X}_5$		1.05	497.6	LC/MS

TABLE 4



Cmp. #	R2	R5	W1	RT	Mass	GC/FID or LC/MS
358		$\text{H}_3\text{C}-\text{CH}_2-\text{X}_5$		9.16		GC/FID
359		$\text{H}_3\text{C}-\text{CH}_2-\text{X}_5$		7.61		GC/FID
360		$\text{H}_3\text{C}-\text{CH}_2-\text{X}_5$		7.17		GC/FID
361		$\text{H}_3\text{C}-\text{CH}_2-\text{X}_5$		7.02		GC/FID
362		$\text{H}_3\text{C}-\text{CH}_2-\text{X}_5$		6.72		GC/FID
363		$\text{H}_3\text{C}-\text{CH}_2-\text{X}_5$		1.05	459.5	LC/MS
364		$\text{H}_3\text{C}-\text{CH}_2-\text{X}_5$		6.79		GC/FID

<div> <div>TABLE 5</div> <div> </div> </div>							
Cpd. #	R2	R5	W1	MW	RT	Mass	GC/FID or LC/MS
365		$H_3C-CH_2-CH_2-X_5$		413.5223	NA		
366		$H_3C-CH_2-CH_2-X_5$		431.5124	1.73	432.1	LC/MS
367		$H_3C-CH_2-CH_2-X_5$		449.5025	1.67	450.4	LC/MS
368		$H_3C-CH_2-CH_2-X_5$		447.9674	1.87	448.3	LC/MS
369		$H_3C-CH_2-CH_2-X_5$		427.5491	1.87	428.4	LC/MS
370		$H_3C-CH_2-CH_2-X_5$		445.5392	1.94	446.4	LC/MS
371		$H_3C-CH_2-CH_2-X_5$		463.5293	1.87	464.4	LC/MS
372		$H_3C-CH_2-CH_2-X_5$		461.9942	2.07	462.4	LC/MS


























TABLE 5							
Cpd. #	R2	R5	W1	MW	RT	Mass	GC/FID or LC/MS
373				429.5213	2.87	430	LC/MS
374				447.5114	1.53	448.3	LC/MS
375				465.5015	1.47	466.5	LC/MS
376				463.9664	1.67	464.4	LC/MS
377				441.5759	2.07	442.6	LC/MS
378				459.566	2.13	460.3	LC/MS
379				477.5561	NA		
380				441.5759	2.00	442.6	LC/MS


























TABLE 5							
Cpd. #	R2	R5	W1	MW	RT	Mass	GC/FID or LC/MS
381				459.566	2.07	460.3	LC/MS
382				445.5883	1.73	446.4	LC/MS
383				463.5784	1.80	464.4	LC/MS
384				481.5685	1.73	482.2	LC/MS
385				480.0334	NA		
386				455.6027	2.13	456.4	LC/MS
387				473.5928	NA		
388				491.5829	2.13	492.4	LC/MS


























TABLE 5							
Cpd. #	R2	R5	W1	MW	RT	Mass	GC/FID or LC/MS
389				490.0478	2.27	490.3	LC/MS
390				455.6027	2.13	456.4	LC/MS
391				473.5928	NA		
392				491.5829	2.13	492.4	LC/MS
393				490.0478	2.27	490.5	LC/MS
394				483.6563	2.40	484.5	LC/MS
395				501.6464	2.47	502.3	LC/MS
396				519.6365	NA		












































































TABLE 5							
Cpd. #	R2	R5	W1	MW	RT	Mass	GC/FID or LC/MS
397				518.1014	NA		
398				485.5849	1.73	486.3	LC/MS
399				503.575	1.73	504.3	LC/MS
400				521.5651	NA		
401				520.03	1.87	520.2	LC/MS
402				481.6405	2.34	482.5	LC/MS
403				499.6306	2.40	500.4	LC/MS
404				517.6207	2.34	518	LC/MS

TABLE 5								
Cpd. #	R2	R5	W1	MW	RT	Mass	GC/FID or LC/MS	
405				516.0856	2.47	516.4	LC/MS	
406				387.4845	1.47	388.5	LC/MS	
407				405.4746	1.53	406.3	LC/MS	
408				423.4647	1.47	424.2	LC/MS	
409				413.5223	1.80	414.4	LC/MS	
410				431.5124	1.87	432.1	LC/MS	
411				449.5025	1.73	450.3	LC/MS	
412				447.9674	1.93	448.5	LC/MS	

<div> <div>TABLE 5</div> <div>  </div> </div>							
Cpd. #	R2	R5	W1	MW	RT	Mass	GC/FID or LC/MS
413				415.5381	1.80	416.4	LC/MS
414				433.5282	1.87	434.5	LC/MS
415				451.5183	1.80	452.4	LC/MS
416				449.9832	2.00	450.3	LC/MS
417				415.5381	1.80	416.4	LC/MS
418				433.5282	1.87	434.5	LC/MS
419				451.5183	1.80	452.4	LC/MS
420				449.9832	2.00	450.3	LC/MS

<div> <div>TABLE 5</div> <div> </div> </div>							
Cpd. #	R2	R5	W1	MW	RT	Mass	GC/FID or LC/MS
421				429.5649	2.00	430.2	LC/MS
422				447.555	2.07	448.6	LC/MS
423				465.5451	2.00	466.3	LC/MS
424				464.01	2.20	464.4	LC/MS
425				429.5649	1.93	430.2	LC/MS
426				447.555	2.00	448.6	LC/MS
427				465.5451	1.93	466.3	LC/MS
428				464.01	2.13	464.4	LC/MS



























TABLE 5							
Cpd. #	R2	R5	W1	MW	RT	Mass	GC/FID or LC/MS
429				439.5601	2.00	440.7	LC/MS
430				457.5502	2.07	458.4	LC/MS
431				475.5403	NA		
432				474.0052	NA		
433				443.5917	2.13	444.6	LC/MS
434				461.5818	2.20	462.4	LC/MS
435				479.5719	NA		
436				478.0368	NA		

TABLE 5



Cpd. #	R2	R5	W1	MW	RT	Mass	GC/FID or LC/MS
437				443.5917	2.13	444.6	LC/MS
438				461.5818	2.20	462.4	LC/MS
439				479.5719	NA		
440				478.0368	NA		
441				455.6027	2.20	456.4	LC/MS
442				473.5928	NA		
443				491.5829	NA		
444				490.0478	2.34	490.3	LC/MS














TABLE 5							
Cpd. #	R2	R5	W1	MW	RT	Mass	GC/FID or LC/MS
445				457.6185	2.34	458.4	LC/MS
446				475.6086	NA		
447				493.5987	2.34	494.5	LC/MS
448				492.0636	2.47	492.4	LC/MS










TABLE 6				
Cpd#	STRUCTURE	IUPAC NAME	NMR	MS (m/z)
449		2-((2-(3-fluorophenyl)-1H-imidazol-1-yl)methyl)-1-propyl-1H-benzimidazole	¹ H NMR (CDCl ₃): 7.80 (m, 1H), 7.41-7.53 (m, 3H), 7.29-7.31 (m, 3H), 7.14-7.22 (m, 2H), 7.06 (d, 1H), 5.51 (s, 2H), 3.69 (t, 2H), 1.43 (m, 2H), 0.74 (t, 3H)	
450		2-((2-(2-fluorophenyl)-1H-imidazol-1-yl)methyl)-1-propyl-1H-benzimidazole	7.78 (m, 1H), 7.66 (m, 1H), 7.51 (m, 1H), 7.23-7.37 (m, 5H), 7.17 (d, 1H), 7.03 (d, 1H), 5.35 (s, 2H), 3.68 (t, 2H), 1.40 (m, 2H), 0.72 (t, 3H)	
451		2-((2-(4-fluorophenyl)-1H-imidazol-1-yl)methyl)-1-propyl-1H-benzimidazole	¹ H NMR (CDCl ₃): 7.80 (m, 1H), 7.65-7.69 (m, 2H), 7.18-7.32 (m, 5H), 7.13 (d, 1H), 7.04 (d, 1H), 5.47 (s, 2H), 3.68 (t, 2H), 1.42 (m, 2H), 0.74 (t, 3H)	
452		2-((2-(3-chloro-4-fluorophenyl)-1H-imidazol-1-yl)methyl)-1-propyl-1H-benzimidazole	¹ H NMR (CDCl ₃): 7.76-7.81 (m, 2H), 7.56 (m, 1H), 7.25-7.32 (m, 4H), 7.14 (d, 1H), 7.05 (d, 1H), 5.46 (s, 2H), 3.74 (t, 2H), 1.47 (m, 2H), 0.78 (t, 3H)	
453		2-((2-(2,5-difluorophenyl)-1H-imidazol-1-yl)methyl)-1-propyl-1H-benzimidazole	7.78 (m, 1H), 7.38 (m, 1H), 7.17-7.31 (m, 6H), 7.03 (d, 1H), 5.36 (s, 2H), 3.73 (t, 2H), 1.44 (m, 2H), 0.76 (t, 3H)	
454		6-chloro-2-((2-(2-fluorophenyl)-1H-imidazol-1-yl)methyl)-1-propyl-1H-benzimidazole	¹ H NMR (CDCl ₃): 7.62-7.69 (m, 2H), 7.51 (m, 1H), 7.22-7.34 (m, 4H), 7.18 (d, 1H), 7.02 (d, 1H), 5.34 (s, 2H), 3.64 (t, 2H), 1.40 (m, 2H), 0.73 (t, 3H)	
455		6-chloro-2-((2-(3-fluorophenyl)-1H-imidazol-1-yl)methyl)-1-propyl-1H-benzimidazole	¹ H NMR (CDCl ₃): 7.69 (d, 1H), 7.38-7.52 (m, 3H), 7.15-7.29 (m, 4H), 7.04 (d, 1H), 5.48 (s, 2H), 3.65 (t, 2H), 1.42 (m, 2H), 0.75 (t, 3H)	
456		6-chloro-2-((2-(4-fluorophenyl)-1H-imidazol-1-yl)methyl)-1-propyl-1H-benzimidazole	¹ H NMR (CDCl ₃): 7.62-7.70 (m, 3H), 7.13-7.28 (m, 5H), 7.02 (d, 1H), 3.64 (t, 2H), 1.41 (m, 2H), 0.75 (t, 3H)	
457		6-chloro-2-((2-(3-chloro-4-fluorophenyl)-1H-imidazol-1-yl)methyl)-1-propyl-1H-benzimidazole	¹ H NMR (CDCl ₃): 7.75 (dd, 1H), 7.69 (dd, 1H), 7.55 (m, 1H), 7.24-7.30 (m, 3H), 7.15 (d, 1H), 7.03 (d, 1H), 5.43 (s, 2H), 3.71 (t, 2H), 1.47 (m, 2H), 0.79 (t, 3H)	

TABLE 6				
Cpd#	STRUCTURE	IUPAC NAME	NMR	MS (m/z)
458		6-chloro-2-([2-(2,5-difluorophenyl)-1H-imidazol-1-yl]methyl)-1-propyl-1H-benzimidazole	¹ H NMR (CDCl ₃): 7.67 (d, 1H), 7.37 (m, 1H), 7.17-7.29 (m, 5H), 7.02 (d, 1H), 5.34 (s, 2H), 3.69 (t, 2H), 1.43 (m, 2H), 0.77 (t, 3H)	
459		2-([2-(3-fluorophenyl)-1H-imidazol-1-yl]methyl)-3-propyl-3H-imidazo[4,5-b]pyridine	¹ H NMR (CDCl ₃): 8.40 (dd, 1H), 8.06 (dd, 1H), 7.39-7.50 (m, 3H), 7.15-7.29 (m, 3H), 7.08 (d, 1H), 5.52 (s, 2H), 3.88 (t, 2H), 1.51 (m, 2H), 0.77 (t, 3H)	
460		2-([2-(2,5-difluorophenyl)-1H-imidazol-1-yl]methyl)-3-propyl-3H-imidazo[4,5-b]pyridine	¹ H NMR (CDCl ₃): 8.40 (dd, 1H), 8.05 (dd, 1H), 7.38 (m, 1H), 7.17-7.28 (m, 4H), 7.08 (d, 1H), 5.39 (s, 2H), 3.89 (t, 2H), 1.50 (m, 2H), 0.79 (t, 3H)	
461		6-chloro-1-(cyclopropylmethyl)-2-([2-(2,5-difluorophenyl)-1H-imidazol-1-yl]methyl)-1H-benzimidazole	¹ H NMR (CDCl ₃): 7.68 (d, 1H), 7.38 (m, 1H), 7.32 (d, 1H), 7.19-7.28 (m, 4H), 7.04 (d, 1H), 5.36 (d, 2H), 3.62 (d, 2H), 0.65 (m, 1H), 0.46-0.51 (m, 2H), 0.07-0.11 (m, 2H)	
462		2-([2-(2,5-difluorophenyl)-1H-imidazol-1-yl]methyl)-3-isobutyl-3H-imidazo[4,5-b]pyridine	¹ H NMR (CDCl ₃): 8.38 (dd, 1H), 8.04 (dd, 1H), 7.38 (m, 1H), 7.16-7.27 (m, 4H), 7.09 (d, 1H), 5.38 (s, 2H), 3.73 (d, 2H), 1.98 (m, 1H), 0.72 (d, 6H)	
463		6-chloro-2-([2-(2,5-difluorophenyl)-1H-imidazol-1-yl]methyl)-1-ethyl-1H-benzimidazole	¹ H NMR (CDCl ₃): 7.67 (d, 1H), 7.37 (m, 1H), 7.17-7.30 (m, 5H), 7.01 (d, 1H), 5.35 (s, 2H), 3.79 (q, 2H), 0.99 (t, 3H)	
464		6-chloro-1-ethyl-2-([2-(3-fluorophenyl)-1H-imidazol-1-yl]methyl)-1H-benzimidazole	¹ H NMR (CDCl ₃): 7.69 (d, 1H), 7.39-7.51 (m, 3H), 7.16-7.30 (m, 4H), 7.03 (d, 1H), 5.49 (s, 2H), 3.75 (q, 2H), 0.98 (t, 3H)	
465		2-([2-(3-fluorophenyl)-1H-imidazol-1-yl]methyl)-3-methyl-3H-imidazo[4,5-b]pyridine	¹ H NMR (CDCl ₃): 8.40 (dd, 1H), 8.05 (dd, 1H), 7.40-7.51 (m, 3H), 7.13-7.29 (m, 3H), 7.02 (d, 1H), 5.53 (s, 2H), 3.51 (s, 3H)	
466		2-([2-(2,5-difluorophenyl)-1H-imidazol-1-yl]methyl)-3-methyl-3H-imidazo[4,5-b]pyridine	¹ H NMR (CDCl ₃): 8.40 (dd, 1H), 8.03 (dd, 1H), 7.36 (m, 1H), 7.15-7.27 (m, 4H), 7.03 (d, 1H), 5.41 (s, 2H), 3.51 (s, 3H)	

TABLE 6				
Cpd#	STRUCTURE	IUPAC NAME	NMR	MS (m/z)
467		2-([2-(3-chloro-4-fluorophenyl)-1H-imidazol-1-yl]methyl)-3-methyl-3H-imidazo[4,5-b]pyridine		
468		3-(cyclopropylmethyl)-2-([2-(2,5-difluorophenyl)-1H-imidazol-1-yl]methyl)-3H-imidazo[4,5-b]pyridine	1H NMR (CDCl3): 8.37 (dd, 1H), 8.04 (dd, 1H), 7.38 (m, 1H), 7.09-7.28 (m, 4H), 7.09 (d, 1H), 5.42 (s, 2H), 3.83 (d, 2H), 0.68 (m, 1H), 0.38-0.43 (m, 2H), 0.24-0.29 (m, 2H)	
469		6-chloro-2-([2-(3-chlorophenyl)-1H-imidazol-1-yl]methyl)-1-ethyl-1H-benzimidazole	1H NMR (CDCl3): 7.67-7.71 (m, 2H), 7.45-7.56 (m, 3H), 7.24-7.30 (m, 2H), 7.16 (d, 2H), 7.04 (d, 1H), 5.47 (s, 2H), 3.76 (q, 2H), 0.99 (t, 3H)	
470		2-([2-(3-chlorophenyl)-1H-imidazol-1-yl]methyl)-3-(cyclopropylmethyl)-3H-imidazo[4,5-b]pyridine	1H NMR (CDCl3): 8.39 (dd, 1H), 8.06 (dd, 1H), 7.68 (m, 1H), 7.40-7.68 (m, 3H), 7.27 (m, 1H), 7.18 (d, 1H), 7.10 (d, 1H), 5.53 (s, 2H), 3.83 (d, 2H), 0.73 (m, 1H), 0.40-0.46 (m, 2H), 0.24-0.28 (m, 2H)	
471		6-chloro-2-([2-(3-chlorophenyl)-1H-imidazol-1-yl]methyl)-1-(cyclopropylmethyl)-1H-benzimidazole	1H NMR (CDCl3): 7.67-7.71 (m, 2H), 7.42-7.56 (m, 3H), 7.24-7.33 (m, 2H), 7.15 (d, 1H), 7.06 (d, 1H), 5.47 (s, 2H), 3.58 (d, 2H), 0.69 (m, 1H), 0.47-0.53 (m, 2H), 0.03-0.09 (m, 2H)	
472		2-([2-(3-chlorophenyl)-1H-imidazol-1-yl]methyl)-1-propyl-1H-benzimidazole	1H NMR (CDCl3): 7.79 (m, 1H), 7.69 (m, 1H), 7.44-7.58 (m, 3H), 7.27-7.30 (m, 3H), 7.14 (d, 1H), 7.07 (d, 1H), 5.49 (s, 2H), 3.69 (t, 2H), 1.42 (m, 2H), 0.75 (t, 3H)	
473		2-([2-(3-chlorophenyl)-1H-imidazol-1-yl]methyl)-3-ethyl-3H-imidazo[4,5-b]pyridine	1H NMR (CDCl3): 8.40 (dd, 1H), 8.06 (dd, 1H), 7.67 (m, 1H), 7.43-7.57 (m, 3H), 7.26 (m, 1H), 7.19 (d, 1H), 7.08 (d, 1H), 5.51 (s, 2H), 4.00 (q, 2H), 1.09 (t, 3H)	
474		2-([2-(3-chlorophenyl)-1H-imidazol-1-yl]methyl)-3-propyl-3H-imidazo[4,5-b]pyridine	1H NMR (CDCl3): 8.40 (dd, 1H), 8.06 (dd, 1H), 7.68 (m, 1H), 7.43-7.56 (m, 3H), 7.26 (m, 1H), 7.18 (d, 1H), 7.09 (d, 1H), 5.50 (s, 2H), 3.89 (t, 2H), 1.52 (m, 2H), 0.79 (t, 3H)	
475		1-ethyl-2-([2-(3-fluorophenyl)-1H-imidazol-1-yl]methyl)-1H-benzimidazole-5-carbonitrile		346.3 [M+1]; 344.2 [M-1]

TABLE 6				
Cpd#	STRUCTURE	IUPAC NAME	NMR	MS (m/z)
476		1-ethyl-2-([2-(3-fluorophenyl)-1H-imidazol-1-yl]methyl)-5-(trifluoromethyl)-1H-benzimidazole		389.2 [M+1]
477		2-([2-(2,5-difluorophenyl)-1H-imidazol-1-yl]methyl)-1-ethyl-5-(trifluoromethyl)-1H-benzimidazole		407.3 [M+1]
478		2-([2-(3-fluorophenyl)-1H-imidazol-1-yl]methyl)-3-(2,2,2-trifluoroethyl)-3H-imidazo[4,5-b]pyridine		376.2 [M+1]
479		2-([2-(2,5-difluorophenyl)-1H-imidazol-1-yl]methyl)-3-(2,2,2-trifluoroethyl)-3H-imidazo[4,5-b]pyridine		394.2 [M+1]
480		2-([2-(3-chlorophenyl)-1H-imidazol-1-yl]methyl)-3-(2,2,2-trifluoroethyl)-3H-imidazo[4,5-b]pyridine		392.2 [M+1]
481		1-ethyl-5-(trifluoromethyl)-2-([2-(3-(trifluoromethyl)phenyl)-1H-imidazol-1-yl]methyl)-1H-benzimidazole	1H NMR (CDCl3): 8.09 (d, 1H), 7.37-7.95 (m, 6H), 7.19 (d, 1H), 7.05 (d, 1H), 5.49 (s, 2H), 3.85 (q, 2H), 1.02 (t, 3H)	
482		5-chloro-3-propyl-2-([2-(3-(trifluoromethyl)phenyl)-1H-imidazol-1-yl]methyl)-3H-imidazo[4,5-b]pyridine	1H NMR (CDCl3): 7.60-7.97 (m, 5H), 7.22-7.27 (m, 2H), 7.09 (d, 1H), 5.45 (s, 2H), 3.85 (t, 2H), 1.51 (m, 2H), 0.76 (t, 3H)	
483		3-(2,2,2-trifluoroethyl)-2-([2-(3-(trifluoromethyl)phenyl)-1H-imidazol-1-yl]methyl)-3H-imidazo[4,5-b]pyridine	1H NMR (CDCl3): 8.44 (dd, 1H), 8.07 (dd, 1H), 7.54-7.89 (m, 4H), 7.24-7.34 (m, 2H), 7.12 (d, 1H), 5.51 (s, 2H), 4.51 (q, 2H)	
484		3-(cyclopropylmethyl)-2-([2-(3-(trifluoromethyl)phenyl)-1H-imidazol-1-yl]methyl)-3H-imidazo[4,5-b]pyridine	1H NMR (CDCl3): 8.39 (dd, 1H), 7.60-8.03 (m, 5H), 7.21-7.26 (m, 2H), 7.11 (d, 1H), 5.51 (s, 2H), 3.84 (d, 2H), 0.70 (m, 1H), 0.44 (m, 2H), 0.24 (m, 2H)	

TABLE 6				
Cpd#	STRUCTURE	IUPAC NAME	NMR	MS (m/z)
485		3-isobutyl-2-((2-[3-(trifluoromethyl)phenyl]-1H-imidazol-1-yl)methyl)-3H-imidazo[4,5-b]pyridine	¹ H NMR (CDCl ₃): 8.39 (dd, 1H), 7.60-8.07 (m, 5H), 7.21-7.26 (m, 2H), 7.12 (d, 1H), 5.49 (s, 2H), 3.71 (d, 2H), 1.98 (m, 1H), 0.69 (d, 6H)	
486		3-propyl-2-((2-[3-(trifluoromethyl)phenyl]-1H-imidazol-1-yl)methyl)-3H-imidazo[4,5-b]pyridine	¹ H NMR (CDCl ₃): 8.40 (dd, 1H), 7.60-8.07 (m, 5H), 7.21-7.26 (m, 2H), 7.11 (d, 1H), 5.49 (s, 2H), 3.89 (t, 2H), 1.51 (m, 2H), 0.76 (t, 3H)	
487		1-propyl-2-((2-[3-(trifluoromethyl)phenyl]-1H-imidazol-1-yl)methyl)-1H-benzimidazole	¹ H NMR (CDCl ₃): 7.61-8.00 (m, 5H), 7.09-7.31 (m, 5H), 5.47 (s, 2H), 3.69 (t, 2H), 1.43 (m, 2H), 0.73 (t, 3H)	
488		6-chloro-1-(cyclopropylmethyl)-2-((2-[3-(trifluoromethyl)phenyl]-1H-imidazol-1-yl)methyl)-1H-benzimidazole	¹ H NMR (CDCl ₃): 7.60-7.96 (m, 5H), 7.05-7.31 (m, 4H), 5.45 (s, 2H), 3.58 (d, 2H), 0.65 (m, 1H), 0.47 (m, 2H), 0.05 (m, 2H)	
489		3-ethyl-2-((2-[3-(trifluoromethyl)phenyl]-1H-imidazol-1-yl)methyl)-3H-imidazo[4,5-b]pyridine	¹ H NMR (CDCl ₃): 8.40 (dd, 1H), 7.60-8.06 (m 5H), 7.21-7.28 (m, 2H), 7.09 (d, 1H), 5.49 (s, 2H), 4.01 (q, 2H), 1.09 (t, 3H)	
490		ethyl 2-([2-(3-chlorophenyl)-1H-imidazol-1-yl]methyl)-1-ethyl-1H-benzimidazole-5-carboxylate		
491		(2-([2-(3-chlorophenyl)-1H-imidazol-1-yl]methyl)-1-ethyl-1H-benzimidazol-5-yl)methanol		
492		2-([2-(3-chlorophenyl)-1H-imidazol-1-yl]methyl)-1-ethyl-5-[(4-methylpiperidin-1-yl)methyl]-1H-benzimidazole		
493		2-([2-(3-chlorophenyl)-1H-imidazol-1-yl]methyl)-1-ethyl-5-(morpholin-4-ylmethyl)-1H-benzimidazole		

TABLE 6				
Cpd#	STRUCTURE	IUPAC NAME	NMR	MS (m/z)
494		2-([2-(3-chlorophenyl)-1H-imidazol-1-yl]methyl)-1-ethyl-1H-benzimidazole-5-carboxylic acid		
495		1-ethyl-2-([2-(3-fluorophenyl)-1H-imidazol-1-yl]methyl)-5-(morpholin-4-ylmethyl)-1H-benzimidazole		420.4 [M+1]; 418.2 [M-1]
496		2-([2-(3-chlorophenyl)-1H-imidazol-1-yl]methyl)-1-ethyl-1H-benzimidazole-5-carboxamide		
497		2-([2-(3-chlorophenyl)-1H-imidazol-1-yl]methyl)-1-ethyl-N-methyl-1H-benzimidazole-5-carboxamide		
498		1-ethyl-2-([2-(3-fluorophenyl)-1H-imidazol-1-yl]methyl)-1H-benzimidazole-5-carboxylic acid		365.1 [M+1]
499		1-ethyl-2-([2-(3-fluorophenyl)-1H-imidazol-1-yl]methyl)-1H-benzimidazole-5-carboxamide		
500		2-([2-(2,5-difluorophenyl)-1H-imidazol-1-yl]methyl)-1-ethyl-1H-benzimidazole-5-carboxylic acid		383.2 [M+1]
501		1-ethyl-2-([2-(3-fluorophenyl)-1H-imidazol-1-yl]methyl)-5-((4-methylpiperidin-1-yl)methyl)-1H-benzimidazole		432.4 [M+1]
502		2-([2-(2,5-difluorophenyl)-1H-imidazol-1-yl]methyl)-N-[2-(dipropylamino)ethyl]-1-ethyl-1H-benzimidazole-5-carboxamide		509.5 [M+1]

TABLE 6				
Cpd#	STRUCTURE	IUPAC NAME	NMR	MS (m/z)
503		5-fluoro-2-([2-(4-fluorophenyl)-1H-imidazol-1-yl]methyl)-1-propyl-1H-benzimidazole	¹ H NMR(CDCl ₃): 7.65-7.71 (m, 2H), 7.47 (d, 1H), 7.03-7.23 (m, 6H), 5.47 (s, 2H), 3.65 (t, 2H), 1.41 (h, 2H), 0.71 (t, 3H)	
504		2-([2-(3-chlorophenyl)-1H-imidazol-1-yl]methyl)-5-fluoro-1-propyl-1H-benzimidazole	¹ H NMR(CDCl ₃): 7.65 (s, 1H), 7.41-7.53 (m, 3H), 7.06-7.24 (m, 5H), 5.47 (s, 2H), 3.65 (t, 2H), 1.47 (h, 2H), 0.71 (t, 3H)	
505		5-fluoro-2-([2-(3-fluorophenyl)-1H-imidazol-1-yl]methyl)-1-propyl-1H-benzimidazole	¹ H NMR(CDCl ₃): 7.35-7.47 (m, 4H), 7.0-7.1 (m, 5H), 5.47 (s, 2H), 3.65 (t, 2H), 1.35 (h, 2H), 0.71 (t, 3H)	
506		5-chloro-2-([2-(2,5-difluorophenyl)-1H-imidazol-1-yl]methyl)-1-ethyl-1H-benzimidazole	¹ H NMR (CDCl ₃): 7.75 (d, 1H), 7.37 (m, 1H), 7.20-7.27 (m, 5H), 7.02 (d, 1H), 5.36 (s, 2H), [0.99 T, 3H), 3.82 (q, 2H)	
507		2-([2-(2,5-difluorophenyl)-1H-imidazol-1-yl]methyl)-1-ethyl-5-fluoro-1H-benzimidazole	¹ H NMR (CDCl ₃): 7.43 (m, 1H), 7.37 (m, 1H), 7.19-7.25 (m, 4H), 7.03-7.10 (m, 2H), 5.35 (s, 2H), 3.82 (q, 2H), 1.00 (t, 3H)	
508		5-chloro-2-([2-(3-chlorophenyl)-1H-imidazol-1-yl]methyl)-1-ethyl-1H-benzimidazole	(L)-Tartrate salt ¹ H NMR (CD ₃ OD): 7.24-7.59 (m, 9H), 5.66 (s, 2H), 4.51 (s, 2H), 4.12 (q, 2H), 1.28 (t, 3H)	
509		2-([2-(3-chlorophenyl)-1H-imidazol-1-yl]methyl)-1-ethyl-1H-benzimidazole-5-carbonitrile	8.00 (s, 1H), 7.38-7.72 (m, 7H), 7.24 (s, 1H), 5.71 (s, 2H), 4.50 (s, 2H), 4.20 (q, 2H), 1.22 (t, 3H)	
510		2-([2-(3-chlorophenyl)-1H-imidazol-1-yl]methyl)-1-ethyl-5-fluoro-1H-benzimidazole	(L)-Tartrate salt ¹ H NMR (CD ₃ OD): 7.08-7.60 (m, 9H), 5.66 (s, 2H), 4.51 (s, 2H), 4.12 (s, 2H), 1.18 (t, 3H)	
511		2-([2-(2,5-difluorophenyl)-1H-imidazol-1-yl]methyl)-1-ethyl-5-(4H-1,2,4-triazol-4-yl)-1H-benzimidazole	Hydrochloride ¹ H NMR (d ₆ DMSO): 9.57(2H, s), 8.11(1H, m), 8.01-7.99(1H, m), 7.86(1H, d), 7.78-7.74(1H, m), 7.66-7.52(3H, m), 5.95(2H, s), 4.31(2H, q), 1.26(3H, t)	406.5 [M+1]

TABLE 6				
Cpd#	STRUCTURE	IUPAC NAME	NMR	MS (m/z)
512		2-([2-(2,5-difluorophenyl)-1H-imidazol-1-yl]methyl)-1-ethyl-1H-benzimidazol-5-amine		354.4 [M+1]
513		1-ethyl-5-fluoro-2-([2-(2-fluorophenyl)-1H-imidazol-1-yl]methyl)-1H-benzimidazole	(L)-Tartrate salt 1H NMR (CD3OD): 7.07-7.58 (m, 9H), 5.53 (s, 2H), 4.51 (s, 2H), 4.03 (q, 2H), 1.07 (t, 3H)	
514		1-ethyl-2-([2-(3-fluorophenyl)-1H-imidazol-1-yl]methyl)-1H-benzimidazol-5-amine	1H NMR(CDCl3): 7.38-7.5 (m, 4h), 7.15 (t, 3H), 7.1 (s, 1H), 7.05 (t, 2H), 6.7 (d, 1H), 5.4 (s, 2H), 3.70 (q, 2H), 0.9 (t, 3H)	
515		1-ethyl-2-([2-(3-fluorophenyl)-1H-imidazol-1-yl]methyl)-5-(4H-1,2,4-triazol-4-yl)-1H-benzimidazole		388.2 [M+1]
516		1-ethyl-2-([2-(3-fluorophenyl)-1H-imidazol-1-yl]methyl)-5-(2H-tetrazol-5-yl)-1H-benzimidazole		389.2 [M+1]
517		5-bromo-1-ethyl-2-([2-(3-fluorophenyl)-1H-imidazol-1-yl]methyl)-1H-benzimidazole	1H NMR (CDCl3): 7.93 (d, 1H), 7.39-7.52 (m, 4H), 7.15-7.22 (m, 3H), 7.025 (d, 1H), 5.49 (s, 2H), 3.76 (q, 2H), 0.97 (t, 3H)	
518		2-([2-(2,5-difluorophenyl)-1H-imidazol-1-yl]methyl)-1-ethyl-1H-benzimidazole-5-carbonitrile	1H NMR (CDCl3): 8.09 (d, 1H), 7.56 (m, 1H), 7.33-7.40 (m, 2H), 7.16-7.24 (m, 3H), 7.04 (d, 1H), 5.41 (s, 2H), 3.89 (q, 2H), 1.03 (t, 3H)	
519		2-([2-(3-bromophenyl)-1H-imidazol-1-yl]methyl)-1-ethyl-5-fluoro-1H-benzimidazole	Hydrochloride 1H NMR (d6-DMSO): 7.41-8.00 (m, 8H), 7.18 (m, 1H), 5.92 (s, 2H), 4.28 (q, 2H), 1.27 (t, 3H)	
520		2-([2-(3-bromophenyl)-1H-imidazol-1-yl]methyl)-5-chloro-1-ethyl-1H-benzimidazole	Hydrochloride 1H NMR (d6-DMSO): 7.32-8.00 (m, 9H), 5.95 (s, 2H), 4.29 (q, 2H), 1.27 (t, 3H)	

TABLE 6				
Cpd#	STRUCTURE	IUPAC NAME	NMR	MS (m/z)
521		3-{1-[(1-ethyl-5-fluoro-1H-benzimidazol-2-yl)methyl]-1H-imidazol-2-yl}benzonitrile		346.11 [M+1]
522		1-ethyl-5-fluoro-2-[[2-(3-methylphenyl)-1H-imidazol-1-yl]methyl]-1H-benzimidazole		
523		1-ethyl-5-fluoro-2-[[2-(3-methoxyphenyl)-1H-imidazol-1-yl]methyl]-1H-benzimidazole		351.15 [M+1]
524		4-{1-[(1-ethyl-5-fluoro-1H-benzimidazol-2-yl)methyl]-1H-imidazol-2-yl}-2-fluorophenol		355.12 [M+1]
525		2-[[2-(3-bromophenyl)-1H-imidazol-1-yl]methyl]-1-ethyl-1H-benzimidazole-5-carbonitrile	1H NMR (CDCl3): 8.11 (d, 1H), 7.81 (s, 1H), 7.55-7.67 (m, 3H), 7.35-7.41 (m, 2H), 7.19 (m, 1H), 7.04 (d, 1H), 5.51 (s, 2H), 3.84 (q, 2H), 1.04 (t, 3H)	
526		2-[[2-(3-fluorophenyl)-1H-imidazol-1-yl]methyl]-1-propyl-1H-benzimidazole-5-carbonitrile		360.2 [M+1]; 358.2 [M-1]
527		2-[[2-(3-chlorophenyl)-1H-imidazol-1-yl]methyl]-1-propyl-1H-benzimidazole-5-carbonitrile		376.2 [M+1]; 374.2 [M-1]
528		2-[[2-(2,5-difluorophenyl)-1H-imidazol-1-yl]methyl]-1-propyl-1H-benzimidazole-5-carbonitrile		378.2 [M+1]
529		2-[[2-(3-chloro-4-fluorophenyl)-1H-imidazol-1-yl]methyl]-1-propyl-1H-benzimidazole-5-carbonitrile		394.4 [M+1]; 392.1 [M-1]

TABLE 6				
Cpd#	STRUCTURE	IUPAC NAME	NMR	MS (m/z)
530		2-([2-(2,5-difluorophenyl)-1H-imidazol-1-yl]methyl)-1-ethyl-5-(methylsulfonyl)-1H-benzimidazole		
531		1-ethyl-5-(3-fluorophenyl)-2-([2-(3-fluorophenyl)-1H-imidazol-1-yl]methyl)-1H-benzimidazole	1H NMR (CDCl3): 7.98 (m, 1H), 7.27-7.54 (m, 8H), 7.16-7.22 (m, 2H), 7.01-7.07 (m, 2H), 5.53 (s, 2H), 3.82 (q, 2H), 1.02 (t, 3H)	
532		1-ethyl-2-([2-(3-fluorophenyl)-1H-imidazol-1-yl]methyl)-5-thien-3-yl-1H-benzimidazole	1H NMR (CDCl3): 8.0 (m, 1H), 7.16-7.59 (m, 10H), 7.07 (d, 1H), 5.52 (s, 2H), 3.81 (q, 2H), 1.01 (t, 3H)	
533		5-bromo-2-([2-(3-chlorophenyl)-1H-imidazol-1-yl]methyl)-1-ethyl-1H-benzimidazole	1H NMR (CDCl3): 7.93 (d, 1H), 7.68 (m, 1H), 7.40-7.57 (m, 2H), 7.15-7.19 (m, 2H), 7.03 (d, 1H), 5.48 (s, 2H), 3.77 (q, 2H), 0.98 (t, 3H)	
534		2-([2-(3-cyanophenyl)-1H-imidazol-1-yl]methyl)-1-ethyl-1H-benzimidazole-5-carbonitrile		353.15 [M+1]
535		3-([1-((1-ethyl-5-fluoro-1H-benzimidazol-2-yl)methyl)-1H-imidazol-2-yl]benzoic acid		364.10 [M+1]
536		1-ethyl-5-fluoro-2-([2-(3-pyridin-4-ylphenyl)-1H-imidazol-1-yl]methyl)-1H-benzimidazole		398.18 [M+1]
537		2-([2-(2,5-difluorophenyl)-1H-imidazol-1-yl]methyl)-1-ethyl-5-nitro-1H-benzimidazole	Hydrochloride 1H NMR (d6 DMSO): 8.42(1H, d), 8.17(1H, dd), 8.00(1H, s), 7.90(1H, s), 7.85(1H, d), 7.70(1H, m), 7.56-7.51(2H, m), 5.90 (2H, s), 4.31(2H, q), 1.24(3H, t)	384.1 [M+1]
538		1-ethyl-2-([2-(3-fluorophenyl)-1H-imidazol-1-yl]methyl)-5-nitro-1H-benzimidazole	1H NMR (d6 DMSO): 8.47(1H, d), 8.19(1H, dd), 8.00(1H, m), 7.96(1H, m), 7.88(1H, d), 7.71-7.50(4H, m), 5.98(2H, s), 4.35(2H, q), 1.30(3H, t)	366.2 [M+1]

TABLE 6				
Cpd#	STRUCTURE	IUPAC NAME	NMR	MS (m/z)
539		2-[[2-(3-chlorophenyl)-1H-imidazol-1-yl]methyl]-1-ethyl-5-nitro-1H-benzimidazole		381.95 [M+1]
540		3-{1-[[1-ethyl-5-(methylsulfonyl)-1H-benzimidazol-2-yl]methyl]-1H-imidazol-2-yl}benzonitrile		406.3 [M+1]; 404.3 [M-1]
541		3-{1-[[1-ethyl-5-fluoro-1H-benzimidazol-2-yl]methyl]-1H-imidazol-2-yl}phenol		337.01 [M+1]
542		2-[[2-(3,4-difluorophenyl)-1H-imidazol-1-yl]methyl]-1-ethyl-1H-benzimidazole-5-carbonitrile		379.11 [M+1]
543		2-[[2-(5-bromo-2-fluorophenyl)-1H-imidazol-1-yl]methyl]-1-ethyl-1H-benzimidazole-5-carbonitrile		425.98 [M+1]
544		2-[[2-(2,5-difluorophenyl)-1H-imidazol-1-yl]methyl]-1-ethyl-5-(trifluoromethoxy)-1H-benzimidazole	1H NMR (CDCl3): 7.65 (s, 1H), 7.18-7.41 (m, 6H), 7.06 (s, 1H), 5.35 (s, 2H), 3.82 (q, 2H), 1.0 (t, 3H)	
545		5-chloro-1-ethyl-2-[[2-(3-fluorophenyl)-1H-imidazol-1-yl]methyl]-1H-benzimidazole	1H NMR (CDCl3): 7.22-7.60 (m, 9H), 5.67 (s, 2H), 4.49 (s, 2H), 4.11 (q, 2H), 1.16 (t, 3H)	
546		3-{1-[[5-bromo-1-ethyl-1H-benzimidazol-2-yl]methyl]-1H-imidazol-2-yl}benzonitrile	Hydrochloride 1H NMR (CD3OD): 7.71-8.25 (m, 9H), 6.18 (s, 2H), 4.50 (br s, 2H), 1.50 (br s, 3H)	
547		1-ethyl-2-[[2-(3-nitrophenyl)-1H-imidazol-1-yl]methyl]-1H-benzimidazole-5-carbonitrile		373.15 [M+1]

TABLE 6				
Cpd#	STRUCTURE	IUPAC NAME	NMR	MS (m/z)
548		6-chloro-1-ethyl-2-([2-(3-fluorophenyl)-1H-imidazol-1-yl]methyl)-1H-benzimidazole-5-carbonitrile		380.4 [M+1]; 378.2 [M-1]
549		5-(3,5-difluorophenyl)-1-ethyl-2-([2-(3-fluorophenyl)-1H-imidazol-1-yl]methyl)-1H-benzimidazole	1H NMR (CDCl3): 7.96 (d, 1H), 7.35-7.70 (m, 6H), 7.06-7.23 (m, 4H), 6.78 (m, 1H), 5.52 (s, 2H), 3.81 (q, 2H), 1.00 (t, 3H)	
550		3-(1-([5-(3,5-difluorophenyl)-1-ethyl-1H-benzimidazol-2-yl]methyl)-1H-imidazol-2-yl)benzonitrile	1H NMR (CDCl3): 7.93-8.02 (m, 2H), 7.37-7.77 (m, 6H), 7.08-7.19 (m, 3H), 6.77 (m, 1H), 5.49 (s, 2H), 3.89 (q, 2H), 1.10 (t, 3H)	
551		1-(1-ethyl-2-([2-(3-fluorophenyl)-1H-imidazol-1-yl]methyl)-1H-benzimidazol-5-yl)ethanone	1H NMR (CDCl3): 8.40 (s, 1H), 8.00 (dd, 1H), 7.33-7.53 (m, 4H), 7.17-7.22 (m, 2H), 7.05 (d, 1H), 5.53 (s, 2H), 3.82 (q, 2H), 2.67 (s, 3H), 1.01 (t, 3H)	
552		5-(3,5-dimethylisoxazol-4-yl)-1-ethyl-2-([2-(3-fluorophenyl)-1H-imidazol-1-yl]methyl)-1H-benzimidazole	1H NMR (CDCl3): 7.66 (d, 1H), 7.34-7.54 (m, 4H), 7.15-7.22 (m, 3H), 7.06 (d, 1H), 5.52 (s, 2H), 3.83 (q, 2H), 2.41 (s, 3H), 2.28 (s, 3H), 1.02 (s, 3H)	
553		1-ethyl-2-([2-(3-(trifluoromethoxy)phenyl)-1H-imidazol-1-yl]methyl)-1H-benzimidazole-5-carbonitrile		411.96 [M+1]
554		(1E)-1-(1-ethyl-2-([2-(3-fluorophenyl)-1H-imidazol-1-yl]methyl)-1H-benzimidazol-5-yl)ethanone O-methyloxime	Syn and anti 1H NMR (CDCl3): 7.98 (d, 1H), 7.94 (d, 1H), 7.75 (dd, 1H+H1), 7.14-7.22 (m, 2H+H1), 7.32-7.52 (m, 4H+H1), [7.03 (d, 1H), 7.01 (d, 1H)], 5.49 (s, 2H+H1), [4.00 (s, 2H), 3.86 (s, 2H)], 3.79 (q, 2H+H1), [2.30 (s, 3H), 2.26 (s, 3H)], 0.98 (t, 3H+H1)	
555		2-([2-(2,5-difluorophenyl)-1H-imidazol-1-yl]methyl)-1-ethyl-5-[5-(trifluoromethyl)-1H-tetrazol-1-yl]-1H-benzimidazole		475.1 [M+1]
556		1-ethyl-2-([2-(3-fluorophenyl)-1H-imidazol-1-yl]methyl)-1H-benzimidazole-5,6-dicarbonitrile		371.0 [M+1]

TABLE 6				
Cpd#	STRUCTURE	IUPAC NAME	NMR	MS (m/z)
557		1-ethyl-2-[(2-pyridin-3-yl-1H-imidazol-1-yl)methyl]-1H-benzimidazole-5-carbonitrile	Dihydrochloride ¹ H NMR (d6 DMSO): 8.96(1H, s), 8.82 (1H, d), 8.24(1H, d), 8.14(1H, s), 8.05(1H, d), 7.98(1H, d), 7.85(1H, d), 7.69-7.65(2H, m), 6.01(2H, s), 4.33(2H, q), 1.29(3H, t)	
558		1-ethyl-2-[(2-pyridin-2-yl-1H-imidazol-1-yl)methyl]-1H-benzimidazole-5-carbonitrile	Dihydrochloride ¹ H NMR (d6 DMSO): 8.55(1H, d), 8.41 (1H, d), 8.10-7.98(4H, m), 7.85(1H, d), 7.64(1H, d), 7.54-7.51(1H, m), 6.39 (2H, s), 4.45(2H, q), 1.41(3H, t)	329.4 [M+1]
559		1-ethyl-5-fluoro-2-[(2-(3-nitrophenyl)-1H-imidazol-1-yl)methyl]-1H-benzimidazole		365.99 [M+1]
560		1-ethyl-2-[(2-(3-methylphenyl)-1H-imidazol-1-yl)methyl]-1H-benzimidazole-5-carbonitrile		343.06 [M+1]
561		1-(3-{1-[(1-ethyl-5-fluoro-1H-benzimidazol-2-yl)methyl]-1H-imidazol-2-yl}phenyl)ethanone		363.05 [M+1]
562		3-{1-[(5-(3,5-dimethylisoxazol-4-yl)-1-ethyl-1H-benzimidazol-2-yl)methyl]-1H-imidazol-2-yl}benzonitrile	¹ H NMR (CDCl ₃): 7.94-8.01 (m, 2H), 7.75 (m, 1H), 7.61-7.66 (m, 2H), 7.38 (dd, 1H), 7.18-7.22 (m, 2H), 7.10 (d, 1H), 5.49 (s, 2H), 3.90 (q, 2H), 2.42 (s, 3H), 2.29 (s, 3H), 1.12 (t, 3H)	
563		2-[(2-(2,5-difluorophenyl)-1H-imidazol-1-yl)methyl]-1-ethyl-5-(1H-imidazol-2-yl)-1H-benzimidazole	Dihydrochloride ¹ H NMR (d6 DMSO): 8.43(1H, s), 8.13-8.11(2H, m), 7.99(1H, d), 7.89(1H, d), 7.79-7.75(3H, m), 7.60-7.51(3H, m), 5.95(2H, s), 4.31(2H, q), 1.26 (3H, t)	405.3 [M+1]
564		1-ethyl-5-nitro-2-[(2-pyridin-2-yl-1H-imidazol-1-yl)methyl]-1H-benzimidazole		349.1 [M+1]
565		2-[(2-(2,5-difluorophenyl)-1H-imidazol-1-yl)methyl]-1-ethyl-5-(1-ethyl-1H-imidazol-2-yl)-1H-benzimidazole		433.2 [M+1]










TABLE 6				
Cpd#	STRUCTURE	IUPAC NAME	NMR	MS (m/z)
566		3-({1-[(5-acetyl-1-ethyl-1H-benzimidazol-2-yl)methyl]-1H-imidazol-2-yl}benzonitrile	¹ H NMR (CDCl ₃): 8.40 (d, 1H), 7.93-8.05 (m, 3H), 7.76 (m, 1H), 7.63 (m, 1H), 7.37 (dd, 1H), 7.21 (d, 1H), 7.09 (d, 1H), 5.50 (s, 2H), 3.91 (q, 2H), 2.68 (s, 3H), 1.11 (t, 3H)	
567		1-ethyl-2-({2-[3-(trifluoromethyl)phenyl]-1H-imidazol-1-yl)methyl}-1H-benzimidazole-5-carbonitrile		396.06 [M+1]
568		1-ethyl-2-({2-thien-2-yl-1H-imidazol-1-yl)methyl}-1H-benzimidazole-5-carbonitrile		334.04 [M+1]
569		3-({1-ethyl-5-(trifluoromethoxy)-1H-benzimidazol-2-yl)methyl}-1H-imidazol-2-yl}benzonitrile		412.2 [M+1]; 410.2 [M-1]
570		3-({1-ethyl-5-fluoro-1H-benzimidazol-2-yl)methyl}-1H-imidazol-2-yl}benzaldehyde		349.17 [M+1]
571		1-ethyl-2-({2-[3-fluorophenyl]-1H-imidazol-1-yl)methyl}-N-methoxy-N-methyl-1H-benzimidazole-5-carboxamide	¹ H NMR (CDCl ₃): 8.19 (d, 1H), 7.71 (dd, 1H), 7.40-7.55 (m, 3H), 7.32 (dd, 1H), 7.16-7.23 (m, 2H), 7.05 (d, 1H), 5.53 (s, 2H), 3.80 (q, 2H), 3.59 (s, 3H), 3.41 (s, 3H), 1.00 (t, 3H)	
572		1-ethyl-2-({2-[3-fluorophenyl]-1H-imidazol-1-yl)methyl}-5-(1,3,4-oxadiazol-2-yl)-1H-benzimidazole		
573		1-ethyl-2-({2-[3-fluorophenyl]-1H-imidazol-1-yl)methyl}-5-(5-methyl-1,3,4-oxadiazol-2-yl)-1H-benzimidazole	Hydrochloride ¹ H NMR (d ₆ DMSO): 8.11 (1H, s), 8.00(1H, d), 7.95(1H, d), 7.90-7.82(2H, m), 7.74-7.72(1H, m), 7.64-7.58(2H, m), 7.53-7.49(1H, m), 5.96(2H, s), 4.329(2H, q), 2.56(3H, s), 1.30(3H, t)	403.3 [M+1]
574		1-ethyl-2-({2-[3-fluorophenyl]-1H-imidazol-1-yl)methyl}-5-(5-methyl-1,2,4-oxadiazol-3-yl)-1H-benzimidazole		403.8 [M+1]; 401.4 [M-1]

TABLE 6				
Cpd#	STRUCTURE	IUPAC NAME	NMR	MS (m/z)
575		1-ethyl-2-([2-(3-fluorophenyl)-1H-imidazol-1-yl]methyl)-5-[[trifluoromethyl]sulfonyl]-1H-benzimidazole		453.8 [M+1]; 451.2 [M-1]
576		1-(2-([2-(3-chlorophenyl)-1H-imidazol-1-yl]methyl)-1-ethyl-1H-benzimidazol-5-yl)ethanone	1H NMR (CDCl3): (8.40 (d, 1H), 8.01 (dd, 1H), 7.68 (m, 1H), 7.55 (m, 1H), 7.45-7.52 (m, 2H), 7.35 (d, 1H), 7.17 (d, 1H), 7.06 (d, 1H), 5.52 (s, 2H), 3.85 (q, 2H), 2.68 (s, 3H), 1.03 (t, 3H)	
577		1-ethyl-2-([2-(3-fluorophenyl)-1H-imidazol-1-yl]methyl)-1H-benzimidazole-6-carbonitrile		346.2 [M+1]; 344.3 [M-1]
578		methyl 3-(1-([5-cyano-1-ethyl-1H-benzimidazol-2-yl]methyl)-1H-imidazol-2-yl)benzoate		386.17 [M+1]
579		1-ethyl-2-([2-(3-fluorophenyl)-1H-imidazol-1-yl]methyl)-1H-imidazo[4,5-c]pyridine	1H NMR (CDCl3): 9.10 (s, 1H), 8.44 (d, 1H), 7.38-7.52 (m, 3H), 7.18-7.20 (m, 2H), 7.16 (s, 1H), 7.04 (d, 1H), 5.53 (s, 2H), 3.79 (q, 2H), 1.00 (t, 3H)	
580				379.05 [M+1]
581		2-([2-(5-aminothien-3-yl)-1H-imidazol-1-yl]methyl)-1-ethyl-1H-benzimidazole-5-carbonitrile		397.12 [M+1]
582		1-ethyl-2-([2-(3-nitrophenyl)-1H-imidazol-1-yl]methyl)-1H-benzimidazole-5-carboxamide		391.17 [M+1]
583		1-ethyl-2-([2-(3-fluorophenyl)-1H-imidazol-1-yl]methyl)-5-(5-methyl-1,3-oxazol-2-yl)-1H-benzimidazole	Hydrochloride 1H NMR (d6 DMSO): 8.05(1H, s), 7.99(1H, d), 7.97(1H, s), 7.88-7.86(1H, m), 7.76-7.72(2H, m), 7.65-7.59(2H, m), 7.54-7.51(1H, m), 6.99(1H, s), 5.939(2H, s), 4.31(2H, q), 2.36(3H, s), 1.29(3H, t).	402.5 [M+1]

TABLE 6				
Cpd#	STRUCTURE	IUPAC NAME	NMR	MS (m/z)
584		1-ethyl-2-([2-(3-fluorophenyl)-1H-imidazol-1-yl]methyl)-5-(5-pyridin-3-yl-1,3,4-oxadiazol-2-yl)-1H-benzimidazole	Hydrochloride ¹ H NMR (d6 DMSO): 9.36(1H, d), 8.84(1H, dd), 8.57(1H, m), 8.44(1H, s), 8.10(1H, dd), 8.03(1H, d), 7.98(1H, d), 7.90(1H, d), 7.75-7.51(5H, m), 5.99(2H, s), 4.35(2H, q), 1.33(3H, t).	466.5 [M+1]
585		(1E)-1-(2-([2-(3-chlorophenyl)-1H-imidazol-1-yl]methyl)-1-ethyl-1H-benzimidazol-5-yl)ethanone oxime	¹ H NMR (CDCl ₃): 11.03 (s, 1H), 7.81 (s, 1H), 7.52-7.67 (m, 4H), 7.40-7.45 (m, 2H), 7.31 (s, 1H), 7.07 (s, 1H), 5.67 (s, 2H), 4.17 (q, 2H), 2.18 (s, 3H), 1.16 (t, 3H)	
586		(1E)-1-(2-([2-(3-chlorophenyl)-1H-imidazol-1-yl]methyl)-1-ethyl-1H-benzimidazol-5-yl)ethanone O-methyloxime	Syn and anti ¹ H NMR (CDCl ₃): 7.99 (dd, 1H), 7.95 (dd, 1H), 7.76 (dd, 1H+H1), 7.65 (m, 1H+H1), 7.44-7.57 (m, 3H+H1), 7.27 (d, 1H+H1), 7.15 (d, 1H+H1), 7.04 (d, 1H+H1), 5.49 (s, 2H+H1), 4.01 (s, 3H), 3.87 (s, 3H), 3.80 (q, 2H+H1), [2.30 (s, 3H), 2.27 (s, 3H)], 1.00 (t, 3H+H1)	
587		(1E)-1-(2-([2-(3-chlorophenyl)-1H-imidazol-1-yl]methyl)-1-ethyl-1H-benzimidazol-5-yl)ethanone O-ethyloxime	Syn and anti ¹ H NMR (CDCl ₃): 8.00 (dd, 1H+H1), 7.76 (dd, 1H+H1), 7.69 (m, 1H+H1), 7.44-7.57 (m, 3H+H1), 7.27 (dd, 1H+H1), 7.15 (d, 1H+H1), 7.04 (d, 1H), 7.03 (d, 1H), 5.49 (s, 2H+H1), 4.26 (q, 2H), 4.13 (q, 2H), 3.80 (q, 2H+H1), 2.27 (s, 3H), 1.34 (t, 3H), 1.25 (t, 3H), 1.00 (t, 3H+H1)	
588		1-ethyl-2-([2-(6-fluoropyridin-2-yl)-1H-imidazol-1-yl]methyl)-1H-benzimidazole-5-carbonitrile		347.07 [M+1]
589		1-ethyl-2-([2-(3-fluorophenyl)-1H-imidazol-1-yl]methyl)-5-(1,2,4-oxadiazol-3-yl)-1H-benzimidazole		
590		1-ethyl-2-([2-(6-methoxypyridin-2-yl)-1H-imidazol-1-yl]methyl)-1H-benzimidazole-5-carbonitrile		359.08 [M+1]
591		(1E)-1-(1-ethyl-2-([2-(3-fluorophenyl)-1H-imidazol-1-yl]methyl)-1H-benzimidazol-5-yl)ethanone oxime	Hydrochloride ¹ H NMR (d6-DMSO): 11.10 s, 1H), 7.52-7.99 (m, 9H+H1), 5.923 (s, 2H+H1): 4.30 (q, 2H+H1), 2.18 (s, 3H), 2.14 (s, 3H), 1.29 (t, 3H+H1)	
592		(1E)-1-(1-ethyl-2-([2-(3-fluorophenyl)-1H-imidazol-1-yl]methyl)-1H-benzimidazol-5-yl)ethanone O-ethyloxime	Hydrochloride ¹ H NMR (d6-DMSO): 7.52-7.99 (m, 9H), 5.93 (s, 2H), 4.28 (q, 2H), 4.14 (q, 2H), 2.20 (s, 3H), 1.23-1.29 (m, 6H)	










TABLE 6				
Cpd#	STRUCTURE	IUPAC NAME	NMR	MS (m/z)
593		1-(1-ethyl-2-([2-(3-fluorophenyl)-1H-imidazol-1-yl]methyl)-1H-benzimidazol-5-yl)propan-1-one	¹ H NMR (CDCl ₃): 8.42 (d, 1H), 8.02 (dd, 1H), 7.32-7.54 (m, 4H), 7.17-7.24 (m, 2H), 7.06 (d, 1H), 5.53 (s, 2H), 3.83 (q, 2H), 3.09 (q, 2H), 1.27 (t, 3H), 1.02 (t, 3H)	
594		ethyl 1-(1-ethyl-2-([2-(3-fluorophenyl)-1H-imidazol-1-yl]methyl)-1H-benzimidazol-5-yl)-1H-1,2,3-triazole-4-carboxylate		
595		3-ethyl-2-([2-(3-fluorophenyl)-1H-imidazol-1-yl]methyl)-3H-imidazo[4,5-c]pyridine		
596		1-(1-ethyl-2-([2-(6-fluoropyridin-2-yl)-1H-imidazol-1-yl]methyl)-1H-benzimidazol-5-yl)ethanone		364.2 [M+1]; 362.6 [M-1]
597		1-ethyl-2-([2-(6-fluoropyridin-2-yl)-1H-imidazol-1-yl]methyl)-5-(trifluoromethyl)-1H-benzimidazole		390.2 [M+1]; 388.2 [M-1]
598		2-([2-(6-fluoropyridin-2-yl)-1H-imidazol-1-yl]methyl)-1-propyl-1H-benzimidazole-5-carbonitrile	¹ H NMR (399.96MHz, CDCl ₃): d 8.17 (dd, J = 2.0, 7.6 Hz, 1H), 8.05 (s, 1H), 7.88 (q, J = 8.0 Hz, 1H), 7.52 (d, J = 8.4 Hz, 1H), 7.41 (d, J = 8.4 Hz, 1H), 7.21 (s, 1H), 7.18 (s, 1H), 6.28 (s, 2H), 4.28 (t, J = 7.6 Hz, 2H), 1.68 (dt, J = 7.6 Hz, 2H), 0.84 (t, J = 7.6 Hz, 3H)	361.2 [M+1]; 359.2 [M-1]
599		1-ethyl-2-([2-(3-fluorophenyl)-1H-imidazol-1-yl]methyl)-1H-imidazo[4,5-b]pyridine		322.2 [M+1]; 320.3 [M-1]
600		2-([2-(3-fluorophenyl)-1H-imidazol-1-yl]methyl)-1-propyl-1H-imidazo[4,5-c]pyridine	¹ H NMR (CDCl ₃): 9.11 (s, 1H), 8.45 (d, 1H), 7.38-7.52 (m, 3H), 7.16-7.26 (m, 3H), 7.06 (s, 1H), 5.53 (s, 2H), 3.71 (t, 2H), 1.43 (q 2H), 0.75 (t, 3H)	
601		2-([2-(3-chlorophenyl)-1H-imidazol-1-yl]methyl)-1-ethyl-5-isoxazol-5-yl-1H-benzimidazole	¹ H NMR (CDCl ₃): 8.30 (d, 1H), 8.21 (dd, 1H), 7.78 (dd, 1H), 7.69 (m, 1H), 7.56 (m, 1H), 7.45-7.48 (m, 2H), 7.39 (dd, 1H), 7.17 (d, 1H), 7.07 (d, 1H), 6.53 (d, 1H), 5.52 (s, 2H), 3.83 (q, 2H), 1.03 (t, 3H)	

TABLE 6				
Cpd#	STRUCTURE	IUPAC NAME	NMR	MS (m/z)
602		1-(2-fluoroethyl)-2-([2-(3-fluorophenyl)-1H-imidazol-1-yl]methyl)-1H-benzimidazole		339.2 [M+1]
603		2-([2-(3-fluorophenyl)-1H-imidazol-1-yl]methyl)-1-propyl-1H-imidazo[4,5-b]pyridine		336.2 [M+1]; 334.2 [M-1]
604		2-([2-(2,5-difluorophenyl)-1H-imidazol-1-yl]methyl)-1-ethyl-1H-imidazo[4,5-b]pyridine		
605		1-ethyl-2-([2-(5-[hydroxy(oxido)amino]thien-3-yl)-1H-imidazol-1-yl]methyl)-1H-benzimidazole-5-carbonitrile		
606		3-propyl-2-([2-(2-pyrimidin-2-yl-1H-imidazol-1-yl)methyl]-3H-imidazo[4,5-b]pyridine	(CDCl3) 8.46 (d, 1H), 8.39 (dd, 1H), 8.03 (dd, 1H), 7.27 (m, 5H), 6.37 (s, 2H), 4.27 (t, 2H), 1.70 (p, 2H), 0.82 (t, 3H)	320.4 [M+1]
607		3-ethyl-2-([2-(2,3,4-trifluorophenyl)-1H-imidazol-1-yl]methyl)-3H-imidazo[4,5-b]pyridine	(CDCl3) 8.40 (dd, 1H), 8.02 (dd, 1H), 7.36 (dq, 1H), 7.25 (dd, 1H), 7.22 (d, 1H), 7.01 (m, 2H), 5.38 (s, 2H), 4.03 (q, 2H), 1.10 (t, 3H)	358.2 [M+1]
608		3-propyl-2-([2-(2-thien-2-yl-1H-imidazol-1-yl)methyl]-3H-imidazo[4,5-b]pyridine	(CDCl3) 8.40 (dd, 1H), 8.05 (dd, 1H), 7.46 (dd, 1H), 7.25 (dd, 1H), 7.15 (m, 2H), 7.03 (d, 1H), 5.61 (s, 2H), 3.96 (t, 2H), 1.55 (p, 2H), 0.80 (t, 3H)	324.3 [M+1]
609		3-ethyl-2-([2-(1-methyl-1H-pyrazol-3-yl)-1H-imidazol-1-yl]methyl)-3H-imidazo[4,5-b]pyridine	(CDCl3) 8.39 (dd, 1H), 8.05 (dd, 1H), 7.43 (dd, 1H), 7.23 (dd, 1H), 7.10 (dd, 1H), 7.04 (dd, 1H), 6.91 (dd, 1H), 6.21 (s, 2H), 4.33 (q, 2H), 3.97 (3, 3H), 1.13 (t, 3H)	m/z 308.3 [M+1]

TABLE 6				
Cpd#	STRUCTURE	IUPAC NAME	NMR	MS (m/z)
610		3-ethyl-2-([2-(2-fluoro-5-methylphenyl)-1H-imidazol-1-yl]methyl)-3H-imidazo[4,5-b]pyridine	(CDCl ₃) 8.39 (dd, J = 4.8, 1.5 Hz, 1H), 8.05 (dd, J = 8.4, 1.5 Hz, 1H), 7.44 (dd, J = 6.6, 2.1 Hz, 1H), 7.25-7.30 (m, 2H), 7.19 (d, J = 1.5 Hz, 1H), 7.12 (dd, J = 9.9, 8.7 Hz, 1H), 7.04 (d, J = 1.5 Hz, 1H), 5.38 (s, 2H), 3.96 (q, J = 7.2 Hz, 2H), 2.37 (s, 3H), 1.03 (t, J = 7.2 Hz, 3H)	336.2 [M+1]
611		3-ethyl-2-([2-(5-fluoro-2-methylphenyl)-1H-imidazol-1-yl]methyl)-3H-imidazo[4,5-b]pyridine	(CDCl ₃) 8.39 (dd, J = 4.6, 1.5 Hz, 1H), 8.04 (dd, J = 8.0, 1.5 Hz, 1H), 7.19-7.32 (m, 3H), 7.07-7.13 (m, 3H), 5.26 (s, 2H), 3.98 (q, J = 7.2 Hz, 2H), 2.24 (s, 3H), 1.11 (t, J = 7.2 Hz, 3H)	m/z 336.2 [M+1]
612		2-([2-(3-chloro-2,6-difluorophenyl)-1H-imidazol-1-yl]methyl)-3-ethyl-3H-imidazo[4,5-b]pyridine	(CDCl ₃) 8.40 (dd, J = 4.7, 1.5 Hz, 1H), 8.01 (dd, J = 8.0, 1.5 Hz, 1H), 7.52 (m, 1H), 7.23-7.30 (m, 2H), 7.11 (d, J = 1.5 Hz, 1H), 7.03 (m, 1H), 5.34 (s, 2H), 4.03 (q, J = 7.2 Hz, 2H), 1.10 (t, J = 7.2 Hz, 3H)	m/z 374.2 [M+1]
613		3-ethyl-2-([2-(2-methoxyphenyl)-1H-imidazol-1-yl]methyl)-3H-imidazo[4,5-b]pyridine	(CDCl ₃) 8.38 (dd, J = 4.8, 1.5 Hz, 1H), 8.03 (dd, J = 8.1, 1.5 Hz, 1H), 7.46-7.57 (m, 2H), 7.25 (m, 1H), 7.16 (d, J = 1.2 Hz, 1H), 7.13 (dt, J = 7.5, 1.2 Hz, 1H), 7.04 (d, J = 8.4 Hz, 1H), 7.00 (d, J = 1.2 Hz, 1H), 5.29 (s, 2H), 3.90 (q, J = 7.2 Hz, 2H), 3.84 (s, 3H), 0.98 (t, J = 7.2 Hz, 3H)	m/z 334.2 [M+1]
614		3-ethyl-2-([2-(2,4,5-trifluorophenyl)-1H-imidazol-1-yl]methyl)-3H-imidazo[4,5-b]pyridine	(CDCl ₃) 8.40 (dd, J = 4.6, 1.4 Hz, 1H), 8.04 (dd, J = 8.2, 1.5 Hz, 1H), 7.50 (m, 1H), 7.25 (dd, J = 7.8, 5.1 Hz, 1H), 7.21 (d, J = 1.5 Hz, 1H), 7.14 (m, 1H), 7.06 (d, J = 1.5 Hz, 1H), 5.37 (s, 2H), 4.02 (q, J = 7.2 Hz, 2H), 1.11 (t, J = 7.2 Hz, 3H)	m/z 358.2 [M+1]
615		4-chloro-2-([2-(6-fluoropyridin-2-yl)-1H-imidazol-1-yl]methyl)-6-methyl-1-propyl-1H-imidazo[4,5-c]pyridine	(CDCl ₃) 8.19(dd, J = 6, 1.2 Hz, 1H), 7.93(q, J = 6 Hz, 1H), 7.21(d, J = 1 Hz, 1H), 7.14(d, J = 1 Hz, 1H), 7.05(s, 1H), 6.93(dd, J = 6, 2 Hz, 1H), 6.38(s, 2H), 4.18(t, J = 5.6 Hz, 2H), 2.63(s, 3H), 1.57(sextet, J = 5.4 Hz, 2H), 0.77(t, J = 5.4 Hz, 3H)	m/z 385 [M+1]
616		4-chloro-6-methyl-1-propyl-2-([2-(pyridin-2-yl)-1H-imidazol-1-yl]methyl)-1H-imidazo[4,5-c]pyridine	(CD ₃ OD) 8.02-7.98(m, 3H), 7.34(d, J = 2 Hz, 1H), 7.14(d, J = 1.8 Hz, 1H), 7.01(d, J = 2 Hz, 1H), 6.47(s, 1H), 6.12(s, 2H), 4.21(t, J = 5.2 Hz, 2H), 2.33(s, 3H), 1.68(sextet, J = 5.4 Hz, 2H), 0.86(t, J = 5.4 Hz, 3H)	m/z 367 [M+1]
617		3-ethyl-2-([2-(5-fluoro-2-methoxyphenyl)-1H-imidazol-1-yl]methyl)-3H-imidazo[4,5-b]pyridine	(CDCl ₃) 8.39 (dd, J = 4.6, 1.4 Hz, 1H), 8.04 (dd, J = 8.0, 1.4 Hz, 1H), 7.15-7.32 (m, 4H), 7.01 (d, J = 1.5 Hz, 1H), 6.96 (dd, J = 9.0, 4.5 Hz, 1H), 5.30 (s, 2H), 3.95 (q, J = 7.2 Hz, 2H), 3.80 (s, 3H), 1.02 (t, J = 7.2 Hz, 3H)	m/z 352.3 [M+1]

TABLE 6				
Cpd#	STRUCTURE	IUPAC NAME	NMR	MS (m/z)
618		2-[[2-(6-fluoropyridin-2-yl)-1H-imidazol-1-yl]methyl]-5-methyl-3-propyl-3H-imidazo[4,5-b]pyridine	(CDCl ₃) 8.18 (dd, J = 7.8, 2.1 Hz, 1H), 7.84-7.92 (m, 2H), 7.21 (d, J = 1.2 Hz, 1H), 7.18 (d, J = 1.2 Hz, 1H), 7.06 (d, J = 7.8 Hz, 1H), 6.89 (dd, J = 8.1, 2.7 Hz, 1H), 6.29 (s, 2H), 4.29 (t, J = 7.5 Hz, 2H), 2.64 (s, 3H), 1.72 (m, 2H), 0.83 (t, J = 7.5 Hz, 3H)	
619		3-ethyl-2-[[2-(2,3,5-trifluorophenyl)-1H-imidazol-1-yl]methyl]-3H-imidazo[4,5-b]pyridine	(CDCl ₃) 8.40 (dd, J = 4.8, 1.2 Hz, 1H), 8.04 (dd, J = 7.8, 1.2 Hz, 1H), 7.05-7.28 (m, 5H), 5.40 (s, 2H), 4.04 (q, J = 7.3 Hz, 2H), 1.10 (t, J = 7.3 Hz, 3H)	
620		2-[[2-(1H-benzimidazol-5-yl)-1H-imidazol-1-yl]methyl]-3-propyl-3H-imidazo[4,5-b]pyridine	(CDCl ₃): 8.38 (d, 1H), 8.02 (m, 2H), 7.96 (s, 1H), 7.66 (s, 1H), 7.42 (d, 1H), 7.25 (m, 2H), 5.56 (s, 2H), 5.29 (1H), 3.87 (t, 2H), 1.48 (m, 2H), 0.68 (t, 3H)	m/z 358 [M+1]
621		3-[[1-[(3-ethyl-3H-imidazo[4,5-b]pyridin-2-yl)methyl]-1H-imidazol-2-yl]-4-fluorobenzonitrile	(CDCl ₃): 8.40 (d, 1H), 7.96 (m, 2H), 7.75 (m, 1H), 7.31 (m, 3H), 7.10 (s, 1H), 5.38 (s, 2H), 4.03 (q, 2H), 1.11 (t, 3H)	
622		2-[[2-(2,6-difluoro-3-methylphenyl)-1H-imidazol-1-yl]methyl]-1-ethyl-1H-benzimidazole-5-carbonitrile	(CDCl ₃) 8.06 (s, 1H), 7.55 (m, 1H), 7.37 (dd, J=8.0, 1.2 Hz, 1H), 7.32-7.26 (m, 2H), 7.04 (s, 1H), 6.94 (t, J=8.8 Hz, 1H), 5.34 (s, 2H), 3.88 (q, J=7.1 Hz, 2H), 2.25 (s, 3H), 1.03 (t, J=7.1 Hz, 3H)	m/z 378 [M+1]
623		1-[[1-[(3-ethyl-3H-imidazo[4,5-b]pyridin-2-yl)methyl]-1H-imidazol-2-yl]isoquinoline	(CDCl ₃) 9.11 (d, 1H), 8.58 (d, 1H), 8.37 (dd, 1H), 8.02 (dd, 1H), 7.86 (dd, 1H), 7.51 (m, 3H), 7.32 (d, 1H), 7.22 (m, 2H), 6.11 (s, 2H), 4.36 (q, 2H), 1.11 (t, 3H)	m/z 355.2 [M+1]
624		2-[[2-(2-fluoro-5-methylphenyl)-1H-imidazol-1-yl]methyl]-1-propyl-1H-imidazo[4,5-c]pyridine	(CDCl ₃) 9.07(s, 1H), 8.41(d, J = 4.5 Hz, 1H), 8.12(dd, J = 5.1, 1.5 Hz, 1H), 7.28-7.24(m, 2H), 7.21(d, J = 4.2 Hz, 1H), 7.10(t, J = 6.6 Hz, 1H), 7.01(s, 1H), 5.37(s, 2H), 3.69(t, J = 5.7 Hz, 2H), 2.34(s, 3H), 1.40(sextet, J = 5.7 Hz, 2H), 0.74(t, J = 5.7 Hz, 3H)	m/z 350 [M+1]
625		1-ethyl-2-[[2-(2-fluoro-5-methylphenyl)-1H-imidazol-1-yl]methyl]-1H-imidazo[4,5-c]pyridine	(CDCl ₃) 9.09(s, 1H), 8.44(d, J = 4.8 Hz, 1H), 7.42(d, J = 5.0 Hz, 1H), 7.29-7.23(m, 3H), 7.12(t, J = 6.3 Hz, 1H), 7.00(s, 1H), 5.39(s, 2H), 3.81(q, J = 5.4 Hz, 2H), 2.36(s, 3H), 0.99(t, J = 5.1 Hz, 3H)	m/z 336 [M+1]

TABLE 6				
Cpd#	STRUCTURE	IUPAC NAME	NMR	MS (m/z)
626		2-((2-(3-chloro-2,6-difluorophenyl)-1H-imidazol-1-yl)methyl)-1-ethyl-1H-imidazo[4,5-c]pyridine	D6dmsol 9.42 (s, 1H), 8.69 (d, J=6.4Hz, 1H), 8.36 (d, J=6.4Hz, 1H), 7.99-7.91 (m, 2H), 7.75 (s, 1H), 7.39 (t, J=8Hz, 1H), 5.95 (s, 2H), 4.43 (q, J=7.2Hz, 2H), 1.25 (t, J=7.2Hz, 3H)	m/z 374 [M+1]
627		1-ethyl-2-(6-((1-ethyl-1H-imidazo[4,5-c]pyridin-2-yl)methyl)-1H-imidazol-2-yl)pyridin-2-yl)-1H-imidazo[4,5-c]pyridine	(D6dmsol) 9.57 (s, 1H), 9.34 (s, 1H), 8.74 (d, J=6.8Hz, 1H), 8.63 (d, J=6.4Hz, 1H), 8.42 (d, J=6.8Hz, 1H), 8.33 (d, J=6.4Hz, 1H), 8.25 (t, J=8Hz, 1H), 8.17 (d, J=6.8Hz, 1H), 8.07 (d, J=7.6Hz, 1H), 7.70 (s, 1H), 7.46 (s, 1H), 6.32 (s, 2H), 4.24-4.16 (m, 4H), 0.97 (t, J=7.2Hz, 3H), 0.89 (t, J=7.2Hz, 3H)	
628		2-((2-(2,6-difluoro-3-methylphenyl)-1H-imidazol-1-yl)methyl)-1-ethyl-1H-imidazo[4,5-c]pyridine	(d6 DMSO) 9.42 (s, 1H), 8.69 (d, J=6.4Hz, 1H), 8.37 (d, J=6.8Hz, 1H), 8.08 (s, 1H), 7.90 (s, 1H), 7.67-7.61 (m, 1H), 7.24 (t, J=8.8Hz, 1H), 5.98 (s, 2H), 4.44 (q, J=7.2Hz, 2H), 2.19 (s, 3H), 1.27(t, J=7.2Hz, 3H)	
629		3-((1-((3-ethyl-3H-imidazo[4,5-b]pyridin-2-yl)methyl)-1H-imidazol-2-yl)pyridin-2(1H)-one	(d6 DMSO) 12.00 (br s, 1H), 8.28 (dd, J=5.2, 1.2Hz, 1H), 7.93 (dd, J=8, 1.2Hz, 1H), 7.56 (dd, J=6.8, 2Hz, 1H), 7.45 (dd, J=6.8, 2Hz, 1H), 7.25 (s, 1H), 7.20 (dd, J=8, 4.8Hz, 1H), 6.98 (s, 1H), 6.19 (t, J=5.2Hz, 1H), 5.64 (s, 2H), 4.14 (q, J=7.2Hz, 2H), 1.07 (t, J=7.2Hz, 3H)	m/z 319 [M - H]
630		2-((2-(6-fluoropyridin-2-yl)-1H-imidazol-1-yl)methyl)-3,5-dimethyl-3H-imidazo[4,5-b]pyridine	(CDCl3) 8.16 (dd, 1H), 7.88(m, 2H), 7.16 (m, 2H), 7.06 (d, 1H), 6.85 (dd, 1H), 6.24 (s, 2H), 3.85 (s, 3H), 2.62 (s, 3H)	
631		8-((2-(6-fluoropyridin-2-yl)-1H-imidazol-1-yl)methyl)-2,9-dimethyl-9H-purine	(CDCl3) 8.92 (s, 1H), 8.18 (dd, 1H), 7.88 (q, 1H), 7.25 (s, 1H), 7.20 (s, 1H), 7.11 (s, 1H), 6.83 (dd, 1H), 6.22 (s, 2H), 3.89 (s, 3H), 2.8 (s, 3H)	m/z 324 [M+1]
632		2-((2-(6-fluoropyridin-2-yl)-1H-imidazol-1-yl)methyl)-1,6-dimethyl-1H-imidazo[4,5-c]pyridine	(CDCl3) 8.91 (s, 1H), 8.16 (dd, J = 7.6, 2.0 Hz, 1H), 7.85 (q, J = 8.0 Hz, 1H), 7.20 (s, 1H), 7.18 (s, 1H), 7.11 (s, 1H), 6.84 (dd, J = 8.0, 2.8 Hz, 1H), 6.22 (s, 2H), 3.80 (s, 3H), 2.66 (s, 3H)	m/z 323 [M+1]
633		6-chloro-1-ethyl-2-((2-(6-fluoropyridin-2-yl)-1H-imidazol-1-yl)methyl)-1H-benzimidazole	(CDCl3) 8.17 (dd, 1H), 7.89 (q, 1H), 7.25 (s, 1H), 7.75 (s, 1H), 7.26 (m, 2H), 7.19 (s, 1H), 7.17 (s, 1H), 6.89 (dd, 1H), 6.28 (s, 2H), 4.33 (q, 2H), 1.20 (s, 3H)	m/z 356 [M+1]

Cpd#	STRUCTURE	IUPAC NAME	NMR	MS (m/z)
634		1-ethyl-2-([2-(6-fluoropyridin-2-yl)-1H-imidazol-1-yl]methyl)-6-(trifluoromethyl)-1H-benzimidazole	(CDCl ₃) 8.20 (m, 1H), 7.87(m, 2H), 7.63 (s, 1H), 7.48 (d, 1H), 7.20 (d, 2H), 6.85 (dd, 1H), 6.25 (s, 2H), 4.38 (q, 2H), 1.24 (t, 3H)	
635		1-ethyl-2-([2-(3-fluoropyridin-2-yl)-1H-imidazol-1-yl]methyl)-6-(trifluoromethyl)-1H-benzimidazole	(CDCl ₃) 8.46 (m, 1H), 7.84(d, 1H), 7.60 (m, 2H), 7.52 (d, 1H), 7.38 (m, 1H), 7.25 (s, 1H), 7.10 (s, 1H), 6.14 (s, 2H), 4.35 (q, 2H), 1.14 (t, 3H)	
636		1-ethyl-2-([2-(3-fluoropyridin-2-yl)-1H-imidazol-1-yl]methyl)-6-methyl-1H-benzimidazole	(CDCl ₃) 8.48 (dt, 1H), 7.59-7.66 (m, 2H), 7.36 (m, 1H), 7.20 (s, 1H), 7.08-7.11 (m, 3H), 6.01 (s, 2H), 4.18 (q, 2H), 2.48 (s, 3H), 1.07 (s, 3H)	
637		6-chloro-1-ethyl-2-([2-(3-fluoropyridin-2-yl)-1H-imidazol-1-yl]methyl)-1H-benzimidazole	(CDCl ₃) 8.48 (dt, 1H), 7.74 (s, 1H), 7.61 (t, 1H), 7.37 (m, 1H), 7.20 (s, 1H), 7.23-7.26 (m, 3H), 7.08 (s, 1H), 6.01 (s, 2H), 4.25 (q, 2H), 1.07 (s, 3H)	
638		2-([2-(3-fluoropyridin-2-yl)-1H-imidazol-1-yl]methyl)-1,6-dimethyl-1H-imidazo[4,5-c]pyridine	(CDCl ₃) 8.95 (s, 1H), 8.45 (dt, J = 5.8, 1.8 Hz, 1H), 7.61 (m, 1H), 7.36 (m, 1H), 7.27 (s, 1H), 7.10 (s, 1H), 7.06 (s, 1H), 6.06 (s, 2H), 3.66 (s, 3H), 2.67 (s, 3H)	m/z 323 [M+1]
639		2-([2-(3-fluorophenyl)-1H-imidazol-1-yl]methyl)-1H-benzimidazole		m/z 293 [M+1]
640		1-benzyl-2-([2-(3-fluorophenyl)-1H-imidazol-1-yl]methyl)-1H-benzimidazole		m/z 383 [M+1]
641		2-([2-(6-fluoropyridin-2-yl)-1H-imidazol-1-yl]methyl)-1-methyl-6-(trifluoromethyl)-1H-benzimidazole	(CDCl ₃) 8.20 (m, 1H), 7.87(m, 2H), 7.63 (s, 1H), 7.48 (d, 1H), 7.20 (s, 2H), 6.84 (dd, 1H), 6.25 (s, 2H), 3.92 (s, 3H)	
642		4-(2-([2-(3-fluorophenyl)-1H-imidazol-1-yl]methyl)-1H-benzimidazol-1-yl)-2-methylbutan-2-ol		m/z 379 [M+1]

TABLE 6				
Cpd#	STRUCTURE	IUPAC NAME	NMR	MS (m/z)
643		1-ethyl-2-([2-(3-fluoropyridin-2-yl)-1H-imidazol-1-yl]methyl)-6-methyl-1H-imidazo[4,5-c]pyridine	(CDCl ₃) 8.96 (s, 1H), 8.46 (dt, 4.8, 1.4 Hz, 1H), 7.61 (m, 1H), 7.37 (m, 1H), 7.24 (d, J = 1.2 Hz, 1H), 7.10 (s, 1H), 7.09 (d, J = 1.2 Hz, 1H), 6.06 (s, 2H), 4.23 (q, J = 7.2 Hz, 2H), 2.66 (s, 3H), 1.10 (t, J = 7.2 Hz, 3H)	m/z 337 [M+1]
644		2-([2-(3-fluorophenyl)-1H-imidazol-1-yl]methyl)-1-methyl-6-(trifluoromethyl)-1H-benzimidazole		m/z 375 [M+1]
645		2-([2-(3-fluoropyridin-2-yl)-1H-imidazol-1-yl]methyl)-1-methyl-6-(trifluoromethyl)-1H-benzimidazole		m/z 376 [M+1]
646		2-([2-(3-fluorophenyl)-1H-imidazol-1-yl]methyl)-1,6-dimethyl-1H-imidazo[4,5-c]pyridine	(CDCl ₃) 8.95 (s, 1H), 7.49-7.37 (m, 3H), 7.16 (m, 2H), 7.06 (s, 1H), 6.97 (s, 1H), 5.50 (s, 2H), 3.33 (s, 3H), 2.66 (s, 3H)	m/z 322 [M+1]
647		1-ethyl-2-([2-(6-fluoropyridin-2-yl)-1H-imidazol-1-yl]methyl)-6-methyl-1H-benzimidazole	(CDCl ₃) 8.15 (dd, 1H), 7.87 (q, 1H), 7.63 (d, 1H), 7.08-7.17 (m, 4H), 6.88 (dd, 1H), 6.26 (s, 1H), 4.27 (q, 2H), 2.46 (s, 1H), 1.16 (t, 3H)	
648		1-(1-ethyl-2-([2-(3-fluoropyridin-2-yl)-1H-imidazol-1-yl]methyl)-1H-benzimidazol-5-yl)ethanone	(CDCl ₃) 8.45 (d, 1H), 8.38 (s, 1H), 7.98 (dd, 1H), 7.61 (t, 1H), 7.37 (m, 2H), 7.24 (s, 1H), 7.21 (s, 1H), 6.08 (s, 2H), 4.29 (q, 2H), 2.66 (s, 3H), 1.14 (t, 3H)	
649		1-ethyl-2-([2-(3-fluoropyridin-2-yl)-1H-imidazol-1-yl]methyl)-1H-imidazo[4,5-c]pyridine	(CDCl ₃) 9.12 (s, 1H), 8.45-8.49 (m, 2H), 7.620 (t, 1H), 7.35-7.41 (m, 1H), 7.30 (d, 1H), 7.27 (s, 1H), 7.11 (s, 1H), 6.13 (s, 2H), 4.30 (q, 2H), 1.12 (t, 3H)	
650		1-ethyl-2-([2-(3-fluoropyridin-2-yl)-1H-imidazol-1-yl]methyl)-1H-benzimidazole-5-carbonitrile	(CDCl ₃) 8.45 (d, 1H), 8.09 (s, 1H), 7.62 (t, 1H), 7.58 (d, 1H), 7.42 (s, 1H), 7.35-7.40 (m, 2H), 7.09 (s, 1H), 6.10 (s, 2H), 4.32 (q, 2H), 1.12 (t, 3H)	
651		5-chloro-1-ethyl-2-([2-(3-fluoropyridin-2-yl)-1H-imidazol-1-yl]methyl)-1H-benzimidazole		m/z 356 [M+1]

TABLE 6				
Cpd#	STRUCTURE	IUPAC NAME	NMR	MS (m/z)
652		1-ethyl-5-fluoro-2-([2-(3-fluoropyridin-2-yl)-1H-imidazol-1-yl]methyl)-1H-benzimidazole		m/z 340 [M+1]
653		5-bromo-1-ethyl-2-([2-(3-fluoropyridin-2-yl)-1H-imidazol-1-yl]methyl)-1H-benzimidazole	(CDCl3) 8.46 (s, 1H), 7.90 (s, 1H), 7.62 (t, 1H), 7.32-7.42 (m, 2H), 7.17-7.27 (m, 2H), 7.06 (s, 1H), 6.02 (s, 2H), 4.22 (q, 2H), 1.06 (t, 3H)	
654		9-ethyl-8-([2-(3-fluoropyridin-2-yl)-1H-imidazol-1-yl]methyl)-2-methyl-9H-purine	(CDCl3) 8.96 (s, 1H), 8.40 (d, 1H), 7.59 (m, 1H), 7.30 (m, 2H), 7.12 (s, 1H), 6.07 (s, 2H), 4.30 (q, 2H), 2.78 (s, 3H), 1.20 (t, 3H)	
655		3-ethyl-2-([2-(6-fluoropyridin-2-yl)-1H-imidazol-1-yl]methyl)-3H-imidazo[4,5-c]pyridine	(CDCl3) 8.84 (s, 1H), 8.47 (d, J = 5.7 Hz, 1H), 8.20 (dd, J = 7.8, 3.0 Hz, 1H), 7.91 (q, J = 8.1 Hz, 1H), 7.28 (s, 1H), 7.25 (d, J = 9.9 Hz, 2H), 6.91 (dd, J = 5.1, 3 Hz, 1H), 6.35 (s, 2H), 4.39 (q, J = 5.4 Hz, 2H), 1.49 (t, J = 5.4 Hz, 3H)	m/z 323 [M+1]
656		6-ethyl-2-([2-(6-fluoropyridin-2-yl)-1H-imidazol-1-yl]methyl)-1-methyl-1H-imidazo[4,5-c]pyridine	(CDCl3) 8.96 (s, 1H), 8.17 (br d, J = 7.6 Hz, 1H), 7.87 (q, J = 7.8 Hz, 1H), 7.21 (s, 1H), 7.19 (s, 1H), 7.12 (s, 1H), 6.87 (m, 1H), 6.24 (s, 2H), 3.83 (s, 3H), 2.95 (q, J = 7.2 Hz, 2H), 1.36 (t, J = 7.2 Hz, 3H)	m/z 337 [M+1]
657		9-ethyl-8-([2-(6-fluoropyridin-2-yl)-1H-imidazol-1-yl]methyl)-2-methyl-9H-purine	(CDCl3) 8.96 (s, 1H), 8.17 (br d, J = 7.6 Hz, 1H), 7.87 (q, J = 7.8 Hz, 1H), 7.21 (s, 1H), 7.19 (s, 1H), 7.12 (s, 1H), 6.87 (m, 1H), 6.24 (s, 2H), 3.83 (s, 3H), 2.95 (q, J = 7.2 Hz, 2H), 1.36 (t, J = 7.2 Hz, 3H)	m/z 337 [M+1]
658		1-ethyl-2-([2-(1,3-thiazol-2-yl)-1H-imidazol-1-yl]methyl)-1H-benzimidazole-5-carbonitrile	(DMSO) 8.05 (s, 1H), 7.80 (d, 1H), 7.73 (d, 1H), 7.65 (d, 1H), 7.60 (d, 1H), 7.48 (s, 1H), 7.10 (s, 1H), 6.20 (s, 2H), 4.40 (q, 2H), 1.35 (t, 3H)	
659		2-([2-(6-fluoropyridin-2-yl)-1H-imidazol-1-yl]methyl)-3-methyl-3H-imidazo[4,5-c]pyridine	(DMSO) 8.95 (s, 1H), 8.22 (d, J = 5.4 Hz, 1H), 7.99 (m, 2H), 7.54 (s, 1H), 7.43 (d, J = 5.4 Hz, 1H), 7.16 (s, 1H), 7.0 (br d, J = 8.1 Hz, 1H), 6.08 (s, 2H), 4.01 (s, 3H)	m/z 309 [M+1]
660		3-ethyl-2-([2-(3-fluoropyridin-2-yl)-1H-imidazol-1-yl]methyl)-3H-imidazo[4,5-c]pyridine	(CDCl3) 8.81 (s, 1H), 8.46 (m, 2H), 7.64 (m, 2H), 7.36 (m, 1H), 7.25 (s, 1H), 7.12 (s, 1H), 6.11 (s, 2H), 4.38 (q, J = 7.2 Hz, 2H), 1.20 (t, J = 7.2 Hz, 3H)	m/z 323 [M+1]

TABLE 6				
Cpd#	STRUCTURE	IUPAC NAME	NMR	MS (m/z)
661		1-ethyl-2-([2-(3-fluoropyridin-2-yl)-1H-imidazol-1-yl]methyl)-5-thien-3-yl-1H-benzimidazole		404.3
662		1-(1-propyl-2-([2-(6-fluoropyridin-2-yl)-1H-imidazol-1-yl]methyl)-1H-benzimidazol-5-yl)ethanone	(CDCl ₃) 8.38 (s, 1H), 8.18 (dd, 1H), 8.00 (d, 1H), 7.91 (q, 1H), 7.39 (d, 1H), 7.24 (s, 1H), 7.18 (s, 1H), 6.89 (dd, 1H), 6.31 (s, 2H), 4.28 (t, 2H), 2.67 (s, 3H), 1.69 (q, 2H), 0.88 (t, 3H)	
663		1-ethyl-2-([2-(1,3-thiazol-2-yl)-1H-imidazol-1-yl]methyl)-6-(trifluoromethyl)-1H-benzimidazole	(CD ₃ Cl ₃) 7.87 (m, 2H), 7.62 (s, 1H), 7.55 (d, 1H), 7.41 (d, 1H), 7.18 (s, 1H), 7.15 (s, 1H), 6.37 (s, 2H), 4.30 (q, 2H), 1.14 (t, 3H)	
664		1-ethyl-2-([2-(pyrimidin-2-yl)-1H-imidazol-1-yl]methyl)-6-(trifluoromethyl)-1H-benzimidazole	(CDCl ₃) 8.86 (d, 2H), 7.85 (d, 1H), 7.62 (s, 1H), 7.55 (d, 1H), 7.29 (m, 2H), 7.16 (s, 1H), 6.39 (s, 2H), 4.28 (q, 2H), 1.18 (t, 3H)	
665		2-([2-(6-(methylamino)pyridin-2-yl)-1H-imidazol-1-yl]methyl)-1-propyl-1H-benzimidazole-5-carbonitrile		m/z 372 [M+1]
666		1-ethyl-2-([2-(2-furyl)-1H-imidazol-1-yl]methyl)-6-(trifluoromethyl)-1H-benzimidazole		m/z 361 [M+1]
667		1-ethyl-2-([2-(1,3-thiazol-2-yl)-1H-imidazol-1-yl]methyl)-5-(trifluoromethyl)-1H-benzimidazole	(CDCl ₃) 8.14(s, 1H), 7.84 (d, 1H), 7.58 (d, 1H), 7.40 (m, 2H), 7.18 (s, 1H), 7.15 (s, 1H), 6.37 (s, 2H), 4.30 (q, 2H), 1.12 (t, 3H)	
668		1-ethyl-2-([2-(pyrimidin-2-yl)-1H-imidazol-1-yl]methyl)-5-(trifluoromethyl)-1H-benzimidazole	(CDCl ₃) 8.85 (d, 2H), 8.06 (s, 1H), 7.54 (d, 1H), 7.40(d, 1H), 7.28 (m, 2H), 7.15 (s, 1H), 6.38 (s, 2H), 4.27 (q, 2H), 1.15 (t, 3H)	
669		5,6-dichloro-1-ethyl-2-([2-(1,3-thiazol-2-yl)-1H-imidazol-1-yl]methyl)-1H-benzimidazole	(CDCl ₃) 7.84 (m, 2H), 7.44 (s, 1H), 7.34 (d, 1H), 7.14 (m, 2H), 6.31 (s, 2H), 4.20 (q, 2H), 1.06 (t, 3H)	

TABLE 6				
Cpd#	STRUCTURE	IUPAC NAME	NMR	MS (m/z)
670		5,6-dichloro-1-ethyl-2-[(2-pyrimidin-2-yl)-1H-imidazol-1-yl)methyl]-1H-benzimidazole	(CDCl ₃) 8.84 (d, 2H), 7.84 (s, 1H), 7.43 (s, 1H), 7.26 (m, 2H), 7.13 (s, 1H), 6.33 (s, 2H), 4.18 (q, 2H), 1.10 (t, 3H)	
671		1-ethyl-2-[(2-(1H-pyrazol-3-yl)-1H-imidazol-1-yl)methyl]-1H-benzimidazole-5-carbonitrile	(CDCl ₃) 7.97 (s, 1H), 7.53 (br s, 1H), 7.46 (d, J = 8.7 Hz, 1H), 7.35 (d, J = 8.1 Hz, 1H), 6.97 (d, J = 6.3 Hz, 2H), 6.79 (br s, 1H), 6.10 (s, 2H), 4.22 (q, J = 7.2 Hz, 2H), 0.88 (t, J = 7.2 Hz, 3H)	m/z 318 [M+1]
672		1-ethyl-6-methyl-2-[(2-(1,3-thiazol-2-yl)-1H-imidazol-1-yl)methyl]-1H-benzimidazole	(CDCl ₃) 7.86 (d, 1H), 7.65 (d, 1H), 7.39 (d, 1H), 7.14 (d, 1H), 7.11 (m, 3H), 6.31 (s, 2H), 4.16 (q, 2H), 1.02 (t, 3H)	
673		3-ethyl-2-[(2-(1,3-thiazol-2-yl)-1H-imidazol-1-yl)methyl]-5-(trifluoromethyl)-3H-imidazo[4,5-b]pyridine	(CD ₃ OD) 8.03 (d, J = 8.4 Hz, 1H), 7.72 (d, J = 3.3 Hz, 1H), 7.65 (d, J = 8.1 Hz, 1H), 7.55 (d, J = 3.3 Hz, 1H), 7.46 (d, J = 1.5 Hz, 1H), 7.20 (d, J = 1.5 Hz, 1H), 6.3 (s, 2H), 4.56 (q, J = 6.9 Hz, 2H), 1.49 (t, J = 6.9 Hz, 3H)	
674		1-ethyl-2-[(2-(1,3-thiazol-2-yl)-1H-imidazol-1-yl)methyl]-6-(trifluoromethyl)-1H-imidazo[4,5-c]pyridine	(CD ₃ OD) 8.83 (s, 1H), 8.14 (s, 1H), 7.71 (d, J = 2.7 Hz, 1H), 7.53 (d, J = 2.7 Hz, 1H), 7.45 (d, J = 0.9 Hz, 1H), 7.19 (d, J = 0.9 Hz, 1H), 6.30 (s, 2H), 4.54 (q, J = 5.4 Hz, 2H), 1.46 (t, J = 5.4 Hz, 3H)	m/z 379 [M+1]
675		3-ethyl-2-[(2-(1,3-thiazol-2-yl)-1H-imidazol-1-yl)methyl]-6-(trifluoromethyl)-3H-imidazo[4,5-c]pyridine	(CD ₃ OD) 9.02 (s, 1H), 7.89 (s, 1H), 7.70 (d, J = 3 Hz, 1H), 7.52 (d, J = 3.3 Hz, 1H), 7.46 (d, J = 1.5 Hz, 1H), 7.19 (d, J = 1.5 Hz, 1H), 6.29 (s, 2H), 4.56 (q, J = 7.5 Hz, 2H), 1.52 (t, J = 7.5 Hz, 3H)	m/z 379 [M+1]
676		1-ethyl-5-fluoro-2-[(2-(1,3-thiazol-2-yl)-1H-imidazol-1-yl)methyl]-1H-benzimidazole	(CDCl ₃) 7.88 (d, 1H), 7.45 (dd, 1H), 7.39 (d, 1H), 7.23 (dd, 1H), 7.18 (s, 1H), 7.12 (s, 1H), 7.06 (td, 1H)	
677		1,6-diethyl-2-[(2-(1,3-thiazol-2-yl)-1H-imidazol-1-yl)methyl]-1H-imidazo[4,5-c]pyridine	(CDCl ₃) 9.01 (d, J = 1.2 Hz, 1H), 7.85 (d, J = 3.6 Hz, 1H), 7.40 (d, J = 3.0 Hz, 1H), 7.16 (d, J = 1.5 Hz, 1H), 7.14 (d, J = 0.9 Hz, 1H), 7.10 (s, 1H), 6.34 (s, 2H), 4.22 (q, J = 7.2 Hz, 2H), 2.93 (q, J = 7.2 Hz, 2H), 1.35 (t, J = 7.2 Hz, 3H), 1.10 (t, J = 7.2 Hz, 3H)	m/z 339 [M+1]











TABLE 6				
Cpd#	STRUCTURE	IUPAC NAME	NMR	MS (m/z)
678		1-(2-((2-(6-fluoropyridin-2-yl)-1H-imidazol-1-yl)methyl)-1-propyl-1H-imidazo[4,5-c]pyridin-4-yl)ethanone	(CDCl ₃) 9.01 (d, J = 1.2 Hz, 1H), 7.85 (d, J = 3.6 Hz, 1H), 7.40 (d, J = 3.0 Hz, 1H), 7.16 (d, J = 1.5 Hz, 1H), 7.14 (d, J = 0.9 Hz, 1H), 7.10 (s, 1H), 6.34 (s, 2H), 4.22 (q, J = 7.2 Hz, 2H), 2.93 (q, J = 7.2 Hz, 2H), 1.35 (t, J = 7.2 Hz, 3H), 1.10 (t, J = 7.2 Hz, 3H)	m/z 339 [M+1]
679		3-ethyl-2-((2-(1,3-thiazol-2-yl)-1H-imidazol-1-yl)methyl)-3H-imidazo[4,5-c]pyridine	(CD ₃ OD) 8.87 (s, 1H), 8.30 (d, J = 5.7 Hz, 1H), 7.75 (d, J = 3.3 Hz, 1H), 7.57 (d, J = 4.8 Hz, 1H), 7.52 (d, J = 3.3 Hz, 1H), 7.39 (d, J = 1.5 Hz, 1H), 7.18 (d, J = 1.5 Hz, 1H), 6.29 (s, 2H), 4.52 (q, J = 7.5 Hz, 2H), 1.44 (t, J = 7.5 Hz, 3H)	m/z 311 [M+1]
680		1-ethyl-5-fluoro-6-methyl-2-((2-(1,3-thiazol-2-yl)-1H-imidazol-1-yl)methyl)-1H-benzimidazole	(CDCl ₃) 7.86 (d, 1H), 7.40 (m, 2H), 7.15 (d, 1H), 7.11 (d, 1H), 7.07 (d, 1H), 6.30 (s, 2H), 4.17 (q, 2H), 2.39 (d, 3H), 1.03 (t, 3H)	
681		4-(1-ethyl-2-((2-(1,3-thiazol-2-yl)-1H-imidazol-1-yl)methyl)-1H-benzimidazol-5-yl)-2-methylbutan-2-ol	(CDCl ₃) 7.85 (d, 1H), 7.60 (s, 1H), 7.38 (d, 1H), 7.22 (m, 1H), 7.15 (m, 2H), 7.09 (s, 1H), 6.31 (s, 2H), 4.17 (q, 2H), 2.80 (m, 2H), 1.81 (m, 2H), 1.22 (d, 6H), 1.01 (t, 3H)	m/z 396.3 [M+1]
682		1-(1-ethyl-2-((2-(1,3-thiazol-2-yl)-1H-imidazol-1-yl)methyl)-1H-benzimidazol-5-yl)-4-hydroxypentan-1-one	(CDCl ₃) 8.44 (d, 1H), 7.80 (dd, 1H), 7.85 (d, 1H), 7.40 (d, 1H), 7.36 (d, 1H), 7.18 (d, 1H), 7.14 (d, 1H), 6.36 (s, 2H), 4.27 (q, 2H), 3.86 (m, 1H), 3.19 (t, 2H), 1.93 (m, 2H), 1.25 (d, 3H), 1.09 (t, 3H)	m/z 410.2 [M+1]
683		1-ethyl-6-isobutyl-2-((2-(1,3-thiazol-2-yl)-1H-imidazol-1-yl)methyl)-1H-imidazo[4,5-c]pyridine	(CDCl ₃) 9.02 (s, 1H), 7.85 (d, 1H), 7.40 (d, 1H), 7.18 (s, 1H), 7.15 (s, 1H), 7.06 (s, 1H), 6.35 (s, 2H), 4.22 (q, 2H), 2.75 (d, 1H), 2.1 (m, 1H), 1.10 (t, 3H), 0.93 (d, 6H)	
684		1-ethyl-6-isopropyl-2-((2-(1,3-thiazol-2-yl)-1H-imidazol-1-yl)methyl)-1H-imidazo[4,5-c]pyridine	(CDCl ₃) 9.01 (s, 1H), 7.83 (d, J = 3.2 Hz, 1H), 7.38 (d, J = 3.2 Hz, 1H), 7.15 (s, 1H), 7.12 (s, 1H), 7.09 (s, 1H), 6.32 (s, 2H), 4.21 (q, J = 7.2 Hz, 2H), 3.17 (sept, J = 6.8 Hz, 1H), 1.34 (d, J = 6.8 Hz, 6H), 1.10 (t, J = 7.2 Hz, 3H)	m/z 353 [M+1]
685		3-ethyl-6-isobutyl-2-((2-(1,3-thiazol-2-yl)-1H-imidazol-1-yl)methyl)-3H-imidazo[4,5-c]pyridine	(CDCl ₃) 8.70 (s, 1H), 7.85 (d, 1H), 7.41 (s, 1H), 7.40 (d, 1H), 7.20 (s, 1H), 7.15 (s, 1H), 6.34 (s, 2H), 4.33 (q, 2H), 2.75 (d, 1H), 2.1 (m, 1H), 1.19 (t, 3H), 0.93 (d, 6H)	
686		1-(1-ethyl-2-((2-(1,3-thiazol-2-yl)-1H-imidazol-1-yl)methyl)-1H-benzimidazol-6-yl)ethanone	(CDCl ₃) 8.05 (d, 1H), 7.92 (dd, 1H), 7.87 (dd, 1H), 7.81 (dd, 1H), 7.40 (d, 1H), 7.20 (d, 1H), 7.14 (d, 1H), 6.17 (s, 2H), 4.33 (q, 2H), 2.67 (s, 3H), 1.13 (t, 3H)	
687		4-(1-ethyl-2-((2-(1,3-thiazol-2-yl)-1H-imidazol-1-yl)methyl)-1H-benzimidazol-6-yl)-2-methylbutan-2-ol	(CDCl ₃) 7.88 (d, 1H), 7.68 (d, 1H), 7.39 (d, 1H), 7.12-7.18 (m, 3H), 7.09 (d, 1H), 6.13 (s, 2H), 4.19 (q, 1H), 2.80-2.89 (m, 2H), 1.80-1.89 (m, 2H), 1.31 (s, 6H), 1.06 (t, 3H)	

TABLE 6				
Cpd#	STRUCTURE	IUPAC NAME	NMR	MS (m/z)
688		1-ethyl-2-[(2-(1,3-thiazol-2-yl)-1H-imidazol-1-yl)methyl]-1H-benzimidazole-6-carbonitrile	(CD3OD) (2HCl Salt) 8.38 (s, 1H), 7.98-8.01 (m, 2H), 7.95 (d, 1H), 7.87 (d, 1H), 7.71 (d, 2H), 6.50 (s, 2H), 4.62 (q, 2H), 1.62 (t, 3H)	
689		1-ethyl-2-[(2-pyrimidin-2-yl-1H-imidazol-1-yl)methyl]-1H-benzimidazole-5-carbonitrile	(CD3OD) (2HCl Salt) 8.88 (d, 2H), 8.08 (d, 1H), 7.95 (d, 1H), 7.93 (d, 1H), 7.90 (s, 1H), 7.73 (dd, 1H), 7.56 (t, 1H), 6.60 (s, 2H), 4.61 (q, 2H), 1.63 (t, 3H)	
690		1-ethyl-2-[(2-pyrimidin-2-yl-1H-imidazol-1-yl)methyl]-1H-benzimidazole-6-carbonitrile	(CDCl3) 8.85 (d, 2H), 7.82 (d, 1H), 7.69 (d, 1H), 7.55 (dd, 1H), 7.26-7.30 (m, 2H), 7.18 (d, 1H), 6.39 (s, 2H), 4.29 (q, 2H), 1.09 (t, 3H)	
691		4-chloro-1-(1-ethyl-2-[(2-(1,3-thiazol-2-yl)-1H-imidazol-1-yl)methyl]-1H-benzimidazol-5-yl)-4-methylpentan-1-one		m/z 442.2 [M-1] 444.1 [M+1]
692		1-ethyl-2-[(2-pyrimidin-2-yl-1H-imidazol-1-yl)methyl]-6-(trifluoromethyl)-1H-imidazo[4,5-c]pyridine	(CD3OD) 8.79 (s, 1 H), 8.70 (d, J = 5.1 Hz, 1H), 8.17 (s, 1H), 7.52 (d, J = 1.2 Hz, 1H), 7.31 (d, J = 1.2 Hz, 1H), 7.27 (t, J = 5.1 Hz, 1H), 6.63 (s, 1H), 6.32 (s, 2H), 4.57 (q, J = 7.5 Hz, 2H), 1.50 (t, J = 7.5 Hz, 3H)	m/z 374 [M+1]
693		2-{1-[(5-acetyl-1-ethyl-1H-benzimidazol-2-yl)methyl]-1H-imidazol-2-yl}-1,3-thiazole-4-carbonitrile	(CDCl3+CD3OD) 8.29 (s, 1H), 8.25 (d, 1H), 7.97 (dd, 1H), 7.52 (d, 1H), 7.42 (s, 1H), 7.22 (s, 1H), 6.17 (s, 2H), 4.44 (q, 2H), 2.64 (s, 3H), 1.43 (t, 3H)	m/z 377 [M+1]
694		5-bromo-1-ethyl-2-[(2-pyrimidin-2-yl-1H-imidazol-1-yl)methyl]-1H-benzimidazole	(CDCl3) 8.85 (d, 2H), 7.91 (d, 1H), 7.39 (dd, 1H), 7.27 (t, 1H), 7.25 (m, 1H), 7.19 (d, 1H), 7.13 (d, 1H), 6.35 (s, 2H), 4.20 (q, 2H), 1.08 (t, 3H)	
695		1-(3-ethyl-2-[(2-(1,3-thiazol-2-yl)-1H-imidazol-1-yl)methyl]-3H-imidazo[4,5-c]pyridin-6-yl)ethanone	(CDCl3) 8.79 (d, 2H), 8.47 (d, 1H), 7.83 (d, 1H), 7.40 (d, 1H), 7.19 (d, 1H), 7.15 (d, 1H), 6.39 (s, 2H), 4.02 (q, 2H), 2.77 (s, 3H), 1.86 (t, 3H)	m/z 353.2 [M+1]
696		3-ethyl-6-methyl-2-[(2-(1,3-thiazol-2-yl)-1H-imidazol-1-yl)methyl]-3H-imidazo[4,5-c]pyridine	(CDCl3) 8.67 (s, 1H), 7.83 (d, J = 3.0 Hz, 1H), 7.5 (s, 1H), 7.38 (d, J = 3.0 Hz, 1H), 7.18 (d, J = 1.2 Hz, 1H), 7.14 (d, J = 1.2 Hz, 1H), 6.33 (s, 2H), 4.32 (q, J = 7.5 Hz, 2H), 1.16 (t, J = 7.5 Hz, 3H)	m/z 325 [M+1]

TABLE 6				
Cpd#	STRUCTURE	IUPAC NAME	NMR	MS (m/z)
697		1-ethyl-5-pyridin-2-yl-2-[(2-pyrimidin-2-yl-1H-imidazol-1-yl)methyl]-1H-benzimidazole		m/z 382.2 [M+1]
698		1-ethyl-6-methyl-2-[(2-(1,3-thiazol-2-yl)-1H-imidazol-1-yl)methyl]-1H-imidazo[4,5-c]pyridine	(CDCl ₃) 8.96 (d, J = 0.9 Hz, 1H), 7.84 (d, J = 3.3 Hz, 1H), 7.39 (d, J = 3.0 Hz, 1H), 7.16 (d, J = 0.9 Hz, 1H), 7.13 (d, J = 1.5 Hz, 1H), 7.10 (s, 1H), 6.33 (s, 2H), 4.21 (q, J = 7.2 Hz, 2H), 2.66 (s, 2H), 1.09 (t, J = 7.2 Hz, 3H)	m/z 325 [M+1]
699		6-(cyclopentylmethyl)-1-ethyl-2-[(2-(1,3-thiazol-2-yl)-1H-imidazol-1-yl)methyl]-1H-imidazo[4,5-c]pyridine	(CDCl ₃) 9.01 (s, 1H), 7.85 (d, 1H), 7.41 (d, 1H), 7.18 (d, 1H), 7.15 (d, 1H), 7.08 (s, 1H), 6.36 (s, 2H), 4.22 (q, 2H), 2.86 (d, 2H), 2.25-2.40 (m, 1H), 1.46-1.78 (m, 6H), 1.15-1.30 (m, 2H), 1.08 (t, 3H)	
700		6-(cyclopentylmethyl)-3-ethyl-2-[(2-(1,3-thiazol-2-yl)-1H-imidazol-1-yl)methyl]-3H-imidazo[4,5-c]pyridine	(CDCl ₃) 8.68 (s, 1H), 7.84 (d, 1H), 7.48 (s, 1H), 7.41 (d, 1H), 7.20 (s, 1H), 7.15 (s, 1H), 6.14 (s, 2H), 4.32 (q, 2H), 2.88 (d, 2H), 2.25-2.38 (m, 1H), 1.45-1.72 (m, 6H), 1.16-1.33 (m, 5H)	
701		3-ethyl-6-isopropyl-2-[(2-(1,3-thiazol-2-yl)-1H-imidazol-1-yl)methyl]-3H-imidazo[4,5-c]pyridine	(CDCl ₃) 8.70 (d, J = 1.2 Hz, 1H), 7.83 (d, J = 3.0 Hz, 1H), 7.53 (d, J = 0.9 Hz, 1H), 7.39 (d, J = 3.0 Hz, 1H), 7.18 (d, J = 0.9 Hz, 1H), 7.14 (d, J = 0.9 Hz, 1H), 6.32 (s, 2H), 4.32 (q, J = 7.2 Hz, 2H), 3.18 (sept, J = 6.9 Hz, 1H), 1.34 (d, J = 6.9 Hz, 6H), 1.17 (t, J =	m/z 353 [M+1]
702		1-ethyl-2-[(2-pyrimidin-2-yl-1H-imidazol-1-yl)methyl]-1H-benzimidazole		m/z 305 [M+1]
703		6-chloro-1-ethyl-2-[(2-pyrimidin-2-yl-1H-imidazol-1-yl)methyl]-1H-benzimidazole		m/z 339 [M+1]
704		1-ethyl-6-methyl-2-[(2-pyrimidin-2-yl-1H-imidazol-1-yl)methyl]-1H-benzimidazole		m/z 319 [M+1]
705		1-ethyl-5-fluoro-2-[(2-pyrimidin-2-yl-1H-imidazol-1-yl)methyl]-1H-benzimidazole		m/z 323 [M+1]

Cpd#	STRUCTURE	IUPAC NAME	NMR	MS (m/z)
706		5-chloro-1-ethyl-2-[(2-pyrimidin-2-yl-1H-imidazol-1-yl)methyl]-1H-benzimidazole		m/z 339 [M+1]
707		3-ethyl-6-propyl-2-[(2-(1,3-thiazol-2-yl)-1H-imidazol-1-yl)methyl]-3H-imidazo[4,5-c]pyridine	(CDCl3) 8.69 (d, J = 0.9 Hz, 1H), 7.83 (d, J = 3.3 Hz, 1H), 7.49 (d, J = 1.2 Hz, 1H), 7.39 (d, J = 3.6 Hz, 1H), 7.18 (d, J = 0.9 Hz, 1H), 7.14 (d, J = 1.5 Hz, 1H), 6.33 (s, 2H), 4.31 (q, J = 7.2 Hz, 2H), 2.86 (m, 2H), 1.78 (sext, J = 7.2 Hz, 2H), 1.17 (t, J = 7.2 Hz, 3H), 0.97 (t, J = 7.2 Hz, 3H)	m/z 353 [M+1]
708		1-ethyl-6-propyl-2-[(2-(1,3-thiazol-2-yl)-1H-imidazol-1-yl)methyl]-1H-imidazo[4,5-c]pyridine	(CDCl3) 9.00 (d, J = 0.9 Hz, 1H), 7.84 (d, J = 3.0 Hz, 1H), 7.39 (d, J = 3.3 Hz, 1H), 7.16 (d, J = 1.5 Hz, 1H), 7.13 (d, J = 1.5 Hz, 1H), 7.09 (s, 1H), 6.33 (s, 2H), 4.22 (q, J = 7.2 Hz, 2H), 2.86 (m, 2H), 1.79 (sext, J = 7.2 Hz, 2H), 1.09 (t, J = 7.2 Hz, 3H), 0.96 (t, J = 7.2 Hz, 3H)	m/z 353 [M+1]
709		3,6-diethyl-2-[(2-(1,3-thiazol-2-yl)-1H-imidazol-1-yl)methyl]-3H-imidazo[4,5-c]pyridine	(CDCl3) 8.69 (d, J = 0.9 Hz, 1H), 7.84 (d, J = 3.6 Hz, 1H), 7.51 (s, 1H), 7.39 (d, J = 3.6 Hz, 1H), 7.18 (d, J = 1.5 Hz, 1H), 7.14 (d, J = 0.9 Hz, 1H), 6.33 (s, 2H), 4.32 (q, J = 7.2 Hz, 2H), 2.93 (q, J = 7.2 Hz, 2H), 1.34 (t, J = 7.2 Hz, 3H), 1.17 (t, J = 7.2 Hz, 3H)	m/z 339 [M+1]
710		1-ethyl-5-fluoro-6-methyl-2-[(2-pyrimidin-2-yl-1H-imidazol-1-yl)methyl]-1H-benzimidazole		m/z 337.4 [M+1]
711		1-ethyl-2-[(2-(2-fluoro-6-methoxyphenyl)-1H-imidazol-1-yl)methyl]-1H-imidazo[4,5-c]pyridine	(CDCl3) 9.04 (d, J = 0.9 Hz, 1H), 8.42 (d, J = 5.4 Hz, 1H), 7.46-7.38 (m, 1H), 7.25-7.22 (m, 2H), 7.00 (d, J = 0.9 Hz, 1H), 6.85-6.76 (m, 2H), 5.31, 5.22 (AB, J = 15.3 Hz, 2H), 3.85 (q, J = 7.2 Hz, 2H), 3.77 (s, 3H), 1.00 (t, J = 7.2 Hz, 3H)	m/z 352 [M+1]
712		1-ethyl-2-[(2-pyrimidin-2-yl-1H-imidazol-1-yl)methyl]-1H-imidazo[4,5-c]pyridine	(CDCl3) 9.09 (s, 1H), 8.85 (d, 2H), 8.46 (d, 1H), 7.26-7.30 (m, 3H), 7.16 (1H), 6.39 (s, 2H), 4.25 (q, 2H), 1.15 (t, 3H)	
713		1-ethyl-6-methyl-2-[(2-pyrimidin-2-yl-1H-imidazol-1-yl)methyl]-1H-imidazo[4,5-c]pyridine	(CDCl3) 8.95 (d, J = 1.2 Hz, 1H), 8.84 (d, J = 4.8 Hz, 2H), 7.27 (t, J = 5.1 Hz, 1H), 7.25 (d, J = 1.2 Hz, 1H), 7.13 (d, J = 1.2 Hz, 1H), 7.10 (s, 1H), 6.35 (s, 2H), 4.19 (q, J = 7.2 Hz, 2H), 2.65 (s, 3H), 1.12 (t, J = 7.2 Hz, 3H)	m/z 320 [M+1]










Cpd#	STRUCTURE	IUPAC NAME	NMR	MS (m/z)
714		3-ethyl-2-[(2-pyrimidin-2-yl-1H-imidazol-1-yl)methyl]-3H-imidazo[4,5-c]pyridine	(DMSO) 8.94 (s, 1H), 8.71 (d, J = 5.1 Hz, 2H), 8.22 (d, J = 5.7 Hz, 1H), 7.53 (d, J = 1.2 Hz, 1H), 7.43 (d, J = 5.4 Hz, 1H), 7.31 (t, J = 4.5 Hz, 1H), 7.17 (d, J = 1.2 Hz, 1H), 6.22 (s, 2H), 4.46 (q, J = 7.5 Hz, 2H), 1.39 (t, J = 7.5 Hz, 3H)	m/z 306 [M+1]
715		3-ethyl-2-[(2-pyrimidin-2-yl-1H-imidazol-1-yl)methyl]-3H-imidazo[4,5-c]pyridine-4-carbonitrile	(CD3OD) 8.70 (d, J = 5.1 Hz, 2H), 8.41 (d, J = 5.4 Hz, 1H), 7.73 (d, J = 5.1 Hz, 1H), 7.52 (d, J = 1.2 Hz, 1H), 7.31 (d, J = 1.2 Hz, 1H), 7.27 (t, J = 4.5 Hz, 1H), 6.35 (s, 2H), 4.75 (q, J = 7.5 Hz, 2H), 1.63 (t, J = 7.2 Hz, 3H)	m/z 331 [M+1]
716		1-ethyl-6-isopropyl-2-[(2-pyrimidin-2-yl-1H-imidazol-1-yl)methyl]-1H-imidazo[4,5-c]pyridine	(CDCl3) 9.01 (d, J = 0.9 Hz, 1H), 8.84 (d, J = 4.8 Hz, 2H), 7.28 (t, J = 4.8 Hz, 1H), 7.25 (d, J = 1.2 Hz, 1H), 7.13 (d, J = 0.9 Hz, 1H), 7.10 (d, J = 0.9 Hz, 1H), 6.35 (s, 2H), 4.21 (q, J = 7.2 Hz, 2H), 3.17 (sept, J = 6.9 Hz, 1H), 1.34 (d, J = 6.9 Hz, 6H), 1.13 (t, J = 7.2 Hz, 3H)	m/z 348 [M+1]
717		{1-ethyl-2-[(2-pyrimidin-2-yl-1H-imidazol-1-yl)methyl]-1H-benzimidazol-5-yl}(phenyl)methanone	(CDCl3) 8.84 (d, 2H), 8.20 (s, 1H), 7.80-7.90 (m, 3H), 7.40-7.60 (m, 4H), 7.3 (m, 3H), 7.18 (s, 1H), 6.38 (s, 2H), 4.25 (q, 2H), 1.22 (t, 3H)	
718		1-{1-ethyl-2-[(2-pyrimidin-2-yl-1H-imidazol-1-yl)methyl]-1H-benzimidazol-6-yl}ethanone	(CDCl3) 8.84 (d, 2H), 8.03 (s, 1H), 7.90 (d, 1H), 7.78 (d, 1H), 7.25-7.28 (m, 2H), 7.18 (s, 1H), 6.38 (s, 2H), 4.48 (q, 2H), 2.66 (s, 3H), 1.08 (t, 3H)	
719		1-ethyl-6-fluoro-2-[(2-pyrimidin-2-yl-1H-imidazol-1-yl)methyl]-1H-benzimidazole	(CDCl3) 8.87 (d, J = 5.1 Hz, 1H), 7.71 (q, J = 4.5 Hz, 1H), 7.27 (m, 3H), 7.14 (s, 1H), 7.03 (m, 2H), 6.34 (s, 2H), 4.18 (q, J = 7.5 Hz, 2H), 1.11 (t, J = 7.5 Hz, 3H)	m/z 323 [M+1]
720		1-ethyl-6-fluoro-2-[(2-(1,3-thiazol-2-yl)-1H-imidazol-1-yl)methyl]-1H-benzimidazole	(CDCl3) 7.86 (d, J = 3.3 Hz, 1H), 7.70 (q, J = 4.8 Hz, 1H), 7.40 (d, J = 3 Hz, 1H), 7.25 (s, 1H), 7.16 (d, J = 1.2 Hz, 1H), 7.01 (m, 2H), 6.32 (s, 2H), 4.19 (q, J = 6.9 Hz, 2H), 1.06 (t, J = 7.5 Hz, 3H)	m/z 328 [M+1]
721		1-ethyl-5-phenyl-2-[(2-pyrimidin-2-yl-1H-imidazol-1-yl)methyl]-1H-benzimidazole		m/z 381.5 [M+1]
722		6-(cyclopentylmethyl)-1-ethyl-2-[(2-pyrimidin-2-yl-1H-imidazol-1-yl)methyl]-1H-imidazo[4,5-c]pyridine	(CDCl3) 9.00 (s, 1H), 8.86 (d, 2H), 7.25-7.31 (m, 2H), 7.15 (s, 1H), 7.08 (s, 1H), 6.37 (s, 2H), 4.20 (q, 2H), 2.87 (d, 2H), 2.25-2.40 (m, 1H), 1.42-1.78 (m, 6H), 1.15-1.30 (m, 2H), 1.13 (t, 3H)	

TABLE 6				
Cpd#	STRUCTURE	IUPAC NAME	NMR	MS (m/z)
723		6-(cyclopentylmethyl)-3-ethyl-2-[(2-pyrimidin-2-yl-1H-imidazol-1-yl)methyl]-3H-imidazo[4,5-c]pyridine	(CDCl ₃) 8.85 (d, 2H), 8.69 (s, 1H), 7.47 (s, 1H), 7.25-7.29 (m, 2H), 7.18 (s, 1H), 6.38 (s, 2H), 4.30 (q, 2H), 2.88 (d, 2H), 2.22-2.39 (m, 1H), 1.42-1.78 (m, 6H), 1.20-1.33 (m, 5H)	
724		1-ethyl-5-isopropyl-2-[(2-pyrimidin-2-yl-1H-imidazol-1-yl)methyl]-1H-benzimidazole	(CDCl ₃) 8.85 (d, 2H), 7.63 (d, 1H), 7.24 (m, 2H), 7.21 (m, 2H), 7.13 (d, 1H), 6.33 (s, 2H), 4.14 (q, 2H), 3.02 (m, 1H), 1.29 (d, 6H), 1.08 (t, 3H)	
725		1-ethyl-6-(2-fluorophenyl)-2-[(2-pyrimidin-2-yl-1H-imidazol-1-yl)methyl]-1H-benzimidazole	(CDCl ₃) 8.87 (d, J = 4.8 Hz, 2H), 7.85 (d, J = 8.4 Hz, 1H), 7.48 (m, 3H), 7.22 (m, 6H), 6.39 (s, 2H), 4.25 (q, J = 7.2 Hz, 2H), 1.16 (t, J = 7.2 Hz, 3H)	m/z 399 [M+1]
726		1-ethyl-2-[(2-(1,3-thiazol-2-yl)-1H-imidazol-1-yl)methyl]-6-[4-(trifluoromethyl)phenyl]-1H-benzimidazole	(DMSO) 7.95 (m, 3H), 7.79 (m, 3H), 7.70 (d, J = 3.3 Hz, 1H), 7.59 (d, J = 8.7 Hz, 1H), 7.52 (s, 1H), 7.48 (s, 1H), 6.19 (s, 2H), 4.43 (q, J = 7.2 Hz, 2H), 1.34 (t, J = 7.2 Hz, 3H)	m/z 454 [M+1]
727		1-ethyl-2-[(2-pyrimidin-2-yl-1H-imidazol-1-yl)methyl]-6-[4-(trifluoromethyl)phenyl]-1H-benzimidazole	(CDCl ₃) 8.87 (d, J = 4.8 Hz, 2H), 8.00 (s, 1H), 7.86 (d, J = 8.4 Hz, 1H), 7.71 (m, 3H), 7.52 (m, 2H), 7.27 (t, J = 4.8 Hz, 1H), 7.25 (s, 1H), 7.17 (s, 1H), 6.39 (s, 2H), 4.28 (q, J = 6.9 Hz, 2H), 1.16 (t, J = 6.9 Hz, 3H)	m/z 449 [M+1]
728		6-isopropyl-3-propyl-2-[(2-(1,3-thiazol-2-yl)-1H-imidazol-1-yl)methyl]-3H-imidazo[4,5-c]pyridine	(CDCl ₃) 8.69 (d, J = 1.2 Hz, 1H), 7.83 (d, J = 3.3 Hz, 1H), 7.52 (s, 1H), 7.39 (d, 3.0 Hz, 1H), 7.21 (d, J = 1.5 Hz, 1H), 7.13 (d, J = 0.9 Hz, 1H), 6.33 (s, 2H), 4.21 (m, 2H), 3.18 (sept, J = 6.9 Hz, 1H), 1.61 (sext, J = 7.2 Hz, 2H), 1.34 (d, J = 6.9 Hz, 6H), 0.77 (t, J = 7.2 Hz, 3H)	m/z 367 [M+1]
729		6-isopropyl-1-propyl-2-[(2-(1,3-thiazol-2-yl)-1H-imidazol-1-yl)methyl]-1H-imidazo[4,5-c]pyridine	(CDCl ₃) 9.01 (d, J = 0.9 Hz, 1H), 7.85 (d, J = 3.6 Hz, 1H), 7.39 (d, J = 3.6 Hz, 1H), 7.18 (d, J = 0.9 Hz, 1H), 7.12 (d, J = 0.9 Hz, 1H), 7.07 (s, 1H), 6.33 (s, 2H), 4.10 (m, 2H), 3.16 (sept, J = 6.9 Hz, 1H), 1.52 (sext, J = 7.2 Hz, 2H), 1.34 (d, J = 6.9 Hz, 6H), 0.75 (t, J = 7.2 Hz, 3H)	m/z 367 [M+1]
730		3-ethyl-6-phenyl-2-[(2-(1,3-thiazol-2-yl)-1H-imidazol-1-yl)methyl]-3H-imidazo[4,5-c]pyridine	(CDCl ₃) 8.85 (d, 1H), 8.08 (d, 1H), 8.02 (d, 1H), 7.99 (m, 1H), 7.84 (d, 1H), 7.48 (m, 2H), 7.40 (m, 2H), 7.22 (d, 1H), 7.17 (d, 1H), 6.36 (s, 2H), 4.38 (q, 2H), 1.22 (t, 3H)	m/z 387.4 [M+1]









TABLE 6				
Cpd#	STRUCTURE	IUPAC NAME	NMR	MS (m/z)
731		3-(2-fluoroethyl)-2-((2-pyrimidin-2-yl-1H-imidazol-1-yl)methyl)-3H-imidazo[4,5-c]pyridine	(CD3OD) 8.91 (s, 1H), 8.69 (d, J = 5.1 Hz, 2H), 8.28 (d, J = 5.7 Hz, 1H), 7.52 (m, 2H), 7.30 (s, 1H), 7.26 (t, J = 4.8 Hz, 1H), 6.25 (s, 2H)	m/z 324 [M+1]
732		3-(1-ethyl-2-((2-(1,3-thiazol-2-yl)-1H-imidazol-1-yl)methyl)-1H-benzimidazol-6-yl)benzonitrile	(DMSO) 8.24 (s, 1H), 8.09 (d, J = 7.8 Hz, 1H), 8.0 (s, 1H), 7.76 (m, 2H), 7.61 (m, 5H), 7.11 (s, 1H), 6.18 (s, 2H), 4.42 (q, J = 7.5 Hz, 2H), 1.34 (t, J = 7.5 Hz, 3H)	m/z 411 [M+1]
733		3-(1-ethyl-2-((2-pyrimidin-2-yl-1H-imidazol-1-yl)methyl)-1H-benzimidazol-6-yl)benzonitrile	(DMSO) 8.76 (d, J = 5.1 Hz, 2H), 8.24 (s, 1H), 8.07 (d, J = 8.4 Hz, 1H), 7.99 (s, 1H), 7.77 (d, J = 7.8 Hz, 1H), 7.64 (t, J = 8.1 Hz, 1H), 7.50 (m, 3H), 7.34 (t, J = 4.8 Hz, 1H), 7.17 (s, 1H), 6.17 (s, 2H), 4.42 (q, J = 6.9 Hz, 2H), 1.34 (t, J = 6.9 Hz, 3H)	m/z 406 [M+1]
734		1-ethyl-2-((2-(1,3-thiazol-2-yl)-1H-imidazol-1-yl)methyl)-5-(trifluoromethoxy)-1H-benzimidazole	(CDCl3) 7.86 (d, 1H), 7.66 (s, 1H), 7.40 (d, 1H), 7.33 (d, 1H), 7.26 (d, 2H), 7.14 (s, 1H), 6.34 (s, 2H), 4.26 (q, 2H), 1.10 (t, 3H)	m/z 394 [M+1]
735		1-ethyl-2-((2-pyrimidin-2-yl-1H-imidazol-1-yl)methyl)-5-(trifluoromethoxy)-1H-benzimidazole	(CDCl3) 8.84 (d, 2H), 7.63 (s, 1H), 7.25-7.28 (m, 3H), 7.15 (d, 2H), 6.35 (s, 2H), 4.22 (q, 2H), 1.13 (t, 3H)	
736		3-ethyl-2-((2-(6-fluoropyridin-2-yl)-1H-imidazol-1-yl)methyl)-6-isopropyl-3H-imidazo[4,5-c]pyridine	(CDCl3) 8.71 (d, J = 1.2 Hz, 1H), 8.17 (dd, J = 7.7, 1.7 Hz, 1H), 7.88 (q, J = 7.9 Hz, 1H), 7.51 (d, J = 0.9 Hz, 1H), 7.21 (d, J = 0.9 Hz, 1H), 7.18 (d, J = 1.5 Hz, 1H), 6.88 (dd, J = 8.3, 2.0 Hz, 1H), 6.26 (s, 2H), 4.42 (q, J = 7.2 Hz, 2H), 3.17 (sept, J = 6.9 Hz, 1H), 1.33 (d, J = 6.9 Hz, 6H), 1.30 (t, J = 7.2 Hz, 3H)	m/z 365 [M+1]
737		1-ethyl-2-((2-(6-fluoropyridin-2-yl)-1H-imidazol-1-yl)methyl)-6-isopropyl-1H-imidazo[4,5-c]pyridine	(CDCl3) 8.99 (d, J = 0.9 Hz, 1H), 8.16 (dd, J = 7.8, 1.5 Hz, 1H), 7.88 (q, J = 7.8 Hz, 1H), 7.18 (d, J = 0.9 Hz, 1H), 7.16 (d, J = 1.5 Hz, 1H), 7.11 (d, 0.9 Hz, 1H), 6.89 (dd, J = 7.8, 2.4 Hz, 1H), 6.27 (s, 2H), 4.33 (q, J = 7.2 Hz, 2H), 3.17 (sept, J = 6.9 Hz, 1H), 1.35 (d, J = 6.9 Hz, 6H), 1.23 (t, J = 7.2 Hz, 3H)	m/z 365 [M+1]
738		3-ethyl-2-((2-(6-fluoropyridin-2-yl)-1H-imidazol-1-yl)methyl)-6-propyl-3H-imidazo[4,5-c]pyridine	(CDCl3) 8.76 (d, J = 0.9 Hz, 1H), 8.17 (dd, J = 8.0, 2.6 Hz, 1H), 7.89 (q, J = 7.8 Hz, 1H), 7.47 (s, 1H), 7.22 (d, J = 0.9 Hz, 1H), 7.18 (d, J = 1.5 Hz, 1H), 6.88 (dd, J = 8.6, 2.9 Hz, 1H), 6.27 (s, 2H), 4.42 (q, J = 7.2 Hz, 2H), 2.86 (m, 2H), 1.78 (sextet, J = 7.2 Hz, 2H), 1.20 (t, J = 7.2 Hz, 3H)	m/z 365 [M+1]

TABLE 6				
Cpd#	STRUCTURE	IUPAC NAME	NMR	MS (m/z)
739			(CD3OD) 8.60 (d, J = 4.5 Hz, 1H), 8.09 (s, 1H), 7.88 (m, 4H), 7.68 (d, J = 8.4 Hz, 1H), 7.57 (d, J = 3.3 Hz, 1H), 7.34 (m, 2H), 7.15 (s, 1H), 6.31 (s, 2H), 4.45 (q, J = 7.5 Hz, 2H), 1.32 (t, J = 7.5 Hz, 3H)	m/z 387 [M+1]
740			(CD3OD) 8.77 (m, 2H), 8.14 (s, 1H), 7.91 (m, 2H), 7.81 (d, J = 8.4 Hz, 1H), 7.62 (d, J = 8.4 Hz, 1H), 7.45 (s, 1H), 7.32 (m, 4H), 6.30 (s, 2H), 4.50 (q, J = 7.2 Hz, 2H), 1.42 (t, J = 7.2 Hz, 3H)	m/z 382 [M+1]
741			(CD3OD) 8.11 (s, 1H), 7.78-7.91 (m, 3H), 7.55-7.62 (m, 4H), 7.42 (d, J = 1.5 Hz, 1H), 6.25 (s, 2H), 4.46 (q, J = 7.5 Hz, 2H), 1.37 (t, J = 7.5 Hz, 3H)	m/z 393 [M+1]
742		1-ethyl-2-((2-(3-fluorophenyl)-1H-imidazol-1-yl)methyl)-1H-benzimidazole		321.3 [M+1]
743		2-((2-(6-fluoropyridin-2-yl)-1H-imidazol-1-yl)methyl)-1H-imidazo[4,5-b]pyridine	(HCl salt) ¹ H NMR (dmso): 1.05 (t, 3H), 2.03 (h, 2H), 4.62 (t, 2H), 6.58 (s, 2H), 7.24(d, 1H), 7.82 (t, 1H), 7.95 (s, 1H), 8.05 (m, 2H), 8.25(q, 1H), 8.62(d, 1H), 8.93 (d, 1H)	337 [M+1]
744		3-ethyl-5-fluoro-2-((2-(3-fluorophenyl)-1H-imidazol-1-yl)methyl)-3H-imidazo[4,5-b]pyridine	dihydrochloride ¹ H NMR (d6 DMSO): 8.18(1H, t), 7.99(1H, d), 7.94(1H, d), 7.72-7.69(1H, m), 7.66-7.61(1H, m), 7.58-7.50(2H, m), 7.03(1H, d), 5.96(2H, s), 4.23(2H, q), 1.31(3H, s)	340.2 [M+1]
745		3-ethyl-2-((2-(6-fluoropyridin-2-yl)-1H-imidazol-1-yl)methyl)-3H-imidazo[4,5-b]pyridine	¹ H NMR, δ ppm (CDCl ₃): 8.39(dd, 1H), 8.19(m, 1H), 8.02(dd, 1H), 7.89(m, 1H), 7.20-7.26(m, 3H), 6.89(m, 1H), 6.32(d, J = 7.97 Hz, 2H), 1.30(t, 3H)	323.3 [M+1]
746		1-ethyl-2-((2-(6-fluoropyridin-2-yl)-1H-imidazol-1-yl)methyl)-1H-imidazo[4,5-b]pyridine	(HCl salt) ¹ H NMR (dmso): 1.63(t, 3H), 4.72(q, 2H), 6.58(s, 2H), 7.30 (d, 1H), 7.82(m, 1H), 7.96(s, 1H), 8.03(m, 2H), 8.24(q, 1H), 8.60(d, 1H), 8.90(d, 1H)	323 [M+1]
747		2-((2-(6-fluoropyridin-2-yl)-1H-imidazol-1-yl)methyl)-N,N-dimethyl-1-propyl-1H-benzimidazole-5-carboxamide	CDCl ₃ : 8.17 (d, J = 7.42 Hz, 1H), 7.85-7.92 (m, 1H), 7.80 (s, 1H), 7.34-7.42 (m, 2H), 7.21 (s, 1H), 7.15 (s, 1H), 6.89 (d, J = 7.97 Hz, 1H), 6.29 (s, 2H), 4.21-4.26 (m, 2H), 3.04-3.11 (br d, 6H), 1.61-1.69 (m, 2H), 0.80(t, J = 7.42 Hz, 3H).	

TABLE 6				
Cpd#	STRUCTURE	IUPAC NAME	NMR	MS (m/z)
748		2-((2-(6-fluoropyridin-2-yl)-1H-imidazol-1-yl)methyl)-5-(5-methyl-1,2,4-oxadiazol-3-yl)-1-propyl-1H-benzimidazole	d6-DMSO: 7.94-8.09 (m, 3H), 7.82 (d, 1H), 7.76 (d, 1H), 7.47 (s, 1H), 7.18 (s, 1H), 7.03 (d, 1H), 6.15 (s, 2H), 4.35 (t, 2H), 2.65 (s, 3H), 1.74-1.82 (m, 2H), 0.94 (t, 3H).	418.3 [M+1]; 416.2 [M-1]
749		2-((2-(6-fluoropyridin-2-yl)-1H-imidazol-1-yl)methyl)-5-(5-methyl-1,3,4-oxadiazol-2-yl)-1-propyl-1H-benzimidazole	CDCl ₃ : 8.36 (s, 1H), 8.18 (dd, 1H), 8.03 (dd, 1H), 7.89 (q, 1H), 7.44 (d, 1H), 7.25 (s, 1H), 7.17 (s, 1H), 6.90 (dd, 1H), 6.33 (s, 2H), 4.29 (t, 2H), 2.59 (s, 3H), 1.59-1.74 (m, 2H), 0.82 (t, 3H).	418.3 [M+1]; 416.2 [M-1]
750		2-((2-(2,5-difluorophenyl)-1H-imidazol-1-yl)methyl)-3-ethyl-5-fluoro-3H-imidazo[4,5-b]pyridine	hydrochloride 1H NMR (d6 DMSO): 8.15(1H, t), 8.08(1H, d), 7.99(1H, d), 7.76-7.72(1H, m), 7.62-7.50(2H, m), 7.02(1H, d), 5.92(2H, s), 4.20(2H, q), 1.26(3H, t)	358.2 [M+1]
751		3-(1-((3-ethyl-5-fluoro-3H-imidazo[4,5-b]pyridin-2-yl)methyl)-1H-imidazol-2-yl)benzonitrile	hydrochloride 1H NMR (d6 DMSO): 8.30(1H, s), 8.16(1H, dd), 8.11(1H, d), 8.04(1H, d), 8.00(1H, d), 7.96(1H, d), 7.74(1H, t), 7.03(1H, d), 5.97(2H, s), 4.22(2H, q), 1.30(3H, t)	347.3 [M+1]
752		2-((2-(6-fluoropyridin-2-yl)-1H-imidazol-1-yl)methyl)-1-propyl-1H-imidazo[4,5-c]pyridine	¹ H NMR (CDCl ₃) δ 9.06 (s, 1 H), 8.42 (d, 1 H), 8.18 (d, 1 H), 7.88 (dd, 1 H), 7.28 (d, 1 H), 7.22 (d, 1 H), 7.18 (d, 1 H), 6.88 (d, 1 H), 6.29 (s, 2 H), 4.26 (t, 2 H), 1.69 (q, 2 H), 0.84 (t, 3 H)	m/e 337
753		2-((2-(6-fluoropyridin-2-yl)-1H-imidazol-1-yl)methyl)-1-isopropyl-1H-imidazo[4,5-c]pyridine	¹ H NMR (CDCl ₃) δ 9.03 (s, 1 H), 8.35 (d, 1 H), 8.14 (d, 1 H), 7.87 (dd, 1 H), 7.42 (d, 1 H), 7.15 (d, 1 H), 7.09 (d, 1 H), 6.87 (d, 1 H), 6.28 (s, 2 H), 5.08 (septet, 1 H), 1.49 (d, 6 H)	m/e 337
754		2-((2-(6-fluoropyridin-2-yl)-1H-imidazol-1-yl)methyl)-1-isobutyl-1H-imidazo[4,5-c]pyridine	¹ H NMR (CDCl ₃) δ 9.03 (s, 1 H), 8.39 (d, 1 H), 8.15 (d, 1 H), 7.85 (dd, 1 H), 7.27 (d, 1 H), 7.24 (s, 1 H), 7.17 (s, 1 H), 6.85 (d, 1 H), 6.25 (s, 2 H), 4.08 (d, 2 H), 2.09 (septet, 1 H), 0.85 (d, 6 H)	m/e 323.
755		2-((2-(6-fluoropyridin-2-yl)-1H-imidazol-1-yl)methyl)-5-(1,2,4-oxadiazol-3-yl)-1-propyl-1H-benzimidazole	Formate d6-DMSO: 9.62 (s, 1H), 7.99-8.09 (m, 3H), 7.90 (d, J = 8.24 Hz, 1H), 7.79 (d, J = 8.79 Hz, 1H), 7.52 (s, 1H), 7.17 (s, 1H), 7.02 (d, J = 7.41 Hz, 1H), 6.16 (s, 2H), 4.34-4.39 (m, 2H), 1.7-1.84 (m, 2H), 0.92 (t, J = 7.14 Hz, 3H).	
756		((2S)-1-((2-((2-(6-fluoropyridin-2-yl)-1H-imidazol-1-yl)methyl)-1-propyl-1H-benzimidazol-5-yl)carbonyl)pyrrolidin-2-yl)methanol	CDCl ₃ : 8.18 (d, J = 7.69 Hz, 1H), 7.86-7.94 (m, 2H), 7.51 (d, J = 8.52 Hz, 1H), 7.38 (d, J = 8.52 Hz, 1H), 7.22 (s, 1H), 7.17 (s, 1H), 6.9 (dd, J = 2.75. 8.24 Hz, 1H), 6.30 (s, 2H), 4.43-4.46 (m, 1H), 4.23-4.28 (m, 2H), 3.76-3.79 (m, 2H), 3.52-3.62 (m, 2H), 2.16-2.19 (m, 1H), 1.25-1.85 (m, 5H), 0.82 (t, J = 7.23 Hz, 2H).	










TABLE 6				
Cpd#	STRUCTURE	IUPAC NAME	NMR	MS (m/z)
757		3-(cyclopropylmethyl)-2-((2-(6-fluoropyridin-2-yl)-1H-imidazol-1-yl)methyl)-3H-imidazo[4,5-b]pyridine	¹ H NMR, δppm (CDCl ₃): 8.37(dd, 1H), 8.18(dd, 1H), 8.01(dd, 1H), 7.88(m, 1H), 7.20-7.26(m, 3H), 6.87(dd, 1H), 6.31(s, 2H), 4.31(d, 2H), 1.22(m, 1H), 0.42-0.53(m, 4H)	349.3 [M+1]
758		2-((2-(6-fluoropyridin-2-yl)-1H-imidazol-1-yl)methyl)-3-propyl-3H-imidazo[4,5-b]pyridine	¹ H NMR, δppm (CDCl ₃): 8.38(dd, 1H), 8.19(m, 1H), 8.02(dd, 1H), 7.89(m, 1H), 7.20-7.26(m, 3H), 6.88(m, 1H), 6.32(s, 2H), 4.35(t, 2H), 1.74(m, 2H), 0.85(t, 3H)	337.2 [M+1]
759		5-chloro-3-ethyl-2-((2-(3-fluorophenyl)-1H-imidazol-1-yl)methyl)-3H-imidazo[4,5-b]pyridine		356.2 [M+1]
760		5-chloro-2-((2-(2,5-difluorophenyl)-1H-imidazol-1-yl)methyl)-3-ethyl-3H-imidazo[4,5-b]pyridine	¹ H NMR (CDCl ₃): 7.96(1H, d), 7.38-7.34(1H, m), 7.26-7.17(4H, m), 7.04(1H, d), 5.37(1H, s), 3.96(2H, q), 1.06(3H, t)	374.2 [M+1]
761		2-((2-(6-fluoropyridin-2-yl)-1H-imidazol-1-yl)methyl)-5-(1,3,4-oxadiazol-2-yl)-1-propyl-1H-benzimidazole	CDCl ₃ : 8.45 9s, 1H), 8.43 (s, 1H), 8.19 (d, J = 7.14 Hz, 1H), 8.10 (d, J = 8.38 Hz, 1H), 7.90(q, J = 7.96 Hz, 1H), 7.48 (d, J = 8.52 Hz, 1H), 7.19 (s, 1H), 6.89-6.92 (m, 1H), 6.33 (s, 2H), 4.3 (t, J = 7.42 Hz, 2H), 1.67-1.74 9m, 2H), 0.86 (t, J = 7.42 Hz, 3H)	404.6 [M+1]; 402.3 [M-1]
762		1-ethyl-2-((2-(6-fluoropyridin-2-yl)-1H-imidazol-1-yl)methyl)-1H-imidazo[4,5-c]pyridine	¹ H NMR (CDCl ₃) δ 8.82 (s, 1 H), 8.45 (d, 1 H), 8.18 (d, 1 H), 7.88 (dd, 1 H), 7.65 (d, 1 H), 7.23 (d, 1 H), 7.20 (d, 1 H), 6.88 (d, 1 H), 6.30 (s, 2 H), 4.47 (q, 2 H), 1.33 (t, 3 H)	m/e 323
763		2-((2-(6-fluoropyridin-2-yl)-1H-imidazol-1-yl)methyl)-1-(2-methoxyethyl)-1H-imidazo[4,5-c]pyridine	¹ H NMR (CDCl ₃) δ 9.02 (s, 1 H), 8.41 (d, 1 H), 8.16 (d, 1 H), 7.84 (dd, 1 H), 7.36 (d, 1 H), 7.25 (d, 1 H), 7.21 (d, 1 H), 6.83 (d, 1 H), 6.18 (s, 2 H), 4.57 (t, 2 H), 3.56 (t, 2 H), 3.20 (s, 3 H)	m/e 353
764		1-ethyl-4-fluoro-2-((2-(3-fluorophenyl)-1H-imidazol-1-yl)methyl)-1H-imidazo[4,5-c]pyridine	¹ H NMR (CDCl ₃): 7.97(1H, dd), 7.53-7.48(1H, m), 7.44-7.38(2H, m), 7.22-7.12(3H, m), 7.04(1H, s), 5.55(2H, s), 3.75(2H, q), 0.96(3H, t)	340.1 [M+1]
765		2-((2-(6-fluoropyridin-2-yl)-1H-imidazol-1-yl)methyl)-3-isobutyl-3H-imidazo[4,5-b]pyridine	¹ H NMR, δppm (CDCl ₃): 8.38(dd, 1H), 8.18(dd, 1H), 8.01(dd, 1H), 7.88(m, 1H), 7.20-7.27(m, 3H), 6.87(dd, 1H), 6.29(s, 2H), 4.21(d, 2H), 2.27(m, 1H), 0.87(d, 6H)	351.3 [M+1]










TABLE 6				
Cpd#	STRUCTURE	IUPAC NAME	NMR	MS (m/z)
766		2-[[2-(6-fluoropyridin-2-yl)-1H-imidazol-1-yl]methyl]-3-isopropyl-3H-imidazo[4,5-b]pyridine	¹ H NMR, δppm (CDCl ₃): 8.36(dd, 1H), 8.18(m, 1H), 8.00(dd, 1H), 7.90(m, 1H), 7.18-7.22(m, 2H), 7.13(d, 1H), 6.89(m, 1H), 6.32(s, 2H), 5.04(m, 1H), 1.63(d, 6H)	337.2 [M+1]
767		5-chloro-3-ethyl-2-[[2-(6-fluoropyridin-2-yl)-1H-imidazol-1-yl]methyl]-3H-imidazo[4,5-b]pyridine	¹ H NMR (CDCl ₃): 8.17(1H, dd), 7.94-7.84(2H, m), 7.23-7.20(3H, m), 6.89-6.85(1H, m), 6.27(2H, s), 4.45(2H, q), 1.31(3H, t)	357.2 [M+1]
768		5-chloro-2-[[2-(4-chloropyridin-2-yl)-1H-imidazol-1-yl]methyl]-3-ethyl-3H-imidazo[4,5-b]pyridine	¹ H NMR (CDCl ₃): 8.41(1H, d), 8.35(1H, dd), 7.94(1H, d), 7.26-7.17(4H, m), 6.36(2H, s), 4.36(2H, q), 1.22(3H, t)	373.2 (M)
769		5-chloro-2-[[2-(6-chloropyridin-2-yl)-1H-imidazol-1-yl]methyl]-3-ethyl-3H-imidazo[4,5-b]pyridine	¹ H NMR (CDCl ₃): 8.22(1H, d), 7.94(1H, d), 7.75(1H, t), 7.29-7.20(4H, m), 6.31(2H, s), 4.45(2H, q), 1.30(3H, t)	373.3 (M)
770		2-[[2-(6-fluoropyridin-2-yl)-1H-imidazol-1-yl]methyl]-1-isobutyl-1H-imidazo[4,5-b]pyridine	(HCl salt) ¹ H-1 NMR (dmso): 1.05 (d, 6H), 2.40 (h, 1H), 4.50 (d, 2H), 6.57 (s, 2H), 7.30 (d, 1H), 7.82 (m, 1H), 7.93(s, 1H), 8.03(m, 2H), 8.23(q, 1H), 8.62(d, 1H), 8.95(d, 1H)	351 [M+1]
771		1-(cyclopropylmethyl)-2-[[2-(6-fluoropyridin-2-yl)-1H-imidazol-1-yl]methyl]-1H-imidazo[4,5-b]pyridine	(free base) ¹ H-1 NMR (CDCl ₃): 0.30 (m, 2H), 0.57(m, 2H), 1.05 (m, 1H), 4.23(d, 2H), 6.38(s, 2H), 6.80(d, 1H), 7.08(s, 1H), 7.22 (m, 1H), 7.39(s, 1H), 7.73(d, 1H), 7.90(q, 1H), 8.20(d, 1H), 8.58(d, 1H)	349 [M+1]
772		1-allyl-2-[[2-(6-fluoropyridin-2-yl)-1H-imidazol-1-yl]methyl]-1H-imidazo[4,5-b]pyridine	(HCl) salt Melting point: 225-231 (decompose) ¹ H-1 NMR (dmso): 5.22-5.53 (m, 4H), 6.21(m, 1H), 6.53 (s, 2H), 7.24(d, 1H), 7.82 (m, 1H), 7.95(s, 1H), 8.05(m, 2H), 8.25(q, 1H), 8.62(d, 1H), 8.82(d, 1H)	335 [M+1]
773		5-fluoro-2-[[2-(6-fluoropyridin-2-yl)-1H-imidazol-1-yl]methyl]-3-propyl-3H-imidazo[4,5-b]pyridine	hydrochloride ¹ H NMR (d ₆ DMSO): 8.42(1H, d), 8.26(1H, q), 8.10-8.05(2H, m), 7.96(1H, d), 7.36(1H, dd), 6.97(1H, d), 6.33(2H, s), 4.30(2H, t), 1.90-1.83(2H, m), 0.93(3H, t).	
774		1-(2-[[2-(2,5-difluorophenyl)-1H-imidazol-1-yl]methyl]-1-ethyl-1H-benzimidazol-5-yl)ethanone	¹ H NMR, δppm (CDCl ₃): 8.39(dd, 1H), 8.01(dd, 1H), 7.34-7.41(m, 2H), 7.18-7.24(m, 3H), 7.04(d, 1H), 5.40(s, 2H), 3.88(q, 2H), 2.69(s, 3H), 1.03(t, 3H)	381.4 [M+1]

TABLE 6				
Cpd#	STRUCTURE	IUPAC NAME	NMR	MS (m/z)
775		methyl 2-([2-(6-fluoropyridin-2-yl)-1H-imidazol-1-yl]methyl)-3-propyl-3H-imidazo[4,5-b]pyridine-6-carboxylate	¹ H NMR (CDCl ₃): 9.05(1H, d), 8.62(1H, d), 8.19(1H, dd), 7.88(1H, q), 7.26-7.21(2H, m), 6.89-6.85(1H, m), 6.31(2H, s), 4.39(2H, t), 3.96(3H, s), 1.79-1.72(2H, m), 0.87(3H, t)	395.3 [M+1]
776		2-([2-(6-fluoropyridin-2-yl)-1H-imidazol-1-yl]methyl)-6-(1,3,4-oxadiazol-2-yl)-3-propyl-3H-imidazo[4,5-b]pyridine	(KC 1084-78-2)	405.3 [M+1]
777		2-([2-(6-fluoropyridin-2-yl)-1H-imidazol-1-yl]methyl)-6-(1,3,4-oxadiazol-2-yl)-1-propyl-1H-benzimidazole	¹ H NMR (CDCl ₃) δ 8.46 (s, 1 H), 8.18 – 8.14 (m, 2 H), 7.95 – 7.83 (m, 3 H), 7.24 (d, 1 H), 7.17 (d, 1 H), 6.87 (d, 1 H), 6.31 (s, 2 H), 4.30 (t, 2 H), 1.71 (q, 2 H), 0.85 (t, 3 H)	m/e 404
778		2-([2-(6-fluoropyridin-2-yl)-1H-imidazol-1-yl]methyl)-6-(5-methyl-1,3,4-oxadiazol-2-yl)-1-propyl-1H-benzimidazole	¹ H NMR (CDCl ₃) δ 8.17 (d, 1 H), 8.07 (s, 1 H), 7.91 – 7.80 (m, 3 H), 7.23 (d, 1 H), 7.16 (d, 1 H), 6.87 (d, 1 H), 6.29 (s, 2 H), 4.28 (t, 2 H), 2.61 (s, 3 H), 1.69 (q, 2 H), 0.84 (t, 3 H)	m/e 418
779		2-([2-(3-fluorophenyl)-1H-imidazol-1-yl]methyl)-6-(1,2,4-oxadiazol-3-yl)-1-propyl-1H-benzimidazole	¹ H NMR (CDCl ₃) δ 8.89 (s, 1 H), 8.15 (s, 1 H), 8.12 – 8.03 (m, 1 H), 7.81 – 7.73 (m, 1 H), 7.40 – 7.02 (m, 6 H), 5.47 (s, 2 H), 3.82 (t, 2 H), 1.55 (q, 2 H), 0.79 (t, 3 H)	m/e 403
780		2-([2-(3-fluorophenyl)-1H-imidazol-1-yl]methyl)-6-(5-methyl-1,3,4-oxadiazol-2-yl)-1-propyl-1H-benzimidazole	¹ H NMR (CDCl ₃) δ 8.04 (s, 1 H), 7.94 – 7.84 (m, 2 H), 7.50 (7.39 (m, 3 H), 7.22 – 7.16 (m, 2 H), 7.08 (d, 1 H), 5.52 (s, 2 H), 3.76 (t, 2 H), 2.63 (s, 3 H), 1.47 (q, 2 H), 0.77 (t, 3 H)	m/e 417
781		2-([2-(3-fluorophenyl)-1H-imidazol-1-yl]methyl)-6-(1,3,4-oxadiazol-2-yl)-1-propyl-1H-benzimidazole	¹ H NMR (CDCl ₃) δ 8.50 (s, 1 H), 8.15 (d, 1 H), 7.98 – 7.89 (m, 2 H), 7.56 – 7.39 (m, 3 H), 7.24 – 7.05 (m, 3 H), 5.57 (s, 2 H), 3.78 (t, 2 H), 1.47 (q, 2 H), 0.80 (t, 3 H)	m/e 403
782		2-([2-(6-chloropyridin-2-yl)-1H-imidazol-1-yl]methyl)-6-(1,3,4-oxadiazol-2-yl)-3-propyl-3H-imidazo[4,5-b]pyridine	¹ H NMR (CDCl ₃): 9.14(1H, d), 8.63(1H, d), 8.51(1H, s), 8.26(1H, d), 7.76(1H, t), 7.28-7.24(3H, m), 6.36(2H, s), 4.42(2H, t), 1.81-1.74(2H, m), 0.88(3H, t)	421.3 [M+1]
783		2-([2-(6-chloropyridin-2-yl)-1H-imidazol-1-yl]methyl)-1-propyl-1H-benzimidazole-5-carbonitrile	¹ H NMR (d ₆ DMSO): 8.02 (1H, d, J = 7.7 Hz), 8.08(1H, s), 7.76(1H, t, J = 8 Hz), 7.55(1H, dd, J = 8, 1.4 Hz), 7.428 (1H, d, J = 8.5 Hz), 7.18-7.32(3H, m), 6.34(1H, s), 4.29(2H, t, J = 8.7 Hz), 1.66(2H, m), 0.81(3H, t, J = 8.7 Hz)	377.3 [M+1]

TABLE 6				
Cpd#	STRUCTURE	IUPAC NAME	NMR	MS (m/z)
784		1-propyl-2-[(2-pyridin-2-yl-1H-imidazol-1-yl)methyl]-1H-benzimidazole-5-carbonitrile		
785		2-[[2-(6-fluoropyridin-2-yl)-1H-imidazol-1-yl]methyl]-5-[[[(2S)-2-(methoxymethyl)pyrrolidin-1-yl]carbonyl]-1-propyl-1H-benzimidazole	CDCl ₃ : 8.17 (d, J = 7.42 Hz, 1H), 7.85-7.93 (m, 2H), 7.51-7.53 (m, 1H), 7.35 (d, J = 8.51 Hz, 1H), 7.21 (s, 1H), 7.16 (s, 1H), 6.89 (dd, J = 2.47, 7.97 Hz, 1H), 6.28-6.3 (m, 2H), 4.46 (bs, 1H), 4.21-4.26 (m, 2H), 3.51-3.68 (m, 4H), 3.40 (s, 3H), 1.95-2.05 (m, 4H), 1.62-1.72 (m, 2H), 0.81 (t, J = 7.42 Hz, 3H).	475.2 [M+1]; 477.2 [M-1]
786		2-[[2-(4-chloropyridin-2-yl)-1H-imidazol-1-yl]methyl]-1-propyl-1H-benzimidazole-5-carbonitrile	¹ H NMR (d ₆ DMSO): 8.42 (1H, d, J = 5 Hz), 8.35(1H, s), 8.08(1H, s), 7.1-7.65(1H, m), 6.41(1H, s), 4.19(2H, t, J = 7.4 Hz), 1.57(2H, m), 0.78(3H, t, J = 7.4 Hz)	377.3 [M+1]
787		2-[[2-(6-fluoropyridin-2-yl)-1H-imidazol-1-yl]methyl]-1-propyl-5-[[2-(pyridin-3-yl)piperidin-1-yl]carbonyl]-1H-benzimidazole	CDCl ₃ : 8.61 9s, 1H), 8.52 (d, J = 4.67 Hz, 1H), 8.15-8.18 (m, 1H), 7.87-7.93 (m, 1H), 7.85 (s, 1H), 7.66 (d, J = 7.69 Hz, 1H), 7.30-7.45 (m, 3H), 7.21 (s, 1H), 7.16 (s, 1H), 6.88-6.91 (m, 1H), 6.29 (s, 2H), 4.25 (t, J = 7.42 Hz, 2H), 2.82-2.91	524.0 [M+1]; 522.4 [M-1]
788		1-(2-[[2-(4-chloropyridin-2-yl)-1H-imidazol-1-yl]methyl]-1-ethyl-1H-benzimidazol-5-yl)ethanone	¹ H NMR, δppm (CDCl ₃): 8.36-8.47(m, 3H), 8.01(dd, 1H), 7.38(d, 1H), 7.29(dd, 1H), 7.16-7.18(m, 2H), 6.43(s, 2H), 4.30(q, 2H), 2.68(s,3H), 1.14(t, 3H)	380.3 [M+1]
789		1-(2-[[2-(6-chloropyridin-2-yl)-1H-imidazol-1-yl]methyl]-1-ethyl-1H-benzimidazol-5-yl)ethanone	¹ H NMR, δppm (CDCl ₃): 8.40(d, 1H), 8.23(d, 1H), 8.01(dd, 1H), 7.77(t, 1H), 7.41(d, 1H), 7.30(d, 1H), 7.24(d, 1H), 7.18(d, 1H), 6.36(s, 2H), 4.41(q, 2H), 2.68(s,3H), 1.25(t, 3H)	380.3 [M+1]
790		2-[[2-(3-fluorophenyl)-1H-imidazol-1-yl]methyl]-6-(5-methyl-1,2,4-oxadiazol-3-yl)-1-propyl-1H-benzimidazole	¹ H NMR (CDCl ₃) δ 8.03 (d, 1 H), 8.02 (s, 1 H), 7.85 (d, 1 H), 7.51 – 7.40 (m, 3 H), 7.21 – 7.16 (m, 2 H), 7.07 (d, 1 H), 5.52 (s, 2 H), 3.75 (t, 2 H), 2.67 (s, 3 H), 1.47 (q, 2 H), 0.76 (t, 3 H)	m/e 417
791		6-(5-ethyl-1,2,4-oxadiazol-3-yl)-2-[[2-(3-fluorophenyl)-1H-imidazol-1-yl]methyl]-1-propyl-1H-benzimidazole	¹ H NMR (CDCl ₃) δ 8.05 (d, 1 H), 8.03 (s, 1 H), 7.85 (d, 1 H), 7.52 – 7.41 (m, 3 H), 7.21 – 7.17 (m, 2 H), 7.08 (d, 1 H), 5.52 (s, 2 H), 3.76 (t, 2 H), 3.00 (q, 2 H), 1.51 – 1.45 (m, 5 H), 0.77 (t, 3 H)	m/e 431
792		2-[[2-(6-chloropyridin-2-yl)-1H-imidazol-1-yl]methyl]-1-propyl-1H-imidazo[4,5-c]pyridine	dihydrochloride ¹ H NMR (DMSO-d ₆) δ 9.32 (s, 1 H), 8.68 (d, 1 H), 8.47 (d, 1 H), 8.45 (d, 1 H), 8.09 – 8.03 (m, 2 H), 7.80 (d, 1 H), 7.54 (d, 1 H), 6.45 (s, 2 H), 4.61 (t, 2 H), 1.88 (q, 2 H), 0.95 (t, 3 H)	m/e 352

TABLE 6				
Cpd#	STRUCTURE	IUPAC NAME	NMR	MS (m/z)
793		2-[[2-(6-chloropyridin-2-yl)-1H-imidazol-1-yl]methyl]-1-ethyl-1H-benzimidazole-5-carbonitrile	Mesylate d6- DMSO: 8.02-8.14 (m, 3H), 7.97 (s, 1H), 7.86 (d, J = 8.52 Hz, 1H), 7.77 (s, 1H), 7.6 (d, J = 8.51 Hz, 1H), 7.57 (d, J = 7.97 Hz, 1H), 6.26 (s, 2H), 4.44-4.51 (m, 2H), 2.30 (s, 3H), 1.39 (t, J = 7.14 Hz, 3H)	363.2 [M+1]
794		2-[[2-(4-chloropyridin-2-yl)-1H-imidazol-1-yl]methyl]-1-propyl-1H-imidazo[4,5-c]pyridine	dihydrochloride ¹ H NMR (DMSO-d6) 8.931 (s, 1 H), 8.66 (d, 1 H), 8.45 (s, 1 H), 8.41 (d, 1 H), 8.32 (d, 1 H), 7.86 (d, 1 H), 7.63 (d, 1 H), 7.52 (d, 1H), 6.42 (s, 2 H), 4.55 (t, 2 H), 1.91 (q, 2 H), 0.97 (t, 3 H)	m/e 353
795		2-[[2-(6-fluoropyridin-2-yl)-1H-imidazol-1-yl]methyl]-6-(5-methyl-1,3,4-oxadiazol-2-yl)-3-propyl-3H-imidazo[4,5-b]pyridine	¹ H NMR (CDCl3): 9.07(1H, d), 8.56(1H, d), 8.20(1H, dd), 7.88(1H, q), 7.26-7.22(2H, m), 6.88(1H, dd), 6.32(2H, s), 4.40(2H, t), 2.65(3H, s), 1.81-1.74(2H, m), 0.88(3H, t)	419.3 [M+1]
796		2-[[2-(2-fluoropyridin-4-yl)-1H-imidazol-1-yl]methyl]-1-propyl-1H-benzimidazole-5-carbonitrile	¹ H NMR (CDCl3): 8.28(1H, d), 8.04(1H, s), 7.54-7.49(2H, m), 7.40(1H, d), 7.25-7.22(2H, m), 7.11(1H, s), 5.55(2H, s), 3.89(2H, t), 1.60-1.53(2H, m), 0.83(3H, t)	361.2 [M+1]
797		2-[[2-(6-ethoxypyridin-2-yl)-1H-imidazol-1-yl]methyl]-N,N-dimethyl-1-propyl-1H-benzimidazole-5-carboxamide	¹ H NMR (CDCl3): 8.49(1H, d), 8.10(1H, d), 7.88(1H, d), 7.71(1H, t), 7.13(1H, d), 7.00(1H, d), 6.74(1H, d), 6.45(2H, s), 4.27(2H, q), 4.08(2H, t), 3.15-3.06(6H, br d), 1.48-1.35(5H, m), 0.67(3H, t)	434.6 [M+1]
798		2-[[2-(4-chloropyridin-2-yl)-1H-imidazol-1-yl]methyl]-3-ethyl-3H-imidazo[4,5-b]pyridine	¹ H NMR, δppm (CDCl3): 8.44(dd, 1H), 8.39(m, 1H), 8.03(dd, 1H), 7.18-7.28(m, 5H), 6.41(s, 2H), 4.39(q, 2H), 1.20(t, 3H)	339.2 [M+1]
799		2-[[2-(6-chloropyridin-2-yl)-1H-imidazol-1-yl]methyl]-3-ethyl-3H-imidazo[4,5-b]pyridine	¹ H NMR, δppm (CDCl3): 8.39(dd, 1H), 8.23(dd, 1H), 8.03(dd, 1H), 7.76(t, 1H), 7.19-7.29(m, 4H), 6.36(s, 2H), 4.48(q, 2H), 1.29(t, 3H)	339.2 [M+1]
800		3-(2-fluoroethyl)-2-[[2-(3-fluorophenyl)-1H-imidazol-1-yl]methyl]-3H-imidazo[4,5-b]pyridine	¹ H NMR (CDCl3): 4.26(t, 1H), 4.35(t, 1H), 4.51(t, 1H), 4.67(t, 1H), 5.54(s, 2H), 7.09-7.45 (m, 4H), 8.06(d, 1H), 8.36(d, 1H)	340 [M+1]
801		3-cyclopropyl-2-[[2-(6-fluoropyridin-2-yl)-1H-imidazol-1-yl]methyl]-3H-imidazo[4,5-b]pyridine	¹ H NMR (CDCl3): line list provided 1.20-1.24 (m, 4H), 3.35 (m, 1H), 6.29 (s, 2H), 6.79 (d, 1H), 7.15 (m, 3H), 7.82 (q, 1H), 7.92 (d, 1H), 8.15 (d, 1H), 8.37(d, 1H)	335 [M+1]

TABLE 6				
Cpd#	STRUCTURE	IUPAC NAME	NMR	MS (m/z)
802		2-((2-(6-chloropyridin-2-yl)-1H-imidazol-1-yl)methyl)-3-(2-fluoroethyl)-3H-imidazo[4,5-b]pyridine	HCl salt ¹ H-NMR (dmso): line list provided 4.77 (s, 2H), 4.90 (m, 2H), 6.28 (s, 2H), 7.24 (m, 1H), 7.62 (d, 1H), 7.93 (m, 2H), 8.12 (m, 2H), 8.36(m, 2H)	357 [M+1]
803		1-(2-((2-(6-fluoropyridin-2-yl)-1H-imidazol-1-yl)methyl)-3-propyl-3H-imidazo[4,5-b]pyridin-6-yl)ethanone	¹ H NMR (CDCl ₃): 9.01(1H, d), 8.53(1H, d), 8.20-8.17(1H, m), 7.87(1H, q), 7.24-7.21(2H, m), 6.88-6.84(1H, m), 6.30(2H, s), 4.40(2H, t), 2.66(3H, s), 1.80-1.73(2H, m), 0.88(3H, t)	379.3 [M+1]
804		5-chloro-2-((2-(6-fluoropyridin-2-yl)-1H-imidazol-1-yl)methyl)-3-propyl-3H-imidazo[4,5-b]pyridine	hydrochloride ¹ H NMR (d6 DMSO): 8.35(1H, dd), 8.24(1H, q), 8.02(1H, s), 7.95(1H, dd), 7.90(1H, s), 7.33(1H, dd), 7.28(1H, d), 7.26(1H, d), 6.32(2H, s), 4.33(2H, t), 1.90-1.85(2H, m), 0.94(3H, t)	371.3 [M+1]
805		2-((2-(3-fluoropyridin-2-yl)-1H-imidazol-1-yl)methyl)-3-propyl-3H-imidazo[4,5-b]pyridine	dihydrochloride ¹ H NMR (d6 DMSO): 8.48-8.46(1H, m), 8.30(1H, dd), 8.12-8.03(2H, m), 7.96(1H, d), 7.90(1H, dd), 7.72-7.65(1H, m), 7.22(1H, dd), 6.18(2H, s), 4.27(2H, t), 1.83-1.75(2H, m), 0.87(3H, t)	337.2
806		2-((2-(3-fluoropyridin-2-yl)-1H-imidazol-1-yl)methyl)-1-propyl-1H-benzimidazole-5-carbonitrile	hydrochloride ¹ H NMR (d6 DMSO): 8.45(1H, m), 8.08-8.02(3H, m), 7.93(1H, s), 7.83(1H, d), 7.69-7.62(2H, m), 6.16(2H, s), 4.29(2H, t), 1.72-1.20(2H, m), 0.86(3H, t)	361.2 [M+1]
807		6-((1-((1-propyl-1H-imidazo[4,5-c]pyridin-2-yl)methyl)-1H-imidazol-2-yl)pyridine-2-carbonitrile	¹ H NMR (CDCl ₃) δ 9.03 (s, 1 H), 8.56 (d, 1 H), 8.44 (d, 1 H), 7.91 (dd, 1 H), 7.60 (d, 1 H), 7.32 (d, 1 H), 7.30 (d, 1 H), 7.25 (d, 1 H), 6.25 (s, 2 H), 4.31 (t, 2 H), 1.78 (q, 2 H), 0.90 (t, 3 H)	m/e 344
808		2-((2-(4-chloropyridin-2-yl)-1H-imidazol-1-yl)methyl)-3-propyl-3H-imidazo[4,5-b]pyridine	¹ H NMR, δppm (CDCl ₃): 8.36-8.45(m, 3H), 8.03(dd, 1H), 7.18-7.29(m, 4H), 6.40(s, 2H), 4.28(t, 2H), 1.66(m, 2H), 0.81(t, 3H)	
809		2-((2-(6-chloropyridin-2-yl)-1H-imidazol-1-yl)methyl)-3-propyl-3H-imidazo[4,5-b]pyridine	¹ H NMR, δppm (CDCl ₃): 8.39(dd, 1H), 8.23(dd, 1H), 8.02(dd, 1H), 7.76(t, 1H), 7.19-7.29(m, 4H), 6.36(s, 2H), 4.34(t, 2H), 1.72(m, 2H), 0.83(t, 3H)	
810		2-((2-(4-chloropyridin-2-yl)-1H-imidazol-1-yl)methyl)-3-cyclopropyl-3H-imidazo[4,5-b]pyridine	HCl salt ¹ H-NMR (DMSO): line list provided 1.24 (m, 4H), 3.48 (m, 1H), 6.43 (s, 2H), 7.19 (m, 1H), 7.67 (m, 1H), 7.86 (d, 1H), 7.97 (s, 1H), 8.06(s, 1H), 8.31 (d, 1H), 8.56 (d, 1H), 8.64 (m, 1H)	351 [M+1]; 349 [M-1]

TABLE 6				
Cpd#	STRUCTURE	IUPAC NAME	NMR	MS (m/z)
811		2-((2-(4-chloropyridin-2-yl)-1H-imidazol-1-yl)methyl)-3-(2-fluoroethyl)-3H-imidazo[4,5-b]pyridine	¹ H NMR (CDCl ₃): 4.60-4.77 (m, 2H), 4.81 (s, 2H), 6.35(s, 2H), 7.20-7.25 (m, 5H), 8.03 (d, 1H), 8.38 (m, 1H)	357 [M+1]
812		6-((1-((3-propyl-3H-imidazo[4,5-b]pyridin-2-yl)methyl)-1H-imidazol-2-yl)pyridine-2-carbonitrile	¹ H NMR, dppm (CDCl ₃): 8.56(dd, 1H), 8.39(dd, 1H), 8.00(dd, 1H), 7.92(dd, 1H), 7.63(dd, 1H), 7.20-7.29(m, 3H), 6.32(s, 2H), 4.36(t, 2H), 1.80(m, 2H), 0.89(t, 3H) LC-MS: calcd: 343.39, found 344.1[M+1]	
813		6-((1-((3-propyl-3H-imidazo[4,5-b]pyridin-2-yl)methyl)-1H-imidazol-2-yl)pyridine-2-carboxamide	¹ H NMR, δppm (CDCl ₃): 8.88(s, 1H), 8.42(dd, 1H), 8.38(dd, 1H), 8.15(dd, 1H), 7.98(dd, 1H), 7.93(d, 1H), 7.22-7.27(m, 3H), 6.18 (s, 2H), 5.67(s, 1H), 4.07(t, 2H), 1.54(m, 2H), 0.83(t, 3H)	362.2 [M+1]
814		3-propyl-2-((2-(6-(trifluoromethyl)pyridin-2-yl)-1H-imidazol-1-yl)methyl)-3H-imidazo[4,5-b]pyridine	¹ H NMR, δppm (CDCl ₃): 8.53(d, 1H), 8.39(dd, 1H), 7.96-8.06(m, 2H), 7.64(dd, 1H), 7.20-7.27(m, 3H), 6.42(s, 2H), 4.26(t, 2H), 1.61(m, 2H), 0.72(t, 3H)	387.3 [M+1]
815		3-(2-fluoroethyl)-2-((2-(3-fluoropyridin-2-yl)-1H-imidazol-1-yl)methyl)-3H-imidazo[4,5-b]pyridine	¹ H NMR (CDCl ₃): line list provided 4.58-4.66 (m, 2H), 4.75 (s, 2H), 6.03 (s, 2H), 7.21-7.35 (m, 4H), 7.57 (m, 1H), 8.04 (d, 1H), 8.41(m, 1H)	341 [M+1]
816		2-((2-(3-fluorophenyl)-1H-imidazol-1-yl)methyl)-1-phenyl-1H-benzimidazole	¹ H NMR (CDCl ₃): 5.38(s, 2H), 6.91-7.17(m, 8H), 7.22-7.45 (m, 6H), 7.84 (d, 1H)	369
817		2-((2-(6-fluoropyridin-2-yl)-1H-imidazol-1-yl)methyl)-1-phenyl-1H-benzimidazole	¹ H NMR: 6.22 (s, 2H), 6.78 (m, 1H), 7.03-7.37 (m, 9H), 7.65-7.92 (m, 3H)	368 [M-1]
818		3-allyl-2-((2-(3-fluorophenyl)-1H-imidazol-1-yl)methyl)-3H-imidazo[4,5-b]pyridine	¹ H NMR (CDCl ₃): 4.59-4.62 (m, 3H), 5.03 (d, 1H), 5.43(s, 2H), 5.70 (m, 1H), 7.03 (s, 1H), 7.17 (m, 1H), 7.25 (m, 2H), 7.43(m, 3H), 8.04 (d, 1H), 8.40 (d, 1H)	334 [M+1]; 332 [M-1]
819		2-((2-(6-fluoropyridin-2-yl)-1H-imidazol-1-yl)methyl)-1-propyl-5-((4-pyridin-2-yl)piperazin-1-yl)carbonyl-1H-benzimidazole	CDCl ₃ : 8.15-8.2 (m, 2H), 7.82-7.94 (m, 2H), 7.44-7.53 (m, 1H), 7.35-7.44 (m, 2H), 7.18 (q, 2H), 6.88 (d, 1H), 6.71 (s, 1H), 6.65 (s, 1H), 6.29 (s, 2H), 4.21-4.29 (m, 2H), 3.53-4.0 (m, 8H), 1.53-1.71 (m, 2H), 0.82 (t, 3H)	

Cpd#	STRUCTURE	IUPAC NAME	NMR	MS (m/z)
820		5-[(2-[(2-(6-fluoropyridin-2-yl)-1H-imidazol-1-yl)methyl]-1-propyl-1H-benzimidazol-5-yl)carbonyl]-2-oxa-5-azabicyclo[2.2.1]heptane	CDCl ₃ : 8.18 (d, 7.97 Hz, 1H), 7.85 (m, 2H), 7.52-7.54 (m, 1H), 7.38-7.42 (m, 1H), 7.17-7.26 (m, 2H), 6.90 (d, J = 7.96 Hz, 1H), 6.30 (s, 2H), 4.73-5.06 (m, 1H), 4.56 (d, 1H), 4.24-4.35 (m, 2H), 4.04-4.10 (m, 1H), 3.81-3.89 (m, 1H), 3.51-3.7 (m, 2H), 1.96 (s, 1H), 1.86 (s, 1H), 1.64-1.71 (m, 2H), 0.83 (t, J = 7.28 Hz, 3H).	
821		2-[(2-(2-fluorophenyl)-1H-imidazol-1-yl)methyl]-3-propyl-3H-imidazo[4,5-b]pyridine	¹ H NMR, δppm (CDCl ₃): 8.39(dd, 1H), 8.03(dd, 1H), 7.65(m, 1H), 7.50(m, 1H), 7.20-7.35(m, 3H), 7.07(d, 1H), 5.38 (s, 2H), 3.84(t, 2H), 1.46(m, 2H), 0.74(t, 3H)	336.2 [M+1]
822		2-[(2-(2-chlorophenyl)-1H-imidazol-1-yl)methyl]-3-propyl-3H-imidazo[4,5-b]pyridine	¹ H NMR, δppm (CDCl ₃): 8.38(dd, 1H), 8.02(dd, 1H), 7.33-7.56(m, 4H), 7.19-7.27(m, 2H), 7.09(d, 1H), 5.30 (s, 2H), 3.85(t, 2H), 1.50(m, 2H), 0.76(t, 3H)	352.2 [M+1]
823		3-propyl-2-[(2-(2-(trifluoromethyl)phenyl)-1H-imidazol-1-yl)methyl]-3H-imidazo[4,5-b]pyridine	¹ H NMR, δppm (CDCl ₃): 8.39(dd, 1H), 8.02(dd, 1H), 7.84 (dd, 1H), 7.53-7.65(m, 2H), 7.45(m, 1H), 7.21-7.27(m, 2H), 7.09(d, 1H), 5.17 (s, 2H), 3.93(t, 2H), 1.59(m, 2H), 0.80(t, 3H)	386.2 [M+1]
824		3-ethyl-2-[(2-(3-fluoropyridin-2-yl)-1H-imidazol-1-yl)methyl]-3H-imidazo[4,5-b]pyridine	¹ H NMR, δppm (CDCl ₃): 8.47(m, 1H), 8.40(dd, 1H), 8.04(dd, 1H), 7.62(m, 1H), 7.36(m, 1H), 7.23-7.28(m, 2H), 7.16(d, 1H), 6.10(s, 2H), 4.37(q, 2H), 1.19(t, 3H)	323.3 [M+1]
825		3-ethyl-2-[(2-(3-fluoropyridin-2-yl)-1H-imidazol-1-yl)methyl]-6-(5-methyl-1,3,4-oxadiazol-2-yl)-3H-imidazo[4,5-b]pyridine	¹ H NMR (CDCl ₃): 9.08(1H, d), 8.58(1H, d), 8.45-8.43(1H, m), 7.65-7.59(1H, m), 7.39-7.30(2H, m), 7.18(1H, s), 6.14(2H, s), 4.43(2H, q), 2.66(3H, s), 1.23(3H, t)	405.3 [M+1]
826		1-allyl-2-[(2-(6-fluoropyridin-2-yl)-1H-imidazol-1-yl)methyl]-1H-imidazo[4,5-c]pyridine	¹ H NMR (CDCl ₃) δ 9.05 (s, 1 H), 8.40 (d, 1 H), 8.14 (d, 1 H), 7.85 (dd, 1 H), 7.26 (d, 1 H), 7.22 (d, 1 H), 7.17 (6.85 (d, 1 H), 6.19 (s, 2 H), 5.82 – 5.70 (m, 1 H), 5.12 (d, 1 H), 4.99 – 4.96 (m, 2 H), 4.82 (d, 1 H)	m/e 335
827		2-[(2-(6-fluoropyridin-2-yl)-1H-imidazol-1-yl)methyl]-1-prop-2-ynyl-1H-imidazo[4,5-c]pyridine	¹ H NMR (CDCl ₃) δ 9.04 (s, 1 H), 8.46 (d, 1 H), 8.15 (d, 1 H), 7.86 (dd, 1 H), 7.41 (d, 1 H), 7.26 (d, 1 H), 7.21 (d, 1 H), 6.85 (d, 1 H), 6.23 (s, 2 H), 5.21 (d, 2 H), 2.33 – 2.32 (m, 1 H)	m/e 333
828		2-[(2-(3-fluorophenyl)-1H-imidazol-1-yl)methyl]-3-prop-2-ynyl-3H-imidazo[4,5-b]pyridine	¹ H NMR (dmsO): 3.47 (s, 1H), 4.78 (s, 2H), 5.65 (s, 2H), 7.07-7.45 (m, 5H), 7.45(m, 2H), 8.07(m, 1H), 8.42(m, 1H)	332 [M+1]; 330 [M-1]

TABLE 6				
Cpd#	STRUCTURE	IUPAC NAME	NMR	MS (m/z)
829		2-((2-(3-fluoropyridin-2-yl)-1H-imidazol-1-yl)methyl)-3-prop-2-ynyl-3H-imidazo[4,5-b]pyridine	HCl salt H-1 NMR (dmso): line list provided 3.51 (s, 1H), 5.25 (s, 2H), 6.25(s, 2H), 7.27(m, 1H), 7.72(m, 1H), 7.94-8.15 (m, 4H), 8.35(m, 1H), 8.50(1H)	333 [M+1]
830		3-cyclopropyl-2-((2-(3-fluoropyridin-2-yl)-1H-imidazol-1-yl)methyl)-3H-imidazo[4,5-b]pyridine	HCl salt H-1 NMR (dmso): line list provided 1.19 (m, 4H), 3.36 (m, 2H), 6.24 (s, 2H), 7.21 (m, 1H), 7.70 (m, 1H), 7.88 (d, 2H), 8.00 (s, 1H), 8.07 (m, 3H), 8.31 (d, 1H), 8.52 (d, 1H)	335 [M+1]
831		3-cyclopropyl-2-((2-(3-fluorophenyl)-1H-imidazol-1-yl)methyl)-3H-imidazo[4,5-b]pyridine	HCl salt H-1 NMR (DMSO): line list provided 1.03-1.22 (m, 4H), 3.37 (m, 1H), 5.96 (s, 2H), 7.22 (m, 1H), 7.45-7.78 (m, 4H), 7.95-8.03 (m, 3H), 8.37 (d, 1H)	334 [M+1]; 332 [M-1]
832		3-ethyl-2-((2-(6-(trifluoromethyl)pyridin-2-yl)-1H-imidazol-1-yl)methyl)-3H-imidazo[4,5-b]pyridine	¹ H NMR, δppm (CDCl ₃): 8.52(d, 1H), 8.39(dd, 1H), 7.90-8.05(m, 2H), 7.64(d, 1H), 7.21-7.27(m, 3H), 6.42(s, 2H), 4.39(q, 2H), 1.17(t, 3H)	373.3 [M+1]
833		1-[1-ethyl-2-((2-(6-(trifluoromethyl)pyridin-2-yl)-1H-imidazol-1-yl)methyl)-1H-benzimidazol-5-yl]ethanone	¹ H NMR, δppm (CDCl ₃): 8.52(d, 1H), 8.40(d, 1H), 7.99-8.03(m, 2H), 7.64(d, 1H), 7.40(d, 1H), 7.20-7.22(m, 2H), 6.41(s, 2H), 4.34(q, 2H), 2.68(s, 3H), 1.15(t, 3H)	414.2 [M+1]
834		3-(2-fluoroethyl)-2-((2-(6-(trifluoromethyl)pyridin-2-yl)-1H-imidazol-1-yl)methyl)-3H-imidazo[4,5-b]pyridine	H-1 NMR (CDCl ₃): line list provided 4.60-4.81 (m, 4H), 6.30 (s, 2H), 7.18-7.30 (m, 3H), 7.54 (d, 1H), 7.88-8.00 (m, 2H), 8.33(d, 1H), 8.49(d, 1H)	391 [M+1]; 389 [M-1]
835		3-allyl-2-((2-(6-fluoropyridin-2-yl)-1H-imidazol-1-yl)methyl)-3H-imidazo[4,5-b]pyridine	H-1 NMR (CDCl ₃): line list provided 4.87 (d, 1H), 5.14 (m, 3H), 5.87 (m, 1H), 6.24 (s, 2H), 6.86 (dd, 1H), 7.23 (m, 3H), 7.84 (m, 1H), 8.02 (d, 1H), 8.15 (m, 1H), 8.38 (d, 1H)	335 [M+1]; 333 [M-1]
836		6-{1-[(3-ethyl-3H-imidazo[4,5-b]pyridin-2-yl)methyl]-1H-imidazol-2-yl}pyridine-2-carbonitrile	¹ H NMR, δppm (CDCl ₃): 8.56(dd, 1H), 8.40(dd, 1H), 8.01(dd, 1H), 7.93(t, 1H), 7.63(dd, 1H), 7.21-7.29(m, 3H), 6.33(s, 2H), 4.44(q, 2H), 1.35(t, 3H)	330.1 [M+1]
837		2-((2-(6-fluoropyridin-2-yl)-1H-imidazol-1-yl)methyl)-N,N-dimethyl-1-propyl-1H-benzimidazole-6-carboxamide	¹ H NMR (CDCl ₃) δ 8.17 (d, 1 H), 7.89 (dd, 1 H), 7.74 (d, 1 H), 7.49 (s, 1 H), 7.30 (d, 1 H), 7.21 (d, 1 H), 7.15 (d, 1 H), 6.89 (d, 1 H), 6.29 (s, 2 H), 4.23 (t, 2 H), 3.07 (br d, 6 H), 1.66 (q, 2 H), 0.81 (t, 3 H)	m/e 407

TABLE 6				
Cpd#	STRUCTURE	IUPAC NAME	NMR	MS (m/z)
838		1-propyl-2-((2-[6-(trifluoromethyl)pyridin-2-yl]-1H-imidazol-1-yl)methyl)-1H-imidazo[4,5-c]pyridine	¹ H NMR (CDCl ₃) δ 9.08 (s, 1 H), 8.52 (d, 1 H), 8.44 (d, 1 H), 7.99 (dd, 1 H), 7.65 (d, 1 H), 7.29 (d, 1 H), 7.26 (d, 1 H), 7.21 (d, 1 H), 6.40 (s, 2 H), 4.19 (t, 2 H), 1.54 (q, 2 H), 0.71 (t, 3 H)	m/e 387
839		2-((2-[6-fluoropyridin-2-yl]-1H-imidazol-1-yl)methyl)-6-(morpholin-4-ylcarbonyl)-1-propyl-1H-benzimidazole	¹ H NMR (CDCl ₃) δ 8.17 (d, 1 H), 7.88 (dd, 1 H), 7.75 (d, 1 H), 7.51 (s, 1 H), 7.28 – 7.25 (m, 1 H), 7.21 (s, 1 H), 7.16 (s, 1 H), 6.89 (d, 1 H), 6.29 (s, 2 H), 4.24 (t, 2 H), 3.59 (br s, 8 H), 1.67 (q, 2 H), 0.82 (t, 3 H)	m/e 449
840		2-((2-[6-fluoropyridin-2-yl]-1H-imidazol-1-yl)methyl)-1-propyl-6-((4-pyridin-2-yl)piperazin-1-yl)carbonyl-1H-benzimidazole	¹ H NMR (CDCl ₃) δ 8.19 – 8.16 (m, 2 H), 7.88 (dd, 1 H), 7.77 (d, 1 H), 7.53 (s, 1 H), 7.48 (d, 1 H), 7.32 (d, 1 H), 7.26 (d, 1 H), 7.17 (d, 1 H), 6.89 (d, 1 H), 6.69 – 6.65 (m, 2 H), 6.30 (s, 2 H), 4.25 (t, 2 H), 3.59 (br s, 8 H), 1.67 (q, 2 H), 0.82 (t, 3 H)	m/e 525
841		methyl 2-((2-[6-fluoropyridin-2-yl]-1H-imidazol-1-yl)methyl)-1-propyl-1H-benzimidazole-6-carboxylate	¹ H NMR (CDCl ₃) δ 8.18 (d, 1 H), 8.09 (s, 1 H), 7.98 – 7.87 (m, 2 H), 7.76 (d, 1 H), 7.24 (s, 1 H), 7.17 (s, 1 H), 6.88 (d, 1 H), 6.31 (s, 2 H), 4.29 (t, 2 H), 3.94 (s, 3 H), 1.69 (q, 2 H), 0.85 (t, 3 H)	m/e 39
842		1-(2-fluoroethyl)-2-((2-[3-fluorophenyl]-1H-imidazol-1-yl)methyl)-1H-imidazo[4,5-c]pyridine	¹ H NMR (CDCl ₃): 4.02 (t, 1H), 4.05 (t, 1H), 4.32 (t, 1H), 4.47 (t, 1H), 5.58 (s, 2H), 7.05-7.33 (m, 5H), 7.44 (m, 2H), 8.47 (d, 1H), 9.10 (s, 1H)	340 [M+1]
843		2-((2-[4-chloro-1-methyl-1H-pyrazol-3-yl]-1H-imidazol-1-yl)methyl)-3-(2-fluoroethyl)-3H-imidazo[4,5-b]pyridine	¹ H NMR (CDCl ₃): line list provided 3.89 (s, 3H), 4.55 (s, 2H), 4.69 (m, 1H), 4.71 (m, 1H), 5.93 (s, 2H), 7.11 (s, 1H), 7.25 (m, 2H), 7.47 (s, 1H), 8.04 (d, 1H), 8.36 (d, 1H)	360 [M+1]
844		(4S)-5-((2-((2-[6-fluoropyridin-2-yl]-1H-imidazol-1-yl)methyl)-1-propyl-1H-benzimidazol-6-yl)carbonyl)-2-oxa-5-azabicyclo[2.2.1]heptane	¹ H NMR (CDCl ₃) δ 8.18 (d, 1 H), 7.89 (dd, 1 H), 7.78-7.72 (m, 1 H), 7.62 (s, 1 H), 7.42-7.33 (m, 1 H), 7.22 (s, 1 H), 7.17 (s, 1 H), 6.89 (d, 1 H), 6.30 (s, 2 H), 5.06-4.51 (m, 2 H), 4.25 (t, 2 H), 4.11-3.48 (m, 4 H), 1.96-1.87 (m, 2 H), 1.68 (q, 2 H), 0.83 (t, 3 H)	m/e 461
845		1-propyl-2-((2-pyrazin-2-yl)-1H-imidazol-1-yl)methyl)-1H-imidazo[4,5-c]pyridine	¹ H NMR (CDCl ₃) δ 9.58 (s, 1 H), 9.09 (d, 1 H), 8.61 – 8.42 (m, 3 H), 7.32 – 7.24 (m, 3 H), 6.33 (s, 2 H), 4.16 (t, 2 H), 1.63 (q, 2 H), 0.81 (t, 3 H)	m/e 320
846		3-ethyl-2-((2-(2-methyl-1,3-thiazol-4-yl)-1H-imidazol-1-yl)methyl)-3H-imidazo[4,5-b]pyridine	¹ H NMR, δppm (CDCl ₃): 8.40(dd, 1H), 8.05(dd, 1H), 7.88(s, 1H), 7.25(m, 1H), 7.11(d, 1H), 7.07(d, 1H), 6.29(s, 2H), 4.35(q, 2H), 2.78(s, 3H), 1.18(t, 3H)	325.2 [M+1]










TABLE 6				
Cpd#	STRUCTURE	IUPAC NAME	NMR	MS (m/z)
847		1-(1-ethyl-2-((2-(2-methyl-1,3-thiazol-4-yl)-1H-imidazol-1-yl)methyl)-1H-benzimidazol-5-yl)ethanone	¹ H NMR, δppm (CDCl ₃): 8.41(d, 1H), 8.01(dd, 1H), 7.89(s, 1H), 7.38(d, 1H), 7.10(d, 1H), 7.05(d, 1H), 6.30(s, 2H), 4.27(q, 2H), 2.79(s, 3H), 2.69(s, 3H), 1.12(t, 3H)	366.2 [M+1]
848		2-((2-(3,5-difluorophenyl)-1H-imidazol-1-yl)methyl)-3-propyl-3H-imidazo[4,5-b]pyridine	¹ H NMR, δppm (CDCl ₃): 8.41(dd, 1H), 8.06(dd, 1H), 7.19-7.29(m, 4H), 7.08(d, 1H), 6.93(m, 1H), 5.52(s, 2H), 3.96(t, 2H), 1.57(m, 3H), 0.82(t, 3H)	354.3 [M+1]
849		2-((2-(3,5-difluorophenyl)-1H-imidazol-1-yl)methyl)-3-ethyl-3H-imidazo[4,5-b]pyridine	¹ H NMR, δppm (CDCl ₃): 8.41(dd, 1H), 8.06(dd, 1H), 7.19-7.31(m, 4H), 7.07(d, 1H), 6.93(m, 1H), 5.52(s, 2H), 4.05(q, 2H), 1.14(t, 3H)	340.3 [M+1]
850		3-propyl-2-((2-(1,3-thiazol-2-yl)-1H-imidazol-1-yl)methyl)-3H-imidazo[4,5-b]pyridine	¹ H NMR, δppm (CDCl ₃): 8.40(dd, 1H), 8.05(dd, 1H), 7.86(d, 1H), 7.40(d, 1H), 7.25(d, 1H), 7.22(d, 1H), 7.15(d, 1H), 6.37(s, 2H), 4.27(t, 2H), 1.62(m, 2H), 0.78(t, 3H)	325.3 [M+1]
851		3-propyl-2-((2-(2,4,5-trifluorophenyl)-1H-imidazol-1-yl)methyl)-3H-imidazo[4,5-b]pyridine	¹ H NMR, δppm (CDCl ₃): 8.41(dd, 1H), 8.03(dd, 1H), 7.50(m, 1H), 7.22-7.29(m, 2H), 7.07-7.15(m, 2H), 5.37(s, 2H), 3.93(t, 2H), 1.53(m, 2H), 0.82(t, 3H)	372.2 [M+1]
852		2-((2-(6-fluoropyridin-3-yl)-1H-imidazol-1-yl)methyl)-3-propyl-3H-imidazo[4,5-b]pyridine	¹ H NMR (CDCl ₃): 8.53-8.51(1H, m), 8.40(1H, dd), 8.16-8.14(1H, m), 8.03(1H, dd), 7.28-7.23(2H, m), 7.11(1H, d), 7.05(1H, dd), 5.46(2H, s), 4.00(2H, t), 1.65-1.59(2H, m), 0.84(3H, t)	337.1 [M+1]
853		2-((2-(1,5-dimethyl-1H-pyrazol-3-yl)-1H-imidazol-1-yl)methyl)-3-propyl-3H-imidazo[4,5-b]pyridine	¹ H NMR (CDCl ₃): 8.37(1H, dd), 8.04(1H, dd), 7.22(1H, dd), 7.06(1H, d), 7.02(1H, d), 6.66(1H, s), 6.20(1H, s), 4.19(2H, t), 3.83(3H, s), 2.33(3H, s), 1.60-1.53(2H, m), 0.76(3H, t)	336.2 [M+1]
854		2-((2-(3,4-difluorophenyl)-1H-imidazol-1-yl)methyl)-3-propyl-3H-imidazo[4,5-b]pyridine	¹ H NMR (CDCl ₃): 8.38(1H, dd), 8.04(1H, dd), 7.56-7.50(1H, m), 7.41-7.36(1H, m), 7.30-7.21(2H, m), 7.15(1H, d), 7.04(1H, d), 5.4592H, s), 3.92(2H, t), 1.60-1.50(2H, m), 0.80(3H, t)	354.3 [M+1]
855		3-(2-fluoroethyl)-2-((2-(3-fluoro-2-methylphenyl)-1H-imidazol-1-yl)methyl)-3H-imidazo[4,5-b]pyridine	¹ H NMR (CDCl ₃): 2.13 (s, 3H), 4.14 (m, 1H), 4.24 (m, 1H), 4.48 (m, 1H), 4.64 (m, 1H), 5.29 (s, 2H), 7.09 7.26 (m, 6H), 8.02 (d, 1H), 8.35 (d, 1H)	354 [M+1]

TABLE 6				
Cpd#	STRUCTURE	IUPAC NAME	NMR	MS (m/z)
856		2-((2-(2,6-difluorophenyl)-1H-imidazol-1-yl)methyl)-3-(2-fluoroethyl)-3H-imidazo[4,5-b]pyridine	¹ H NMR (CDCl ₃): line list provided 4.14 (t, 1H), 4.24 (t, 1H), 4.46 (t, 1H), 4.64 (t, 1H), 5.40 (s, 2H), 6.98 (m, 2H), 7.11 (s, 1H), 7.26 (m, 3H), 8.01 (d, 1H), 8.34 (d, 1H)	358 [M+1]
857		3-(2-fluoroethyl)-2-((2-(3-fluoro-4-methylphenyl)-1H-imidazol-1-yl)methyl)-3H-imidazo[4,5-b]pyridine	¹ H NMR (CDCl ₃): line list provided 2.35 (s, 3H), 4.26 (t, 1H), 4.35 (t, 1H), 4.51 (t, 1H), 4.67 (t, 1H), 5.53 (s, 2H), 7.07 (s, 1H), 7.18 (s, 1H), 7.26-7.34 (m, 4H), 8.07 (d, 1H), 8.36 (d, 1H)	354 [M+1]
858		3-ethyl-2-((2-(3-fluorophenyl)-1H-imidazol-1-yl)methyl)-6-(5-methyl-1,3,4-oxadiazol-2-yl)-3H-imidazo[4,5-b]pyridine	dihydrochloride ¹ H NMR (d ₆ -DMSO): 9.96(1H, d), 8.50(1H, d), 7.98-7.91(2H, m), 7.72-7.50(4H, m), 6.01(2H, s), 4.35(2H, q), 2.58(3H, s), 1.35(3H, t)	404.3 [M+1]
859		2-((2-(2-methyl-1,3-thiazol-4-yl)-1H-imidazol-1-yl)methyl)-3-propyl-3H-imidazo[4,5-b]pyridine	¹ H NMR, δppm (CDCl ₃): 8.40(dd, 1H), 8.05(dd, 1H), 7.91(s, 1H), 7.24(m, 1H), 7.11(d, 1H), 7.07(d, 1H), 6.30(s, 2H), 4.22(t, 2H), 2.78(s, 3H), 1.62(m, 2H), 0.79(t, 3H)	339.2 [M+1]
860		2-((2-(2-methoxypyridin-3-yl)-1H-imidazol-1-yl)methyl)-3-propyl-3H-imidazo[4,5-b]pyridine	¹ H NMR, δppm (CDCl ₃): 8.38(dd, 1H), 8.30(dd, 1H), 8.03(dd, 1H), 7.87(dd, 1H), 7.19-7.27(m, 2H), 7.01-7.05(m, 2H), 5.33(s, 2H), 3.97(s, 3H), 3.86(t, 2H), 1.47(m, 2H), 0.77(t, 3H).	349.3 [M+1]
861		2-((2-(2,3-dichlorophenyl)-1H-imidazol-1-yl)methyl)-3-propyl-3H-imidazo[4,5-b]pyridine	¹ H NMR, δppm (CDCl ₃): 8.38(dd, 1H), 8.02(dd, 1H), 7.59(dd, 1H), 7.39(dd, 1H), 7.21-7.29(m, 3H), 7.10(d, 1H), 5.29(s, 2H), 3.90 (t, 2H), 1.53 (m, 2H), 0.79(t, 3H).	386.3
862		2-((2-(2-fluoropyridin-3-yl)-1H-imidazol-1-yl)methyl)-3-propyl-3H-imidazo[4,5-b]pyridine	¹ H NMR, δppm (CDCl ₃): 8.39(dd, 1H), 8.35(m, 1H), 8.12(m, 1H), 8.02(dd, 1H), 7.35(m, 1H), 7.22-7.29(m, 2H), 7.10(d, 1H), 5.43(s, 2H), 3.95 (t, 2H), 1.54 (m, 2H), 0.80(t, 3H).	337.2
863		2-((2-(2,3-difluorophenyl)-1H-imidazol-1-yl)methyl)-3-ethyl-3H-imidazo[4,5-b]pyridine	¹ H NMR, δppm (CDCl ₃): 8.40(dd, 1H), 8.04(dd, 1H), 7.22-7.43(m, 5H), 7.08(d, 1H), 5.40(s, 2H), 3.99 (q, 2H), 1.05(t, 3H).	340.1
864		2-((2-(2,3-difluorophenyl)-1H-imidazol-1-yl)methyl)-3-propyl-3H-imidazo[4,5-b]pyridine	¹ H NMR, δppm (CDCl ₃): 8.39(dd, 1H), 8.03(dd, 1H), 7.22-7.42(m, 5H), 7.09(d, 1H), 5.39(s, 2H), 3.86(t, 2H), 1.50(m, 2H), 0.78(t, 3H)	354.2










TABLE 6				
Cpd#	STRUCTURE	IUPAC NAME	NMR	MS (m/z)
865		2-((2-(4-chloro-1-methyl-1H-pyrazol-3-yl)-1H-imidazol-1-yl)methyl)-3-propyl-3H-imidazo[4,5-b]pyridine	¹ H NMR (CDCl ₃): 8.38(1H, dd), 8.04(1H, dd), 7.52(1H, s), 7.26-7.21(2H, m), 7.05(1H, d), 5.98(2H, s), 4.15(2H, t), 3.94(3H, s), 1.62-1.52(2H, m), 0.80(3H, t)	356.2 [M+1]
866		3-cyclobutyl-2-((2-(6-fluoropyridin-2-yl)methyl)-1H-imidazol-1-yl)methyl)-3H-imidazo[4,5-b]pyridine	dihydrochloride ¹ H NMR (d ₆ DMSO): 8.39(1H, dd), 8.33(1H, dd), 8.25(1H, q), 8.03(1H, d), 7.95(1H, d), 7.87(1H, dd), 7.33(1H, dd), 7.21(1H, dd), 6.34(2H, s), 5.18(1H, m), 3.28-1.20(2H, m), 2.58-2.42(2H, m), 2.00-1.86(2H, m)	349.3 [M+1]
867		2-((1-((3-propyl-3H-imidazo[4,5-b]pyridin-2-yl)methyl)-1H-imidazol-2-yl)isonicotinonitrile	dihydrochloride ¹ H NMR (d ₆ DMSO): 8.78(1H, s), 8.72(1H, d), 8.32(1H, dd), 7.99(1H, d), 7.93-7.86(3H, m), 7.21(1H, dd), 6.34(2H, s), 4.37(2H, t), 1.91-1.82(2H, m), 0.93(3H, t)	344.2 [M+1]
868		2-((1-((3-propyl-1H-imidazo[4,5-c]pyridin-2-yl)methyl)-1H-imidazol-2-yl)isonicotinonitrile	¹ H NMR (CDCl ₃) δ 9.07 (s, 1 H), 8.68 (d, 1 H), 8.60 (s, 1 H), 8.44 (d, 1 H), 7.45 (d, 1 H), 7.29 (d, 1 H), 7.26 (d, 1 H), 7.22 (d, 1 H), 6.37 (s, 2 H), 4.16 (t, 2 H), 1.63 (q, 2 H), 0.81 (t, 3 H)	m/e 344
869		2-((2-(2-methoxyphenyl)-1H-imidazol-1-yl)methyl)-3-propyl-3H-imidazo[4,5-b]pyridine	dihydrochloride ¹ H NMR (d ₆ DMSO): 8.33(1H, dd), 8.02-7.99(2H, m), 7.93(1H, d), 7.65-7.59(1H, m), 7.51-7.48(1H, m), 7.29-7.21(2H, m), 7.11-7.06(1H, m), 5.75(2H, s), 4.12(2H, t), 3.64(3H, s), 1.69-1.61(2H, m), 0.74(3H, t)	348.3 [M+1]
870		2-((2-(3-methoxyphenyl)-1H-imidazol-1-yl)methyl)-3-propyl-3H-imidazo[4,5-b]pyridine	dihydrochloride ¹ H NMR (d ₆ DMSO): 8.35(1H, dd), 8.04-7.98(2H, m), 7.93(1H, d), 7.48(1H, t), 7.35-7.19(4H, m), 5.95(2H, s), 4.22(2H, t), 3.69(3H, s), 1.79-1.68(2H, m), 0.81(3H, t)	348.39 [M+1]
871		6-((1-((3-(2-fluoroethyl)-3H-imidazo[4,5-b]pyridin-2-yl)methyl)-1H-imidazol-2-yl)pyridine-2-carbonitrile	¹ H NMR (CDCl ₃): line list provided 4.74 (m, 1H), 4.84-4.95 (m, 3H), 6.16 (s, 2H), 7.19-7.28 (m, 2H), 7.39 (s, 1H), 7.54 (d, 1H), 7.83 (t, 1H), 7.89 (d, 1H), 8.34 (m, 1H), 8.53 (d, 1H)	348 [M+1]
872		2-((2-(3,4-difluorophenyl)-1H-imidazol-1-yl)methyl)-3-(2-fluoroethyl)-3H-imidazo[4,5-b]pyridine	¹ H NMR (CDCl ₃): line list provided 4.33 (t, 1H), 4.42 (t, 1H), 4.56 (t, 1H), 4.73 (t, 1H), 5.51 (s, 2H), 7.07 (s, 1H), 7.18-7.26 (m, 2H), 7.39 (m, 1H), 7.55 (m, 1H), 8.07 (d, 1H), 8.37 (s, 1H)	358 [M+1]
873		3-(2-fluoroethyl)-2-((2-(2,4,5-trifluorophenyl)-1H-imidazol-1-yl)methyl)-3H-imidazo[4,5-b]pyridine	¹ H NMR (CDCl ₃): line list provided 4.23 (t, 1H), 4.31 (t, 1H), 4.50 (t, 1H), 4.66 (t, 1H), 5.40 (s, 2H), 7.01-7.09 (m, 2H), 7.20-7.26 (m, 2H), 7.43 (m, 1H), 7.99 (d, 1H), 8.32 (d, 1H)	376 [M+1]










TABLE 6				
Cpd#	STRUCTURE	IUPAC NAME	NMR	MS (m/z)
874		3-(2-fluoroethyl)-2-((2-(5-fluoro-2-methylphenyl)-1H-imidazol-1-yl)methyl)-3H-imidazo[4,5-b]pyridine	¹ H NMR (CDCl ₃): line list provided 2.16 (s, 3H), 4.19 (t, 1H), 4.26 (t, 1H), 4.49 (t, 1H), 4.57 (t, 1H), 5.28 (s, 2H), 7.02-7.07 (2H), 7.14 (s, 1H), 7.21-7.26 (m, 3H), 8.02 (d, 1H), 8.35 (d, 1H)	354
875		1-propyl-2-((2-(1,3-thiazol-2-yl)-1H-imidazol-1-yl)methyl)-1H-imidazo[4,5-c]pyridine	¹ H NMR (CDCl ₃) δ 9.07 (s, 1 H), 8.41 (d, 1 H), 7.82 (d, 1 H), 7.38 (d, 1 H), 7.25 (d, 1 H), 7.19 (s, 1 H), 7.12 (s, 1 H), 6.35 (s, 2 H), 4.14 (t, 2 H), 1.53 (q, 2 H), 0.74 (t, 3 H)	m/e 325
876		1-ethyl-2-((2-(1,3-thiazol-2-yl)-1H-imidazol-1-yl)methyl)-1H-imidazo[4,5-c]pyridine	¹ H NMR (CDCl ₃) δ 9.08 (s, 1 H), 8.43 (d, 1 H), 7.85 (d, 1 H), 7.38 (d, 1 H), 7.38 – 7.26 (m, 1 H), 7.17 (s, 1 H), 7.14 (s, 1 H), 6.35 (s, 2 H), 4.25 (q, 2 H), 1.11 (t, 3 H)	m/e 311
877		2-((2-(6-fluoropyridin-2-yl)-1H-imidazol-1-yl)methyl)-6-methyl-1-propyl-1H-imidazo[4,5-c]pyridine	¹ H NMR (CDCl ₃) δ 8.93 (s, 1 H), 8.16 (d, 1 H), 7.87 (dd, 1 H), 7.20 (d, 1 H), 7.16 (d, 1 H), 7.10 (d, 1 H), 6.87 (d, 1 H), 6.26 (s, 2 H), 4.19 (t, 2 H), 2.65 (s, 3 H), 1.65 (q, 2 H), 0.83 (t, 3 H)	m/e 351
878		2-((2-(3-fluorophenyl)-1H-imidazol-1-yl)methyl)-6-methyl-1-propyl-1H-imidazo[4,5-c]pyridine	¹ H NMR (CDCl ₃) δ 8.95 (s, 1 H), 7.49 – 7.36 (m, 3 H), 7.19 – 7.15 (m, 2 H), 7.04 (m, 2 H), 5.48 (s, 2 H), 3.64 (t, 2 H), 2.64 (s, 3 H), 1.40 (q, 2 H), 0.73 (t, 3 H)	
879		3-(cyclopropylmethyl)-2-((2-(1,3-thiazol-2-yl)-1H-imidazol-1-yl)methyl)-3H-imidazo[4,5-b]pyridine	dihydrochloride ¹ H NMR (d ₆ DMSO): 8.35(1H, dd), 7.95(1H, dd), 7.90-7.87(2H, m), 7.81(1H, s), 7.52(1H, s), 7.27(1H, dd), 6.35(2H, s), 4.31(2H, d), 1.42-1.38(1H, m), 0.55-0.52(4H, m)	335.1 [M-1]
880		2-((1-((3-propyl-3H-imidazo[4,5-b]pyridin-2-yl)methyl)-1H-imidazol-2-yl)benzonitrile	dihydrochloride ¹ H NMR (d ₆ DMSO): 8.31(1H, d), 8.15(1H, s), 8.05-8.03(2H, m), 7.94-7.92(2H, m), 7.85-7.77(2H, m), 7.27-7.24(1H, m), 5.97(1H, s), 4.14(2H, t), 1.72-1.64(2H, m), 0.75(3H, t)	343.3 [M+1]
881		2-((2-(5-methylisoxazol-3-yl)-1H-imidazol-1-yl)methyl)-3-propyl-3H-imidazo[4,5-b]pyridine	¹ H NMR, δppm (CDCl ₃): 8.39(dd, 1H), 8.04(dd, 1H), 7.19-7.27(m, 3H), 6.65(d, 1H), 6.10(s, 2H), 4.26(t, 2H), 1.63(m, 2H), 0.83(t, 3H)	323.2
882		3-isopropyl-2-((2-(1,3-thiazol-2-yl)-1H-imidazol-1-yl)methyl)-3H-imidazo[4,5-b]pyridine	¹ H NMR, δppm (CDCl ₃): 8.37(dd, 1H), 8.02(dd, 1H), 7.84 (d, 1H), 7.39(d, 1H), 7.11-7.23(m, 3H), 6.32(s, 2H), 4.92(m, 1H), 1.56(d, 6H)	325.4

TABLE 6				
Cpd#	STRUCTURE	IUPAC NAME	NMR	MS (m/z)
883		1-(1-ethyl-2-[[2-(1,3-thiazol-2-yl)-1H-imidazol-1-yl]methyl]-1H-benzimidazol-5-yl)ethanone	¹ H NMR, δppm (CDCl ₃): 8.39(d, 1H), 8.00(dd, 1H) 7.84 (d, 1H), 7.39(d, 1H), 7.37(d, 1H), 7.18(d, 1H), 7.13(d, 1H), 6.35(s, 2H), 4.27(q, 2H), 2.67(s, 3H), 1.12(t, 3H)	352.2
884		2-[[2-(2,3-difluorophenyl)-1H-imidazol-1-yl]methyl]-1-ethyl-1H-benzimidazole-5-carbonitrile	¹ H NMR, δppm (CDCl ₃): 8.09(dd, 1H) 7.55 (dd, 1H), 7.19-7.40(m, 5H), 7.04(d, 1H), 5.40(s, 2H), 3.86(q, 2H), 1.02(t, 3H)	
885		2-[[2-(5-fluoro-2-methylphenyl)-1H-imidazol-1-yl]methyl]-3-propyl-3H-imidazo[4,5-b]pyridine	¹ H NMR, δppm (CDCl ₃): 8.39(dd, 1H), 8.03(dd, 1H), 7.20-7.31(m, 3H), 7.07-7.13(m, 3H), 5.24(s, 2H), 3.88(t, 2H), 2.22(s, 3H), 1.54(m, 2H), 0.78(t, 3H)	350.3
886		2-[[2-(2-fluoro-5-methylphenyl)-1H-imidazol-1-yl]methyl]-3-propyl-3H-imidazo[4,5-b]pyridine	¹ H NMR, δppm (CDCl ₃): 8.38(dd, 1H), 8.04(dd, 1H), 7.44(dd, 1H), 7.22-7.30(m, 2H), 7.19(d, 1H), 7.12(dd, 1H), 7.05(d, 1H), 5.37(s, 2H), 3.84(t, 2H), 2.36 (s, 3H), 1.46(m, 2H), 0.76(t, 3H)	350.3
887		2-[[2-(5-fluoro-2-methoxyphenyl)-1H-imidazol-1-yl]methyl]-3-propyl-3H-imidazo[4,5-b]pyridine	¹ H NMR, δppm (CDCl ₃): 8.39(dd, 1H), 8.03(dd, 1H), 7.15-7.32(m 4H), 7.03(d, 1H), 6.96(dd, 1H), 5.30(s, 2H), 3.84(t, 2H), 3.79(s, 3H), 1.46(m, 2H), 0.76(t, 3H)	366.2
888		1-ethyl-2-[[2-[3-(trifluoromethyl)phenyl]-1H-imidazol-1-yl]methyl]-1H-imidazo[4,5-c]pyridine	dihydrochloride ¹ H NMR (d ₆ DMSO): 10.02(1H, s), 8.70(1H, d), 8.37(1H, d), 8.08(1H, s), 7.97-7.94(3H, m), 7.82(1H, s), 7.75(1H, t), 6.09(2H, s), 4.45(2H, q), 1.32(3H, t).	
889		1-ethyl-2-[[2-(5-fluoro-2-methylphenyl)-1H-imidazol-1-yl]methyl]-1H-benzimidazole-5-carbonitrile	hydrochloride ¹ H NMR (d ₆ DMSO): 8.10(1H, s), 8.07(1H, d), 7.95(1H, d), 7.78(1H, d), 7.64(1H, dd), 7.52(1H, dd), 7.38-7.35(2H, m), 5.82(2H, s), 4.23(2H, q), 2.17(3H, s), 1.18(3H, t).	
890		2-[[2-(2,6-difluorophenyl)-1H-imidazol-1-yl]methyl]-1-propyl-1H-imidazo[4,5-c]pyridine	¹ H NMR (CDCl ₃) δ 9.05 (s, 1H), 8.43 (d, 1 H), 7.5 – 7.4 (m, 1 H), 7.27-7.23 (m, 2 H), 7.07 – 7.01 (m, 3 H), 5.34 (s, 2 H), 3.75 (t, 2H), 1.47 (q, 2 H), 0.77 (t, 3 H)	m/e 354
891		2-[[2-(2,6-difluorophenyl)-1H-imidazol-1-yl]methyl]-1-ethyl-1H-imidazo[4,5-c]pyridine	¹ H NMR (CDCl ₃) δ 9.05 (s, 1 H), 8.43 (d, 1 H), 7.52 – 7.42 (m, 1 H), 7.27 – 7.23 (m, 2 H), 7.08 – 7.01 (m, 3 H), 5.35 (s, 2 H), 3.85 (q, 2 H), 1.02 (t, 3 H)	m/e 340

TABLE 6				
Cpd#	STRUCTURE	IUPAC NAME	NMR	MS (m/z)
892		2-((2-(3,4-difluorophenyl)-1H-imidazol-1-yl)methyl)-1-ethyl-1H-imidazo[4,5-c]pyridine	¹ H NMR (CDCl ₃) δ 9.10 (s, 1 H), 8.46 (d, 1 H), 7.56 – 7.51 (m, 1 H), 7.42 – 7.30 (m, 1 H), 7.31 – 7.26 (m, 2 H), 7.16 (d, 1 H), 7.03 (d, 1 H), 5.48 (s, 2 H), 3.83 (q, 2 H), 1.05 (t, 3 H)	m/e 340
893		2-((2-(5-fluoro-2-methylphenyl)-1H-imidazol-1-yl)methyl)-1-propyl-1H-imidazo[4,5-c]pyridine	¹ H NMR (CDCl ₃) δ 9.07 (s, 1 H), 8.43 (d, 1 H), 7.31 – 7.23 (m, 3 H), 7.12 – 7.04 (m, 3 H), 5.24 (s, 2 H), 3.73 (t, 2 H), 2.21 (s, 3 H), 1.49 (q, 2 H), 0.75 (t, 3 H)	m/e 350
894		1-ethyl-2-((2-(5-fluoro-2-methylphenyl)-1H-imidazol-1-yl)methyl)-1H-imidazo[4,5-c]pyridine	¹ H NMR (CDCl ₃) δ 9.08 (s, 1 H), 8.44 (d, 1 H), 7.31 – 7.24 (m, 3 H), 7.12 – 7.04 (m, 3 H), 5.25 (s, 1 H), 3.83 (q, 2 H), 2.22 (s, 3 H), 1.06 (t, 3 H)	m/e 336
895		2-((2-(1,3-thiazol-2-yl)-1H-imidazol-1-yl)methyl)-3-(2,2,2-trifluoroethyl)-3H-imidazo[4,5-b]pyridine	¹ H NMR (CDCl ₃): line list provided 5.32 (q, 2H), 6.19 (s, 2H), 7.19 (s, 1H), 7.26 (m, 2H), 7.43 (s, 1H), 7.79 (d, 1H), 8.04 (d, 1H), 8.41 (m, 1H)	365 [M+1]
896		2-((2-(6-fluoropyridin-2-yl)-1H-imidazol-1-yl)methyl)-3-(2,2,2-trifluoroethyl)-3H-imidazo[4,5-b]pyridine	HCl salt ¹ H NMR (CD ₃ OD): 5.42 (q, 2H), 6.40 (s, 2H), 7.24 (d, 1H), 7.37 (m, 1H), 7.90 (m, 2H), 8.03 (m, 2H), 8.22 (m, 1H), 8.43 (m, 1H)	377 [M+1]
897		2-((2-(3,4-difluorophenyl)-1H-imidazol-1-yl)methyl)-3-ethyl-3H-imidazo[4,5-b]pyridine	¹ H NMR, δppm (CDCl ₃): 8.39(dd, 1H), 8.04(dd, 1H), 7.54(m 1H), 7.40(m, 1H), 7.26-7.32(m, 2H), 7.16(d, 1H), 7.04(d, 1H), 5.47(s, 2H), 4.02(q, 2H), 1.09(t, 3H)	340.2
898		2-((2-(1-methyl-1H-pyrazol-3-yl)-1H-imidazol-1-yl)methyl)-3-propyl-3H-imidazo[4,5-b]pyridine	¹ H NMR (CDCl ₃): 8.38(1H, dd), 8.05(1H, dd), 7.42(1H, d), 7.23(1H, dd), 7.09(1H, d), 7.04(1H, d), 6.91(1H, d), 6.21(2H, s), 4.19(2H, t), 3.97(3H, s), 1.61-1.50(2H, m), 0.76(3H, t)	322.3 [M+1]
899		2-((2-(2,3-difluorophenyl)-1H-imidazol-1-yl)methyl)-1-ethyl-1H-imidazo[4,5-c]pyridine	dihydrochloride ¹ H NMR (d ₆ DMSO): 9.44(1H, s), 8.70(1H, d), 8.37(1H, d), 8.04(1H, s), 7.89(1H, s), 7.73(1H, q), 7.49(1H, d), 7.37(1H, t), 6.05(2H, s), 4.44(2H, q), 1.30(3H, t)	340.2
900		1-propyl-2-((2-quinolin-2-yl)-1H-imidazol-1-yl)methyl)-1H-benzimidazole-5-carbonitrile	dihydrochloride ¹ H NMR (d ₆ DMSO): 8.60(1H, d), 8.359(1H, d), 8.02-7.99(2H, m), 7.88-7.85(2H, m), 7.79-7.60(5H, m), 6.54(2H, s), 4.47(2H, t), 1.85-1.77(2H, m), 0.89(3H, t)	393.4 [M+1]

TABLE 6				
Cpd#	STRUCTURE	IUPAC NAME	NMR	MS (m/z)
901		2-[(2-(2,4-difluorophenyl)-1H-imidazol-1-yl)methyl]-3-ethyl-3H-imidazo[4,5-b]pyridine	dihydrochloride ¹ H NMR (d6 DMSO): 8.35-8.33(1H, m), 8.07-8.06(1H, m), 7.99-7.96(2H, m), 7.86-7.80(1H, m), 7.63-7.57(1H, m), 7.35-7.32(1H, m), 7.28-7.25(1H, m), 5.89(2H, s), 4.27(2H, q), 1.28(3H, t)	340.2 [M+1]
902		1-Propyl-2-(2-pyrimidin-2-yl-imidazol-1-ylmethyl)-1H-benzimidazole-5-carbonitrile	dihydrochloride ¹ H NMR (d6 DMSO): 8.92(2H, d), 8.18(1H, d), 8.00-7.99(2H, m), 7.88(1H, dd), 7.65-7.62(2H, m), 6.46(2H, s), 4.42(2H, t), 1.93-1.84(2H, m), 0.97(3H, t)	344.2 [M+1]
903		3-ethyl-2-[(2-(2,3,6-trifluorophenyl)-1H-imidazol-1-yl)methyl]-3H-imidazo[4,5-b]pyridine	¹ H NMR, δppm (CDCl3): 8.40(dd, 1H), 8.02(dd, 1H), 7.24-7.32(m, 3H), 7.11(d, 1H), 7.00(m, 1H), 5.36(s, 2H), 4.02(q, 2H), 1.09(t, 3H)	358.2
904		3-propyl-2-[(2-thien-3-yl-1H-imidazol-1-yl)methyl]-3H-imidazo[4,5-b]pyridine	¹ H NMR, δppm (CDCl3): 8.40(dd, 1H), 8.05(dd, 1H), 7.66(dd, 1H), 7.43-7.49(m, 2H), 7.25(dd, 1H), 7.14(d, 1H), 7.02(d, 1H), 5.55(s, 2H), 3.91(t, 2H), 1.51(m, 2H), 0.79(t, 3H)	324.3
905		1-(1-ethyl-2-[(2-pyrimidin-2-yl-1H-imidazol-1-yl)methyl]-1H-benzimidazol-5-yl)ethanone	¹ H NMR, δppm (CDCl3): 8.87(d, 2H), 8.41(dd, 1H), 8.02(dd, 1H), 7.40(dd, 1H), 7.26-7.31(m, 2H), 7.16(d, 1H), 6.40(s, 2H), 4.27(q, 2H), 2.69(s, 3H), 1.17(t, 3H)	
906		1-ethyl-2-[(2-(6-fluoropyridin-2-yl)-1H-imidazol-1-yl)methyl]-6-methyl-1H-imidazo[4,5-c]pyridine	¹ H NMR (CDCl3) δ 8.93 (s, 1 H), 8.16 (d, 1 H), 7.87 (dd, 1 H), 7.18 (d, 1 H), 7.12 (s, 1 H), 6.87 (d, 1 H), 6.26 (s, 2 H), 4.30 (q, 2 H), 2.65 (s, 3 H), 1.22 (t, 3 H)	m/e 337
907		1-ethyl-2-[(2-(3-fluorophenyl)-1H-imidazol-1-yl)methyl]-6-methyl-1H-imidazo[4,5-c]pyridine	dihydrochloride ¹ H NMR (DMSO-d6) δ 9.28 (s, 1 H), 8.24 (s, 1 H), 8.23 (d, 1 H), 7.94 (d, 1 H), 7.69-7.48 (m, 4 H), 6.12 (s, 2 H), 4.39 (q, 2 H), 2.77 (s, 3 H), 1.35 (t, 3 H);	m/e 336
908		3-ethyl-2-[(2-[2-fluoro-5-(trifluoromethyl)phenyl]-1H-imidazol-1-yl)methyl]-3H-imidazo[4,5-b]pyridine	¹ H NMR (CDCl3): line list provided 1.08 (t, 3H), 3.99 (q, 2H), 5.38 (s, 2H), 7.10 (s, 1H), 7.22-7.39 (m, 2H), 7.75 (m, 1H), 7.93 (m, 1H), 8.02 (d, 1H), 8.38 (d, 1H)	390 (M + 1)
909		2-[(2-(2-methylphenyl)-1H-imidazol-1-yl)methyl]-3-propyl-3H-imidazo[4,5-b]pyridine	¹ H NMR (CDCl3): line list provided 0.73 (t, 3H), 1.49 (m, 2H), 2.24 (s, 3H), 3.81 (m, 2H), 5.22 (s, 2H), 7.10 (s, 1H), 7.18-7.36 (m, 6H), 8.02 (d, 1H), 8.37 (d, 1H)	332 [M+1]

TABLE 6				
Cpd#	STRUCTURE	IUPAC NAME	NMR	MS (m/z)
910		6-{1-[(3-cyclopropyl-3H-imidazo[4,5-b]pyridin-2-yl)methyl]-1H-imidazol-2-yl}pyridine-2-carbonitrile	H-1 NMR (CDCl ₃): line list provided 1.28 (m, 4H), 3.37 (m, 1H), 3.48 (s, 2H), 6.36 (s, 2H), 7.15-7.28 (m, 3H), 7.56 (d, 1H), 7.86 (m, 2H), 8.38 (m, 1H), 8.54 (d, 1H)	342 [M+1]; 340 (M-1)
911		4-fluoro-3-{1-[(3-propyl-3H-imidazo[4,5-b]pyridin-2-yl)methyl]-1H-imidazol-2-yl}benzonitrile	CDCl ₃ : 8.38 (dd, J = 1.48, 4.81 Hz, 1H), 7.98 (dt, J = 1.48, 7.35 Hz, 2H), 7.72-7.78 (m, 1H), 7.35 (t, J = 9.07 Hz, 1H), 7.22-7.26 (m, 2H), 7.11 (d, J = 1.1 Hz, 1H), 5.37 (s, 2H), 3.93 (t, J = 7.69 Hz, 2H), 1.54 (dq, J = 7.42, 7.69 Hz, 2H),	361.2 [M+1]
912		1-propyl-2-[(2-{3-(trifluoromethyl)phenyl}-1H-imidazol-1-yl)methyl]-1H-imidazo[4,5-c]pyridine	¹ H NMR (CDCl ₃): 9.09(1H, s), 8.43(1H, d), 7.93(1H, s), 7.84(1H, d), 7.72(1H, d), 7.62(1H, t), 7.25(1H,), 7.19(1H, d), 7.09(1H, d), 5.50(2H, s), 3.63(2H, t), 1.49-1.39(2H, m), 0.75(3H, t).	
913		2-{[2-(6-fluoropyridin-2-yl)-1H-imidazol-1-yl)methyl]-1-propyl-1H-benzimidazole-5-carbohydrazide	CD ₃ OD: 8.10-8.29 (m, 6H), 8.03 (s, 1H), 7.29 (d, 1H), 6.4 (s, 2H), 4.59-4.65 (m, 2H), 2.0-2.12 (m, 2H), 1.06-1.07 (m, 3H).	394.3 [M+1]

TABLE 7				
Cmp. #	STRUCTURE	IUPAC NAME	NMR	MS (m/z)
914		1-ethyl-2-([1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl)-5-(morpholin-4-ylmethyl)-1H-benzimidazole		420.3 [M+1]
915		ethyl 1-ethyl-2-([1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl)-1H-benzimidazole-5-carboxylate		393.3 [M+1]
916		1-ethyl-2-([1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl)-1H-benzimidazole-5-carboxylic acid		365.2 [M+1]
917		N-[2-(dipropylamino)ethyl]-1-ethyl-2-([1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl)-1H-benzimidazole-5-carboxamide		491.4 [M+1]
918		N,1-diethyl-2-([1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl)-1H-benzimidazole-5-carboxamide		392.3 [M+1]
919		1-ethyl-2-([1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl)-N-phenyl-1H-benzimidazole-5-carboxamide		392.3 [M+1]
920		1-ethyl-2-([1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl)-5-[(4-methylpiperazin-1-yl)methyl]-1H-benzimidazole	¹ H NMR (CDCl ₃): 7.65 (d, 1H), 7.61 (d, 1H), 7.45 (m, 1H), 7.22-7.31 (m, 4H), 7.11 (m, 1H), 6.16 (d, 1H), 4.31 (s, 2H), 3.96 (q, 2H), 3.62 (s, 2H), 2.40-2.60 (m, 8H), 2.25 (s, 3H), 1.22 (t, 3H)	
921		1-ethyl-5-[(4-ethylpiperazin-1-yl)methyl]-2-([1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl)-1H-benzimidazole	¹ H NMR (CDCl ₃): 7.65 (d, 1H), 7.62 (d, 1H), 7.45 (m, 1H), 7.23-7.30 (m, 4H), 7.13 (m, 1H), 6.16 (d, 1H), 4.31 (s, 2H), 3.96 (q, 2H), 3.62 (s, 2H), 2.40-2.60 (m, 10H), 1.21 (t, 3H), 1.078 (t, 3H)	

TABLE 7				
Cmp. #	STRUCTURE	IUPAC NAME	NMR	MS (m/z)
922		5-[(4-cyclopentylpiperazin-1-yl)methyl]-1-ethyl-2-[[1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl]-1H-benzimidazole	¹ H NMR (CDCl ₃): 7.65 (d, 1H), 7.61 (d, 1H), 7.45 (m, 1H), 7.22-7.31 (m, 4H), 7.13 (m, 1H), 6.15 (d, 1H), 4.31 (s, 2H), 3.96 (q, 2H), 3.63 (s, 2H), 2.40-2.60 (m, 9H), 1.31-1.90 (m, 8H), 1.21 (t, 3H.)	
923		5-[(4-cycloheptyl) piperazin-1-yl]methyl]-1-ethyl-2-[[1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl]-1H-benzimidazole		515.5 [M+1]
924		2-[[1-(3-chlorophenyl)-1H-pyrazol-5-yl]methyl]-1-ethyl-5-(morpholin-4-ylmethyl)-1H-benzimidazole		436.3 [M+1]; 434.3 [M-1]
925		1'-[[1-ethyl-2-[[1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl]-1H-benzimidazol-5-yl]methyl]-1,4'-bipiperidine	¹ H NMR (CDCl ₃): 7.63 (d, 1H), 7.61 (d, 1H), 7.44 (m, 1H), 7.2307.31 (m, 4H), 7.13 (m, 1H), 6.16 (d, 1H), 4.31 (s, 2H), 3.96 (q, 2H), 3.61 (s, 2H), 2.96 (m, 2H), 2.50 (m, 4H), 1.40-2.31 (m, 13H), 1.23 (t, 3H)	
926		1-ethyl-2-[[1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl]-5-[(4-pyrrolidin-1-yl)piperidin-1-yl]methyl]-1H-benzimidazole		
927		1-ethyl-2-[[1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl]-5-[[4-phenylpiperazin-1-yl]methyl]-1H-benzimidazole	¹ H NMR (CDCl ₃): 7.69 (d, 1H), 7.62 (s, 1H), 7.45 (m, 1H), 7.21-7.33 (m, 6H), 7.13 (m, 1H), 6.82-6.94 (m, 3H), 6.16 (d, 1H), 4.33 (s, 2H), 3.97 (q, 2H), 3.69 (s, 2H), 3.15-3.23 (m, 4H), 2.59-2.67 (m, 4H), 1.21 (t, 3H)	
928		2-[[1-(3-chlorophenyl)-1H-pyrazol-5-yl]methyl]-5-[(4-cycloheptyl)piperazin-1-yl]carbonyl]-1-ethyl-1H-benzimidazole		545.4 [M+1]; 543.4 [M-1]
929		1-ethyl-2-[[1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl]-1H-benzimidazole-5-carboxamide		
930		1-ethyl-5-fluoro-2-[[1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl]-1H-benzimidazole	(L)-Tartrate salt ¹ H NMR (CD ₃ OD): 7.67 (s, 1H), 7.05-7.46 (m, 7H), 6.31 (s, 1H), 4.53 (s, 2H), 4.47 (s, 2H), 4.14 (q, 2H), 1.21 (t, 3H)	

TABLE 7				
Cmp. #	STRUCTURE	IUPAC NAME	NMR	MS (m/z)
931		5-chloro-1-ethyl-2-([1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl)-1H-benzimidazole	(L)-Tartrate salt 1H NMR (d6-DMSO): 7.65 (s, 1H), 7.21-7.58 (m, 7H), 6.26 (s, 1H), 4.48 (s, 2H), 4.30 (s, 2H), 4.16 (q, 2H), 1.15 (t, 3H)	
932		5-bromo-1-ethyl-2-([1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl)-1H-benzimidazole	(L)-Tartrate salt 1H NMR (d6-DMSO): 7.72 (d, 1H), 7.65 (d, 1H), 7.21-7.53 (m, 6H), 6.26 (s, 1H), 4.48 (s, 2H), 4.30 (s, 2H), 4.16 (q, 2H), 1.14 (t, 3H)	
933		1-ethyl-2-([1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl)-5-phenyl-1H-benzimidazole	1H NMR (CDCl3): 7.96 (d, 1H), 7.27-7.66 (m, 11H), 7.14 (m, 1H), 6.21 (d, 1H), 4.38 (s, 2H), 4.01 (q, 2H), 1.25 (t, 3H)	
934		2-([1-(2,5-difluorophenyl)-1H-pyrazol-5-yl]methyl)-1-ethyl-1H-benzimidazole-5-carbonitrile		364.2 [M+1]
935		2-([1-(3-chlorophenyl)-1H-pyrazol-5-yl]methyl)-1-ethyl-1H-benzimidazole-5-carbonitrile	1H NMR (CDCl3): 8.05 (d, 1H), 7.36-7.65 (m, 7H), 6.21 (d, 1H), 4.36 (s, 2H), 4.02 (q, 2H), 1.24 (t, 3H)	
936		2-([1-(2,5-difluorophenyl)-1H-pyrazol-5-yl]methyl)-1-ethyl-1H-benzimidazole-5-amine		354.2 [M+1]
937		2-([1-(2,5-difluorophenyl)-1H-pyrazol-5-yl]methyl)-1-ethyl-5-(4H-1,2,4-triazol-4-yl)-1H-benzimidazole		405.95 [M+1]
938		1-ethyl-2-([1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl)-5-(1H-1,2,4-triazol-1-ylmethyl)-1H-benzimidazole		402.4 [M+1]
939		1-ethyl-2-([1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl)-5-(1H-tetrazol-1-ylmethyl)-1H-benzimidazole		403.2 [M+1]; 401.4 [M-1]

TABLE 7				
Cmp. #	STRUCTURE	IUPAC NAME	NMR	MS (m/z)
940		1-ethyl-2-([1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl)-5-(2H-tetrazol-2-ylmethyl)-1H-benzimidazole		403.4 [M+1]; 401.4 [M-1]
941		1-ethyl-2-([1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl)-5-[(methylthio)methyl]-1H-benzimidazole		381.3 [M+1]; 379.3 [M-1]
942		2-([1-(2,5-difluorophenyl)-1H-pyrazol-5-yl]methyl)-1-ethyl-5-nitro-1H-benzimidazole	Mesylate 1H NMR (d6 DMSO) 8.38 (1H, d), 8.15-8.12(1H,m), 7.77(1H,d), 7.71(1H,s), 7.51-7.34(3H, m), 6.40(1H, s), 4.21 (2H, q), 2.32(3H, s), 1.11(3H, t).	384.2 [M+1]
943		1-ethyl-2-([1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl)-5-(1,3-thiazol-5-yl)-1H-benzimidazole	Hydrochloride 1H NMR (CD3OD): 9.725 (s, 1H), 8.67 (s, 1H), 8.13 (s, 1H), 8.01-8.08 (m, 2H), 7.79 (s, 1H), 7.21-7.57 (m, 4H), 6.59 (d, 1H), 4.97 (s, 2H), 4.50 (q, 2H), 1.39 (t, 3H)	
944		1-ethyl-2-([1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl)-N-methoxy-N-methyl-1H-benzimidazole-5-carboxamide	1H NMR (CDCl3): 8.13 (d, 1H), 7.63-7.70 (m, 2H), 7.24-7.49 (m, 4H), 7.13 (m, 1H), 6.19 (d, 1H), 4.36 (s, 2H), 4.00 (q, 2H), 3.60 (s, 3H), 3.40 (s, 3H), 1.23 (t, 3H)	
945		3-(1-ethyl-2-([1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl)-1H-benzimidazol-5-yl)benzonitrile	1H NMR (CDCl3): 7.84-7.92 (m, 3H), 7.40-7.64 (m, 6H), 7.26-7.33 (m, 2H), 7.14 (m, 1H), 6.21 (d, 1H), 4.37 (s, 2H), 4.03 (q, 2H), 1.26 (t, 3H)	
946		N-[3-(1-ethyl-2-([1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl)-1H-benzimidazol-5-yl)phenyl]acetamide	1H NMR (CDCl3): 7.93 (d, 1H), 7.26-7.72 (m, 10H), 7.14 (m, 1H), 6.20 (d, 1H), 4.36 (s, 2H), 4.01 (q, 2H), 2.21 (s, 3H), 1.24 (t, 3H)	
947		1-ethyl-2-([1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl)-5-[4-(trifluoromethoxy)phenyl]-1H-benzimidazole	1H NMR (CDCl3): 7.92 (d, 1H), 7.27-7.65 (m, 10H), 7.14 (m, 1H), 6.20 (d, 1H), 4.37 (s, 2H), 4.08 (q, 2H), 1.25 (t, 3H)	
948		5-(3-chloro-4-fluorophenyl)-1-ethyl-2-([1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl)-1H-benzimidazole	1H NMR (CDCl3): 7.897 (d, 1H), 7.63-7.66 (m, 2H), 7.11-7.50 (m, 8H), 6.21 (d, 1H), 4.36 (s, 2H), 4.02 (q, 2H), 1.25 (t, 3H)	

TABLE 7				
Cmp. #	STRUCTURE	IUPAC NAME	NMR	MS (m/z)
949		1-ethyl-5-(3-fluorophenyl)-2-([1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl)-1H-benzimidazole	¹ H NMR (CDCl ₃): 7.95 (d, 1H), 7.64 (d, 1H), 7.27-7.53 (m, 8H)/ 7.14 (m, 2H), 7.03 (m, 1H), 6.20 (d, 1H), 4.36 (s, 2H), 4.02 (q, 2H), 1.25 (t, 3H)	
950		1-ethyl-2-([1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl)-5-pyridin-3-yl-1H-benzimidazole	¹ H NMR (CDCl ₃): 8.90 (d, 1H), 8.59 (m, 1H), 7.90-7.95 (m, 2H), 7.64 (d, 1H), 7.25-7.53 (m, 6H), 7.14 (m, 1H), 6.21 (d, 1H), 4.37 (s, 2H), 4.03 (q, 2H), 1.26 (t, 3H)	
951		1-ethyl-2-([1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl)-5-pyridin-4-yl-1H-benzimidazole	¹ H NMR (CDCl ₃): 8.67 (d, 1H), 8.65 (d, 1H), 8.03 (d, 1H), 7.26-7.65 (m, 8H), 7.15 (m, 1H), 6.21 (d, 1H), 4.37 (s, 2H), 4.03 (q, 2H), 1.26 (t, 3H)	
952		2-([1-(2,5-difluorophenyl)-1H-pyrazol-5-yl]methyl)-1-ethyl-5-(methylidyne-lambda-5-azanyl)-1H-benzimidazole	¹ H NMR (CDCl ₃): 7.70-7.67(2H, m), 7.31-7.14(5H, m), 6.16(1H, s), 4.28(2H, s), 4.02(2H, q), 1.22(3H, t)	364.1 [M+1]
953		N-(2-([1-(2,5-difluorophenyl)-1H-pyrazol-5-yl]methyl)-1-ethyl-1H-benzimidazol-5-yl)-2,2,2-trifluoroacetamide		449.96 [M+1]
954		2-([1-(2,5-difluorophenyl)-1H-pyrazol-5-yl]methyl)-1-ethyl-5-(1H-tetrazol-1-yl)-1H-benzimidazole	¹ H NMR (DMSO): 10.07 (s, 1H), 8.07 (s, 1H), 7.86 (d, 1H), 7.79 (d, 1H), 7.71 (s, 1H), 7.36-7.57 (m, 3H), 6.4 (s, 2H), 4.46 (s, 2H), 4.21 (q, 2H), 1.14 (t, 3H)	
955		1-(1-ethyl-2-([1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl)-1H-benzimidazol-5-yl)ethanol		
956		2-([1-(3-chlorophenyl)-1H-pyrazol-5-yl]methyl)-1-ethyl-N-methoxy-N-methyl-1H-benzimidazole-5-carboxamide	¹ H NMR (CDCl ₃): 8.13 (d, 1H), 7.63-7.70 (m, 2H), 7.51 (m, 1H), 7.31-7.41 (m, 4H), 6.20 (d, 1H), 4.35 (s, 2H), 4.00 (q, 2H), 3.60 (s, 3H), 3.40 (s, 3H), 1.23 (t, 3H)	

TABLE 7				
Cmp. #	STRUCTURE	IUPAC NAME	NMR	MS (m/z)
957		2-([1-(2,5-difluorophenyl)-1H-pyrazol-5-yl]methyl)-1-ethyl-5-(5-methyl-1,3,4-oxadiazol-2-yl)-1H-benzimidazole		437.02 [M+1]
958		1-(2-([1-(3-chlorophenyl)-1H-pyrazol-5-yl]methyl)-1-ethyl-1H-benzimidazol-5-yl)ethanone	¹ H NMR (CDCl ₃): 8.37 (d, 1H), 8.03 (dd, 1H), 7.65 (d, 1H), 7.38-7.50 (m, 5H), 6.24 (d, 1H), 4.49 (s, 2H), 4.03 (q, 2H), 2.68 (s, 3H), 1.24 (t, 3H)	
959		2-([1-(2,5-difluorophenyl)-1H-pyrazol-5-yl]methyl)-1-ethyl-5-[5-(trifluoromethyl)-1H-tetrazol-1-yl]-1H-benzimidazole		474.99 [M+1]
960		5-(3,5-difluorophenyl)-1-ethyl-2-([1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl)-1H-benzimidazole	¹ H NMR (CDCl ₃): 7.91 (d, 1H), 7.64 (d, 1H), 7.27-7.50 (m, 5H), 7.10-7.18 (m, 3H), 6.77 (m, 1H), 6.20 (d, 1H), 4.36 (s, 2H), 4.02 (q, 2H), 1.25 (t, 3H)	
961		(1Z)-1-(2-([1-(3-chlorophenyl)-1H-pyrazol-5-yl]methyl)-1-ethyl-1H-benzimidazol-5-yl)ethanone O-methyloxime	¹ H NMR (CDCl ₃): 7.94 (d, 1H), 7.73 (dd, 1H), 7.73 (dd, 1H), 7.63 (d, 1H), 7.52 (m, 1H), 7.38-7.40 (m, 3H), 7.27 (m, 1H), 6.19 (d, 1H), 4.32 (s, 2H), 3.87-4.10 (m, 5H), 2.30 (s, 3H), 1.22 (t, 3H)	
962		2-([1-(2,5-difluorophenyl)-1H-pyrazol-5-yl]methyl)-1-ethyl-5-(1,3,4-oxadiazol-2-yl)-1H-benzimidazole		425.10 [M+1]
963		(1E)-1-(1-ethyl-2-([1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl)-1H-benzimidazol-5-yl)ethanone oxime	¹ H NMR (CD ₃ OD): 7.81 (d, 1H), 7.64-7.68 (m, 2H), 7.42-7.53 (m, 2H), 7.16-7.32 (m, 3H), 6.28 (d, 1H), 4.44 (s, 2H), 4.12 (q, 2H), 2.30 (s, 3H), 1.23 (t, 3H)	
964		(1E)-1-(1-ethyl-2-([1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl)-1H-benzimidazol-5-yl)ethanone O-ethyloxime	Syn and anti ¹ H NMR (CDCl ₃): 7.95 (d, 1H+H1), 7.73 (dd, 1H+H1), 7.63 (d, 1H+H1), 7.44 (m, 1H+H1), 7.24-7.33 (m, 3H+H1), [6.18 (d, 1H+H1), 7.13 (d, 1H+H1)], 4.33 (s, 2H), [4.26 (q, 2H), 4.13 (q, 2H)], 3.98 (q, 2H+2H1), [2.31 (s, 3H), 2.27 (s, 3H)], [1.35 (t, 3H), 1.28 (t, 3H)], 1.22 (t, 3H+H1)]	

TABLE 7				
Cmp. #	STRUCTURE	IUPAC NAME	NMR	MS (m/z)
965		2-([1-(2,5-difluorophenyl)-1H-pyrazol-5-yl]methyl)-1-ethyl-5-(5-pyridin-3-yl-1,3,4-oxadiazol-2-yl)-1H-benzimidazole	1H NMR (d6 DMSO): 9.40(1H, s), 8.87(1H, s), 8.63(1H, d), 8.41(1H, s), 8.20(1H, d), 8.01(1H, d), 7.76(2H, m), 7.59(1H, m), 7.48-7.41(2H, m), 6.50(1H, s), 4.61(2H, s), 4.31(2H, q), 1.19(3H, t)	484.5 [M+1]
966		2-(1-ethyl-2-([1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl)-1H-benzimidazol-5-yl)-4-methoxybenzaldehyde	1H NMR (CDCl3): 9.86 (s, 1H), 8.03 (d, 1H), 7.26-7.78 (m, 7H), 6.94-7.18 (m, 3H), 6.22 (dd, 1H), 4.38 (s, 2H), 4.04 (q, 2H), 3.91 (s, 3H), 1.27 (t, 3H)	
967		1-ethyl-2-([1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl)-N,N-dimethyl-1H-benzimidazole-5-carboxamide	1H NMR (CDCl3): 7.78 (s, 1H), 7.63 (s, 1H), 7.10-7.48 (m, 6H), 6.18 (s, 1H), 4.35 (s, 2H), 4.00 (q, 2H), 3.13 (s, 3H), 3.05 (s, 3H), 1.22 (t, 3H)	
968		5-(3,5-dimethylisoxazol-4-yl)-1-ethyl-2-([1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl)-1H-benzimidazole	1H NMR (CDCl3): 7.62-7.64 (m, 2H), 7.25-7.50 (m, 4H), 7.11-7.18 (m, 2H), 6.21 (d, 1H), 4.36 (s, 2H), 4.02 (q, 2H), 2.42 (s, 3H), 2.29 (s, 3H), 1.26 (t, 3H)	
969		5-bromo-2-([1-(3-chlorophenyl)-1H-pyrazol-5-yl]methyl)-1-ethyl-1H-benzimidazole	1H NMR (CDCl3): 7.87 (d, 1H), 7.63 (d, 1H), 7.50 (m, 1H), 7.35-7.42 (m, 4H), 7.18 (d, 1H), 6.18 (d, 1H), 4.31 (s, 2H), 3.96 (q, 2H), 1.21 (t, 3H)	
970		2-([1-(2,5-difluorophenyl)-1H-pyrazol-5-yl]methyl)-1-ethyl-5-(5-pyridin-4-yl-1,3,4-oxadiazol-2-yl)-1H-benzimidazole	Dihydrochloride 1H NMR (d6 DMSO): 8.98(2H, m), 8.43(1H, s), 8.35(2H, m), 8.19(1H, d), 7.99(1H, d), 7.75(1H, d), 7.59-7.56(1H, m), 7.50-7.41(2H, m), 6.49(1H, s), 4.58(2H, s), 4.30(2H, q), 1.19(3H, t)	484.5 [M+1]
971		(1E)-1-(2-([1-(3-chlorophenyl)-1H-pyrazol-5-yl]methyl)-1-ethyl-1H-benzimidazol-5-yl)ethanone oxime	1H NMR (CDCl3): 10.92 (br s, 1H), 8.58 (s, 1H), 7.52-7.64 (m, 3H), 7.29 (d, 1H), 6.20 (d, 1H), 4.36 (s, 2H), 3.97 (q, 2H), 2.38 (s, 3H), 1.21 (t, 3H)	
972		(1E)-1-(2-([1-(3-chlorophenyl)-1H-pyrazol-5-yl]methyl)-1-ethyl-1H-benzimidazol-5-yl)ethanone O-ethyloxime	Syn and anti 1H NMR (CDCl3): 7.95 (d, 1H+H1), 7.73 (dd, 1H+H1), 7.63 (d, 1H+H1), 7.44 (m, 1H+H1), 7.24-7.33 (m, 3H+H1), [6.18 (d, 1H+H1), 7.13 (d, 1H+H1)], 4.33 (s, 2H), [4.26 (q, 2H), 4.13 (q, 2H1)], 3.98 (q, 2H+2H1), [2.31 (s, 3H), 2.27 (s, 3H1)], [1.35 (t, 3H), 1.28 (t, 3H1)], 1.22 (t, 3H+H1)]	









TABLE 7				
Cmp. #	STRUCTURE	IUPAC NAME	NMR	MS (m/z)
973		(1E)-1-(2-([1-(3-chlorophenyl)-1H-pyrazol-5-yl]methyl)-1-ethyl-1H-benzimidazol-5-yl)ethanone O-(tert-butyl)oxime	Syn and anti ¹ H NMR (CDCl ₃): 8.00 (d, 1H+H1), 7.75 (dd, 1H+H1), 7.63 (d, 1H+H1), 7.27 (d, 1H+H1), 7.38-7.40 (m, 3H+H1), 6.20 (d, 1H)[6.18(d, 1H1)], 4.32 (s, 2H+H1), 3.99 (q, 2H+H1), 2.28 (s, 3H), [2.27(s, 3H1)], 1.37 (s, 9H), [1.32 (s, 9H1)], 1.22 (t, 3H, t3H)	
974		(1E)-1-(2-([1-(3-chlorophenyl)-1H-pyrazol-5-yl]methyl)-1-ethyl-1H-benzimidazol-5-yl)ethanone O-tetrahydro-2H-pyran-2-yloxime	¹ H NMR (CDCl ₃): 7.95 (d, 1H), 7.79 (dd, 1H), 7.63 (d, 1H), 7.52 (m, 1H), 7.37-7.40 (m, 3H), 7.26 (dd, 1H), 6.19 (d, 1H), 5.43 (m, 1H), 4.32 (s, 2H), 3.85-4.02 (m, 4H), 3.66 (m, 2H), 1.62-1.92 (m, 4H), 1.22 (t, 3H)	
975		(1-ethyl-2-([1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl)-1H-benzimidazol-5-yl)acetonitrile	7.66 (m, 1H), 7.63 (d, 1H), 7.24-7.49 (m, 5H), 7.13 (m, 1H), 6.17 (d, 1H), 4.34 (s, 2H), 3.99 (q, 2H), 3.87 (s, 2H), 1.22 (t, 3H)	
976		1-ethyl-2-([1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl)-5-(1,3,4-oxadiazol-2-yl)-1H-benzimidazole	¹ H NMR (d ₆ DMSO): 8.17(1H, s), 8.04-7.98(2H, m), 7.71(1H, d), 7.55-7.50(2H, m), 7.45-7.43(1H, m), 7.29-7.24(1H, m), 6.40(1H, d), 4.77(2H, s), 4.36(2H, q), 2.58(3H, s), 1.24(3H, t)	403.6 [M+1]
977		1-ethyl-2-([1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl)-5-(5-methyl-1,3,4-oxadiazol-2-yl)-1H-benzimidazole	¹ H NMR (CD ₃ OD): 8.79 (s, 1H), 8.31 (d, 1H), 7.69 (s, 1H), 7.60 (d, 1H), 7.46 (dd, 1H), 7.30 (d, 2H), 7.16-7.20 (m, 1H), 6.37 (s, 1H), 4.55 (s, 2H), 4.20 (q, 2H), 1.24 (t, 3H)	
978		1-ethyl-2-([1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl)-1H-imidazo[4,5-c]pyridine	¹ H NMR (CD ₃ OD): 8.79 (s, 1H), 8.31 (d, 1H), 7.69 (s, 1H), 7.60 (d, 1H), 7.46 (dd, 1H), 7.30 (d, 2H), 7.16-7.20 (m, 1H), 6.37 (s, 1H), 4.55 (s, 2H), 4.20 (q, 2H), 1.24 (t, 3H)	
979		1-ethyl-2-([1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl)-1H-benzimidazole-6-carbonitrile		
980		1-(1-ethyl-2-([1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl)-1H-benzimidazol-5-yl)propan-1-one		

TABLE 7				
Cmp. #	STRUCTURE	IUPAC NAME	NMR	MS (m/z)
981		2-[[1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl]-1-(2-methoxyethyl)-1H-benzimidazole-5-carbonitrile		
982		2-[[1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl]-1-(3-methoxypropyl)-1H-benzimidazole-5-carbonitrile		390.1 [M+1]; 388.3 [M-1]
983		1-(2-cyanoethyl)-2-[[1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl]-1H-benzimidazole-5-carbonitrile		371.3 [M+1]
984		1-ethyl-5-(5-ethyl-1,3,4-oxadiazol-2-yl)-2-[[1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl]-1H-benzimidazole	1 H NMR (d6DMSO): 8.17(1H, s), 8.01-7.94(2H, m), 7.71(1H, s), 7.53-7.51(2H, m), 7.44-7.42(1H, m), 7.28-7.24(1H, m), 6.39(1H, s), 4.73(2H, s), 4.33(2H, q), 2.94(2H, q), 1.32(3H, t), 1.23(3H, t)	417.5 [M+1]
985		5-(1-ethyl-2-[[1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl]-1H-benzimidazol-5-yl)-1,3,4-oxadiazol-2-amine	1 H NMR (d6DMSO): 8.17(1H, s), 8.01-7.94(2H, m), 7.71(1H, s), 7.53-7.51(2H, m), 7.44-7.42(1H, m), 7.28-7.24(1H, m), 6.39(1H, s), 4.73(2H, s), 4.33(2H, q), 2.94(2H, q), 1.32(3H, t), 1.23(3H, t)	417.5 [M+1]
986		2-[[1-(2,5-difluorophenyl)-1H-pyrazol-5-yl](hydroxy)methyl]-1-ethyl-1H-benzimidazole-5-carbonitrile	1H NMR (CDCl3): 7.93 (d, 1H), 7.67 (d, 1H), 7.55 (dd, 1H), 7.40 (dd, 1H); 7.24 (m, 1H), 7.10-7.15 (m, 2H), 6.20 (d, 1H), 6.01 (s, 1H), 4.98 (br, S, 1H); 4.04 (q, 2H), 1.21 (t, 3H)	
987		1-ethyl-2-[[1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl]-1H-imidazo[4,5-b]pyridine		322.3 [M+1]; 320.3 [M-1]
988		2-[[1-(2,5-difluorophenyl)-1H-pyrazol-5-yl](fluoro)methyl]-1-ethyl-1H-benzimidazole-5-carbonitrile	1H NMR (CDCl3): 8.07 (d, 1H), 7.77 (d, 1H), 7.60 (dd, 1H), 7.47 (dd, 1H), 7.11-7.23 (m, 3H), 6.85 [d, 1H, J(H,F)=47.4], 6.50 (t, 1H), 4.27 (m, 2H), 1.36 (t, 3H)	

TABLE 7				
Cmp. #	STRUCTURE	IUPAC NAME	NMR	MS (m/z)
989		1-ethyl-2-([1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl)-5-(5-pyridin-3-yl-1,3,4-oxadiazol-2-yl)-1H-benzimidazole	¹ H NMR (d ₆ DMSO): 9.41(1H, s), 8.88(1H, d), 8.67(1H, d), 8.46(1H, s), 8.25(1H, d), 8.10(1H, d), 7.81-7.72(2H, m), 7.55-7.46(3H, m), 7.30-7.26(1H, m), 6.43(1H, s), 4.84(2H, s), 4.40(2H, q), 1.27(3H, t)	
990		1-ethyl-2-([1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl)-5-isoxazol-5-yl-1H-benzimidazole	¹ H NMR (CDCl ₃): 8.30 (d, 1H), 8.16 (dd, 1H), 7.76 (dd, 1H), 7.64 (d, 1H), 7.38-7.50 (m, 2H), 7.25-7.32 (m, 2H), 7.14 (m, 1H), 6.52 (d, 1H), 6.21 (d, 1H), 4.37 (s, 2H), 4.01 (q, 2H), 1.25 (t, 3H)	
991		1-ethyl-2-([1-(3-fluorophenyl)-1H-pyrazol-5-yl](hydroxy)methyl)-1H-benzimidazole-5-carbonitrile	¹ H NMR (CDCl ₃): 8.02 (dd, 1H), 7.47-7.62 (m 5H), 7.39 (d, 1H), 7.18 (m, 1H), 6.12 (d, 1H), 6.00 (s, 1H), 5.05 (brs, 1H), 3.83 (q, 2H), 1.08 (t, 3H)	
992		1-ethyl-2-(fluoro[1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl)-1H-benzimidazole-5-carbonitrile	¹ H NMR (CDCl ₃): 8.12 (dd, 1H), 7.73 (d, 1H), 7.62 (dd, 1H), 7.44-7.53 (m, 2H), 7.31-7.39 (m, 2H), 7.17 (m, 1H), 6.83 (d, 1H, J _{H,F} =46.8 MZ), 4.39 (m, 2H), 1.43 (t, 3H)	
993		methyl 5-([1-(3-cyano-1-ethyl-1H-benzimidazol-2-yl)oxy]-1-(3-fluorophenyl)-1H-pyrazole-4-carboxylate		406.3 [M+1]
994		2-([1-(2,5-difluorophenyl)-1H-pyrazol-5-yl](difluoro)methyl)-1-ethyl-1H-benzimidazole-5-carbonitrile	¹ H NMR (CDCl ₃): 8.09 (m, 1H), 7.84 (M, 1H), 7.64 (dd, 1H), 7.48 (dd, 1H), 6.96-7.18 (m, 3H), 6.85 (d, 1H), 4.33 (q, 2H), 1.38 (t, 3H)	
995		1-ethyl-2-([1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl)-6-(1,3,4-oxadiazol-2-yl)-1H-benzimidazole		389.2 [M+1]
996		1-ethyl-2-([1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl)-6-(5-methyl-1,3,4-oxadiazol-2-yl)-1H-benzimidazole		403.3 [M+1]









TABLE 7				
Cmp. #	STRUCTURE	IUPAC NAME	NMR	MS (m/z)
997		1-(2-((1-(3-fluorophenyl)-1H-pyrazol-5-yl)methyl)-1H-benzimidazol-1-yl)acetone		349.3 [M+1]; 347.2 [M-1]
998		2-(difluoro[1-(3-fluorophenyl)-1H-pyrazol-5-yl)methyl]-1-ethyl-1H-benzimidazole-5-carbonitrile	1H NMR (CDCl3): 8.078 (m, 1H), 7.80 (m, 1H), 7.63 (dd, 1H), 7.48 (dd, 1H), 7.13-7.28 (m, 3H), 6.99 (m, 1H), 6.81 (d, 1H), 4.35 (q, 2H), 1.39 (t, 3H)	
999		2-((1-(3-fluorophenyl)-1H-pyrazol-5-yl)methyl)-1-(3-fluoropropyl)-1H-benzimidazole		353.2 [M+1]; 351.2 [M-1]
1000		1-(2-fluoroethyl)-2-((1-(3-fluorophenyl)-1H-pyrazol-5-yl)methyl)-1H-benzimidazole		339.2 [M+1]
1001		4-(2-((1-(3-fluorophenyl)-1H-pyrazol-5-yl)methyl)-1H-benzimidazol-1-yl)butanenitrile		360.2 [M+1]; 358.3 [M-1]
1002		1-(1-ethyl-2-((1-(3-fluorophenyl)-1H-pyrazol-5-yl)methyl)-1H-benzimidazol-5-yl)-2,2,2-trifluoroethanone		417.2 [M+1]
1003		1-(2-((1-(3-chlorophenyl)-1H-pyrazol-5-yl)methyl)-1-ethyl-1H-benzimidazol-5-yl)-2,2,2-trifluoroethanone		
1004		5-chloro-2-((1-(2,5-difluorophenyl)-1H-pyrazol-5-yl)(difluoro)methyl)-1-ethyl-1H-benzimidazole		

TABLE 7				
Cmp. #	STRUCTURE	IUPAC NAME	NMR	MS (m/z)
1005		(1E)-1-(1-ethyl-2-([1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl)-1H-benzimidazol-5-yl)ethanone O-(2-ethoxyethyl)oxime	¹ H NMR, δppm (CDCl ₃): 7.95(m, 1H), 7.72(dd, 1H), 7.63(d, 1H), 7.45(m, 1H), 7.11-7.31(m, 4H), 6.18(d, 1H), 4.34-4.37(m, 4H), 3.98(q, 2H), 3.77(t, 2H), 3.58(q, 2H), 2.33(s, 3H), 1.18-1.26(m, 6H)	
1006		1-ethyl-2-([1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl)-N,N-dimethyl-1H-benzimidazole-6-carboxamide	¹ H NMR, δppm (CDCl ₃): 7.72(dd, 1H), 7.63(d, 1H), 7.40-7.49 (m, 2H), 7.24-7.34(m, 3H), 7.13(d, 1H), 6.18(d, 1H), 4.35(s, 2H), 3.99(q, 2H), 3.13(s, 2H), 3.07(s, 3H), 1.23(t, 3H)	
1007		3-ethyl-2-([1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl)-3H-imidazo[4,5-b]pyridine		
1008		2-([1-(2,5-difluorophenyl)-1H-pyrazol-5-yl]methyl)-3-ethyl-3H-imidazo[4,5-b]pyridine	¹ H NMR (CDCl ₃): 8.19 (1H, d), 7.61 (1H, d), 7.48-7.44(1H,m), 7.28-7.13(4H, m), 6.14(1H, m), 4.44(2H, s), 3.93(2H, q), 1.17(3H, t),	356.2 [M+1]
1009		4-chloro-1-ethyl-2-([1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl)-1H-imidazo[4,5-c]pyridine	¹ H nmr (CDCl ₃): 8.19(1H, d), 7.61(1H, d), 7.48-7.44(1H, m), 7.28-7.13(4H, m), 6.14(1H, m), 4.44(2H, s), 3.93(2H, q), 1.17(3H, t)	356.2 [M+1]
1010		3-ethyl-2-([1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl)-5-methoxy-3H-imidazo[4,5-b]pyridine		392.3 [M+1]
1011		2-([1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl)-3-isopropyl-3H-imidazo[4,5-b]pyridine		
1012		1-(cyclopropylmethyl)-2-([1-(2,5-difluorophenyl)-1H-pyrazol-5-yl]methyl)-1H-imidazo[4,5-b]pyridine	H-1 NMR (CDCl ₃): 0.22 (m, 2H), 0.54(m, 2H), 0.95(m, 1H), 3.84(d, 2H), 4.32(s, 2H), 6.18(s, 2H), 7.13-7.25 (m, 4H), 7.65 (m, 2H), 8.52 (d, 1H)	366 [M+1]
1013		1-(cyclopropylmethyl)-2-([1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl)-1H-imidazo[4,5-b]pyridine	H-1 NMR (CDCl ₃): 0.22 (m, 2H), 0.54(m, 2H), 0.95(m, 1H), 3.84(d, 2H), 4.32(s, 2H), 6.18(s, 2H), 7.13-7.25 (m, 4H), 7.65 (m, 2H), 8.52 (d, 1H)	366 [M+1]

TABLE 7				
Cmp. #	STRUCTURE	IUPAC NAME	NMR	MS (m/z)
1014		2-([1-(2,5-difluorophenyl)-1H-pyrazol-5-yl]methyl)-1-propyl-1H-imidazo[4,5-b]pyridine	H-1 NMR (CDCl ₃): 0.82(t, 3H), 1.62(h, 2H), 3.85(t, 2H), 4.33 (s, 2H), 6.20 (s, 1H), 7.05-7.25(m, 4H), 7.82(m, 2H), 8.53(d, 1H)	354 [M+1]
1015		2-([1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl)-3-propyl-3H-imidazo[4,5-b]pyridine	(CDCl ₃) 8.43(dd, J = 4.8, 1.5 Hz, 1H), 8.10 (dd, J = 8.0, 1.4 Hz, 1H), 7.64 (d, J = 2.1 Hz, 1H), 7.47 (m, 1H), 7.23-7.34 (m, 3H), 7.12 (m, 1H), 6.31 (d, J = 1.8 Hz, 1H), 4.60 (s, 2H), 4.03 (t, J = 7.7 Hz, 2H), 1.65 (m, 2H), 0.84 (t, J = 7.5 Hz, 3H)	m/z 366.2 [M+1]
1016		2-([1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl)-5-methyl-3-propyl-3H-imidazo[4,5-b]pyridine	(CDCl ₃) 7.85 (d, J = 7.8 Hz, 1H), 7.64 (d, J = 1.5 Hz, 1H), 7.43 (m, 1H), 7.24-7.29 (m, 2H), 7.12 (m, 1H), 7.06 (d, J = 8.1 Hz, 1H), 6.22 (d, J = 1.5 Hz, 1H), 4.33 (s, 2H), 4.01 (t, J = 7.5 Hz, 2H), 2.64 (s, 3H), 1.67 (m, 2H), 0.84 (t, J = 7.3 Hz, 3H)	m/z 350.3 [M+1]
1017		3-ethyl-2-([1-(3-(trifluoromethyl)phenyl)-1H-pyrazol-5-yl]methyl)-3H-imidazo[4,5-b]pyridine	(CDCl ₃) 8.36 (dd, J = 4.8, 1.3 Hz, 1H), 7.98 (dd, J = 8.0, 1.3 Hz, 1H), 7.78 (m, 1H), 7.56-7.72 (m, 4H), 7.22 (dd, J = 8.1, 4.8 Hz, 1H), 6.29 (d, J = 1.5 Hz, 1H), 4.36 (s, 2H), 4.17 (q, J = 7.2 Hz, 2H), 1.27 (t, J = 7.2 Hz, 3H)	m/z 372.2 [M+1]
1018		3-ethyl-2-([1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl)-5-methyl-3H-imidazo[4,5-b]pyridine	(CDCl ₃) 7.85 (d, J = 8.1 Hz, 1H), 7.64 (d, J = 1.8 Hz, 1H), 7.44 (m, 1H), 7.24-7.29 (m, 2H), 7.13 (m, 1H), 7.06 (d, J = 8.1 Hz, 1H), 6.21 (d, J = 1.5 Hz, 1H), 4.33 (s, 2H), 4.12 (q, J = 7.3 Hz, 2H), 2.65 (s, 3H), 1.25 (t, J = 7.3 Hz, 3H)	m/z 336.2 [M+1]
1019		3-ethyl-2-([1-(6-fluoropyridin-2-yl)-1H-pyrazol-5-yl]methyl)-5-methyl-3H-imidazo[4,5-b]pyridine		m/z 337.2 [M+1]
1020		8-([1-(6-fluoropyridin-2-yl)-1H-pyrazol-5-yl]methyl)-2,9-dimethyl-9H-purine	(CDCl ₃) 8.81 (s, 1H), 7.87 (m, 2H), 7.69 (d, 1H), 6.67 (dd, 1H), 6.33 (s, 1H), 4.86 (s, 2H), 3.90 (s, 3H), 2.79 (s, 3H)	
1021		2-([1-(6-fluoropyridin-2-yl)-1H-pyrazol-5-yl]methyl)-1,6-dimethyl-1H-imidazo[4,5-c]pyridine	(CDCl ₃) 8.82 (s, 1H), 7.88 (m, 1H), 7.84 (m, 1H), 7.67 (s, 1H), 7.11 (s, 1H), 6.69 (m, 1H), 6.27 (s, 1H), 4.85 (s, 2H), 3.81 (s, 3H), 2.66 (s, 3H)	m/z 323 [M+1]

TABLE 7				
Cmp. #	STRUCTURE	IUPAC NAME	NMR	MS (m/z)
1022		8-[[1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl]-2,9-dimethyl-9H-purine	(CDCl ₃) 8.90 (s, 1H), 7.63(d, 1H), 7.42 (m, 1H), 7.26 (m, 2H), 7.12 (m, 1H), 6.16 (d, 1H), 4.32 (s, 2H), 3.60 (s, 3H), 2.78 (s, 3H)	
1023		2-[[1-(3-fluorophenyl)-1H-pyrazol-5-yl]methyl]-1-methyl-6-(trifluoromethyl)-1H-benzimidazole	(CDCl ₃) 7.79 (d, 1H), 7.59 (d, 2H), 7.54 (m, 2H), 7.30 (m, 2H), 7.16 (m, 1H), 6.12 (d, 1H), 4.38 (s, 2H), 3.61 (s, 3H),	
1024		9-ethyl-8-[[1-(6-fluoropyridin-2-yl)-1H-pyrazol-5-yl]methyl]-2-methyl-9H-purine	(CDCl ₃) 8.82 (s, 1H), 7.88 (m, 1H), 7.83 (q, 1H), 7.69 (d, 1H), 6.69 (m, 1H), 6.35 (s, 1H), 4.95 (s, 2H), 4.44 (q, 2H), 2.82 (s, 3H), 1.52 (t, 3H)	
1025		3-ethyl-2-[[1-(6-fluoropyridin-2-yl)-1H-pyrazol-5-yl]methyl]-3H-imidazo[4,5-c]pyridine	(CDCl ₃) 8.78 (s, 1H), 8.38 (d, J = 3.9 Hz, 1H), 7.83-7.89 (m, 2H), 7.66 (s, 1H), 7.56 (d, J = 3.9 Hz, 1H), 7.25 (d, J = 9.9 Hz, 1H), 6.70 (d, J = 9.9 Hz, 1H), 4.81 (s, 2H), 4.39 (q, J = 5.4 Hz, 2H), 1.49 (t, J = 5.4 Hz, 3H)	m/z 323 [M+1]
1026		1-ethyl-2-[[1-(3-fluoropyridin-2-yl)-1H-pyrazol-5-yl]methyl]-6-methyl-1H-imidazo[4,5-c]pyridine	(CDCl ₃) 8.82 (s, 1H), 8.28 (br d, J = 4.0 Hz, 1H), 7.70 (d, J = 1.6 Hz, 1H), 7.61 (m, 1H), 7.35 (m, 1H), 7.07 (s, 1H), 6.23 (s, 1H), 4.56 (s, 2H), 4.09 (q, J = 7.2 Hz, 2H), 2.65 (s, 3H), 1.25 (t, J = 7.2 Hz, 3H)	m/z 337 [M+1]
1027		2-[[1-(6-fluoropyridin-2-yl)-1H-pyrazol-5-yl]methyl]-3-methyl-3H-imidazo[4,5-c]pyridine	(CDCl ₃) 8.78 (br s, 1H), 8.38 (br s, 1H), 7.87 (m, 2H), 7.68 (s, 1H), 7.55 (br s, 1H), 7.25 (s, 1H), 6.69 (br d, J = 7.5 Hz, 1H), 6.30 (s, 2H), 4.88 (s, 3H)	m/z 309.2 [M+1]
1028		6-ethyl-2-[[1-(3-fluoropyridin-2-yl)-1H-pyrazol-5-yl]methyl]-1-methyl-1H-imidazo[4,5-c]pyridine	(CDCl ₃) 8.86 (s, 1H), 8.30 (d, J = 4.4 Hz, 1H), 7.71 (s, 1H), 7.62 (m, 1H), 7.36 (m, 1H), 7.07 (s, 1H), 6.18 (s, 1H), 4.58 (s, 2H), 3.64 (s, 3H), 2.94 (q, J = 7.6 Hz, 2H), 1.36 (t, J = 7.6 Hz, 3H)	m/z 337 [M+1]
1029		2-[[1-(6-fluoropyridin-2-yl)-1H-pyrazol-5-yl]methyl]-3,5-dimethyl-3H-imidazo[4,5-b]pyridine	(CDCl ₃) 8.86 (s, 1H), 8.30 (d, J = 4.4 Hz, 1H), 7.71 (s, 1H), 7.62 (m, 1H), 7.36 (m, 1H), 7.07 (s, 1H), 6.18 (s, 1H), 4.58 (s, 2H), 3.64 (s, 3H), 2.94 (q, J = 7.6 Hz, 2H), 1.36 (t, J = 7.6 Hz, 3H)	m/z 337 [M+1]

TABLE 7				
Cmp. #	STRUCTURE	IUPAC NAME	NMR	MS (m/z)
1030		1-ethyl-2-([1-(1,3-thiazol-2-yl)-1H-pyrazol-5-yl]methyl)-1H-imidazo[4,5-c]pyridine	(CDCl ₃) 9.00 (s, 1H), 8.43 (d, 1H), 7.63 (d, 1H), 7.43 (d, 1H), 7.15 (d, 1H), 7.06 (d, 1H), 6.25 (d, 1H), 4.99 (s, 2H), 4.25 (q, 2H), 1.36 (t, 3H)	
1031		1-ethyl-2-([1-(1,3-thiazol-2-yl)-1H-pyrazol-5-yl]ethyl)-1H-imidazo[4,5-c]pyridine	(CDCl ₃) 9.05 (s, 1H), 8.39 (d, 1H), 7.59 (d, 1H), 7.49 (d, 1H), 7.25 (d, 1H), 7.09 (d, 1H), 6.28 (d, 1H), 5.85 (q, 1H), 4.19 (q, 2H), 1.85 (d, 3H), 1.29 (t, 3H)	
1032		1-ethyl-2-([1-(1,3-thiazol-2-yl)-1H-pyrazol-5-yl]methyl)-6-(trifluoromethyl)-1H-imidazo[4,5-c]pyridine	(CD ₃ OD) 8.7 (s, 1H), 8.17 (s, 1H), 7.74 (d, J = 3.6 Hz, 1H), 7.33 (d, J = 3.6 Hz, 1H), 7.21 (d, J = 3.6 Hz, 1H), 7.21 (d, J = 3.6 Hz, 1H), 6.53 (s, 1H), 5.01 (s, 2H), 4.54 (q, J = 7.2 Hz, 2H), 1.52 (t, J = 7.2 Hz, 3H)	
1033		3-ethyl-2-([1-(1,3-thiazol-2-yl)-1H-pyrazol-5-yl]methyl)-3H-imidazo[4,5-c]pyridine	(CD ₃ OD) 9.42 (s, 1H), 8.44 (d, J = 6.6 Hz, 1H), 7.9 (d, J = 6.3 Hz, 1H), 7.7 (s, 1H), 7.26 (d, J = 3.6 Hz, 1H), 7.20 (d, J = 3.6 Hz, 1H), 6.63 (s, 1H), 5.09 (s, 2H), 4.68 (q, J = 7.5 Hz, 2H), 1.62 (t, J = 7.5 Hz, 3H)	m/z 311 [M+1]









Cmp. #	STRUCTURE	IUPAC NAME	NMR	MS (m/z)
1042		3-(cyclopropylmethyl)-2-([2-(2,5-difluorophenyl)-4-methyl-1H-imidazol-1-yl]methyl)-3H-imidazo[4,5-b]pyridine	¹ H NMR (CDCl ₃): 8.38 (dd, 1H), 8.04 (dd, 1H), 7.40 (m, 1H), 7.15-7.27 (m, 3H), 6.79 (d, 1H), 5.35 (s, 2H), 3.82 (d, 2H), 2.22 d, 3H), 0.75 (m, 1H), 0.39-0.44 (m, 2H), 0.23-0.28 (m, 2H)	
1043		1-ethyl-2-([2-(3-fluorophenyl)-4-methyl-1H-imidazol-1-yl]methyl)-1H-benzimidazole-5-carbonitrile		m/z 360 [M+1]
1044		1-ethyl-2-([2-(3-fluorophenyl)-5-methyl-1H-imidazol-1-yl]methyl)-1H-benzimidazole-5-carbonitrile		m/z 360 [M+1]
1045		2-([2-(2-methyl-4-phenyl-1,3-thiazol-5-yl)methyl]-1-propyl-1H-imidazo[4,5-c]pyridine	(CDCl ₃) 9.06 (s, 1H), 8.39 (d, J = 5.6 Hz, 1H), 7.61 (m, 2H), 7.47 (m, 2H), 7.41 (m, 1H), 7.20 (d, J = 5.6 Hz, 1H), 4.54 (s, 2H), 3.77 (t, J = 7.6 Hz, 2H), 2.68 (s, 3H).	m/z 349 [M+1]
1046		5-bromo-1-ethyl-2-([3-(3-fluorophenyl)-1H-pyrazol-4-yl]methyl)-1H-benzimidazole		401.0 [M+1]
1047		5-bromo-1-ethyl-2-([5-(3-fluorophenyl)isoxazol-4-yl]methyl)-1H-benzimidazole	(CDCl ₃) 8.40 (s, 1H), 7.88 (s, 1H), 7.31-7.52 (m, 4H), 7.18-7.22 (m, 2H), 4.14 (s, 2H), 4.04 (q, 2H), 1.18 (t, 3H)	
1048		5-bromo-1-ethyl-2-([5-(3-fluorophenyl)-1-methyl-1H-pyrazol-4-yl]methyl)-1H-benzimidazole	(CDCl ₃) 7.81 (s, 1H), 7.38-7.45 (m, 2H), 7.32 (dd, 1H), 7.00-7.18 (m, 4H), 4.00 (s, 2H), 3.90 (q, 2H), 3.78 (s, 3H), 1.15 (t, 3H)	
1049		5-bromo-1-ethyl-2-([3-(3-fluorophenyl)-1-methyl-1H-pyrazol-4-yl]methyl)-1H-benzimidazole	(CDCl ₃) 7.88 (s, 1H), 7.27-7.42 (m, 4H), 7.00-7.20 (m, 3H), 4.21 (s, 2H), 3.95 (t, 2H), 3.83 (s, 3H), 1.15 (t, 3H)	

TABLE 8				
Cmp. #	STRUCTURE	IUPAC NAME	NMR	MS (m/z)
1050		5-bromo-1-ethyl-2-([4-(3-fluorophenyl)-1,3-thiazol-5-yl]methyl)-1H-benzimidazole	(CDCl ₃) 8.78 (s, 1H), 7.91 (s, 1H), 7.38-7.48 (m, 4H), 7.11-7.19 (m, 2H), 4.59 (s, 2H), 3.90 (q, 2H), 1.12 (t, 3H)	
1051		1-ethyl-2-([4-(6-fluoropyridin-2-yl)-1,3-thiazol-5-yl]methyl)-6-methyl-1H-imidazo[4,5-c]pyridine	(CDCl ₃) 8.92 (s, 1H), 8.75 (s, 1H), 8.16 (d, 1H), 7.90 (dd, 1H), 7.08 (s, 1H), 6.87 (d, 1H), 5.28 (s, 2H), 4.32 (q, 2H), 2.63 (s, 3H), 1.32 (t, 3H)	
1052		2-([3,5-bis(3-fluorophenyl)isoxazol-4-yl]methyl)-1-ethyl-6-(trifluoromethyl)-1H-benzimidazole		m/z 484 [M+1]
1053		2-(2,4'-bi-1,3-thiazol-5'-ylmethyl)-1-ethyl-1H-imidazo[4,5-c]pyridine	(CDCl ₃) 9.05 (s, 1H), 8.76 (s, 1H), 8.41 (d, 1H), 7.88 (d, 1H), 7.41 (d, 1H), 7.25 (d, 1H), 5.35 (s, 2H), 4.26 (q, 2H), 1.25 (t, 3H)	
1054		1-ethyl-2-([2-(1,3-thiazol-2-yl)-4-(trifluoromethyl)-1H-imidazol-1-yl]methyl)-1H-imidazo[4,5-c]pyridine	(CDCl ₃) 9.10 (s, 1H), 8.47 (d, J = 6.0 Hz, 1H), 7.88 (d, J = 3.2 Hz, 1H), 7.58 (s, 1H), 7.48 (d, J = 3.2 Hz, 1H), 7.31 (d, J = 5.6 Hz, 1H), 6.37 (s, 2H), 4.31 (q, J = 7.2 Hz, 2H), 1.24 (t, J = 7.2 Hz, 3H)	m/z 379 [M+1]
1055		3-ethyl-6-isopropyl-2-([2-(1,3-thiazol-2-yl)-4-(trifluoromethyl)-1H-imidazol-1-yl]methyl)-3H-imidazo[4,5-c]pyridine	(CDCl ₃) 8.75 (s, 1H), 7.87 (d, J = 3.3 Hz, 1H), 7.59 (d, J = 0.9 Hz, 1H), 7.54 (s, 1H), 7.47 (d, J = 3.6 Hz, 1H), 6.32 (s, 2H), 4.38 (q, J = 7.2 Hz, 2H), 3.21 (sept, J = 6.9 Hz, 1H), 1.35 (d, J = 6.9 Hz, 6H), 1.30 (t, J = 7.2 Hz, 3H)	m/z 421 [M+1]
1056		1-ethyl-6-isopropyl-2-([2-(1,3-thiazol-2-yl)-4-(trifluoromethyl)-1H-imidazol-1-yl]methyl)-1H-imidazo[4,5-c]pyridine	(CDCl ₃) 9.02 (s, 1H), 7.88 (d, J = 2.7 Hz, 1H), 7.55 (d, J = 0.9 Hz, 1H), 7.48 (d, J = 3.3 Hz, 1H), 7.13 (s, 1H), 6.34 (s, 2H), 4.28 (q, J = 7.2 Hz, 2H), 3.20 (sept, J = 6.9 Hz, 1H), 1.36 (d, J = 6.9 Hz, 6H), 1.22 (t, J = 7.2 Hz, 3H)	m/z 421 [M+1]

Example 51

5 Preparation of radiolabeled probe compounds of the invention

10 The compounds of the invention are prepared as radiolabeled probes by carrying out their synthesis using precursors comprising at least one atom that is a radioisotope. The radioisotope is preferably selected from at least one of carbon (preferably ^{14}C), hydrogen (preferably ^3H), sulfur (preferably ^{35}S), or iodine (preferably ^{125}I). Such radiolabeled probes are conveniently synthesized by a radioisotope supplier specializing in custom synthesis of radiolabeled probe compounds. Such suppliers include Amersham Corporation, Arlington Heights, IL; Cambridge Isotope Laboratories, Inc. Andover, MA; SRI International, Menlo Park, CA; Wizard Laboratories, West Sacramento, CA; ChemSyn Laboratories, Lexena, KS; American Radiolabeled Chemicals, Inc., St. Louis, MO; and Moravsek Biochemicals Inc., Brea, CA.

20 Tritium labeled probe compounds are also conveniently prepared catalytically via platinum-catalyzed exchange in tritiated acetic acid, acid-catalyzed exchange in tritiated trifluoroacetic acid, or heterogeneous-catalyzed exchange with tritium gas. Such preparations are also conveniently carried out as a custom radiolabeling by any of the suppliers listed in the preceding paragraph using the compound of the invention as substrate. In addition, certain precursors may be subjected to tritium-halogen exchange with tritium gas, tritium gas reduction of unsaturated bonds, or reduction using sodium borotritide, as
30 appropriate.

Example 52

Receptor autoradiography

Receptor autoradiography (receptor mapping) is carried out in vitro as described by Kuhar in sections 8.1.1 to 8.1.9 of Current Protocols in Pharmacology (1998) John Wiley & Sons, New York, using radiolabeled compounds of the invention prepared as described in the preceding Example.

Example 53

Binding Assay

The high affinity and high selectivity of preferred compounds of the invention for the benzodiazepine site of the GABA_A receptor can be confirmed using the binding assay described by Thomas and Tallman (*J. Bio. Chem.* 1981; 156:9838-9842, and *J. Neurosci.* 1983; 3:433-440).

Rat cortical tissue is dissected and homogenized in 25 volumes (w/v) of Buffer A (0.05 M Tris HCl buffer, pH 7.4 at 4 °C). The tissue homogenate is centrifuged in the cold (4 °C) at 20,000 x g for 20 minutes. The supernatant is decanted, the pellet rehomogenized in the same volume of buffer, and centrifuged again at 20,000 x g. The supernatant of this centrifugation step is decanted. The resulting pellet may be stored at -20 °C overnight. The pellet is then thawed and resuspended in 25 volumes of Buffer A (original wt/vol), centrifuged at 20,000 x g and the supernatant decanted. This wash step is repeated once. The pellet is finally resuspended in 50 volumes of Buffer A.

Incubations contain 100 µl of tissue homogenate, 100 µl of radioligand, (0.5 nM ³H-Ro15-1788 [³H-Flumazenil], specific activity 80 Ci/mmol), and test compound or control (see below), and are brought to a total volume of 500 µl with Buffer A. Incubations are carried for 30 minutes at 4°C and then rapidly filtered through Whatman GFB filters to separate free and bound

ligand. Filters are washed twice with fresh Buffer A and counted in a liquid scintillation counter. Nonspecific binding (control) is determined by displacement of ^3H Ro15-1788 with 10 μM Diazepam (Research Biochemicals International, Natick, MA). Data are collected in triplicate, averaged, and percent inhibition of total specific binding (Total Specific Binding = Total - Nonspecific) is calculated for each compound.

A competition binding curve may be obtained with up to 11 points spanning the compound concentration range from 10^{-12}M to 10^{-5}M obtained per curve by the method described above for determining percent inhibition. K_i values are calculated according to the Cheng-Prusoff equation. Each of the compounds disclosed in Tables 6-8 was tested in this fashion and each was found to have a K_i of $< 4\mu\text{M}$. Preferred compounds of the invention exhibit K_i values of less than 100 nM and more preferred compounds of the invention exhibit K_i values of less than 10 nM.

Example 54

Electrophysiology

The following assay can be used to determine if a compound of the invention acts as an agonist, an antagonist, or an inverse agonist at the benzodiazepine site of the GABA_A receptor.

Assays are carried out as described in White and Gurley (NeuroReport 6: 1313-1316, 1995) and White, Gurley, Hartnett, Stirling, and Gregory (Receptors and Channels 3: 1-5, 1995) with modifications. Electrophysiological recordings are carried out using the two electrode voltage-clamp technique at a membrane holding potential of -70 mV. *Xenopus Laevis* oocytes are enzymatically isolated and injected with non-polyadenylated cRNA mixed in a ratio of 4:1:4 for α , β and γ subunits, respectively. Of the nine combinations of α , β and γ subunits described in the White et al. publications, preferred combinations are $\alpha_1\beta_2\gamma_2$,

These net values are used for the calculation of average efficacy and EC_{50} values by standard methods. To evaluate average efficacy and EC_{50} values, the concentration/effect data are averaged across cells and fit to the logistic equation.

5

It is to be understood that the foregoing describes preferred embodiments of the present invention and that modifications may be made therein without departing from the scope of the present invention as set forth in the following claims.

10

10038069 122101